Interntial statistics


Covariance and correlation

Covanance -

9ge hight
5 80cm
10 100cm
15 120cm

X1 X2 X3 X4 X5 X6

XT YT | XI YI = Positive cova. XI YT | XT YI = Negative cova. XT Y- | XI Y- = Zero Cova.



A co-relation—
Relation as well as streamyth.

$$(ox(X,y) = \frac{(ov(X,y))}{\sigma_X \sigma_Y}$$

Range co-reletion = -1 to · +1

regulive con = -1

Type correlation

Dearson's correlation coefficient

2) spearman's Rank correlation

coefficient

Pearson's -It describe the linear relation blw quantitative/Number variable.

 $P(x,y) = \frac{(6v(x,y))}{\sigma_{x}\sigma_{y}}$

Disadvantage - It doesn't work on non-linear selation variable +089 Spealman's Rank com coefficient formula $Y(X,Y) = \frac{Cov(X,Y)}{Rox Roy}$

X	- 1	XL	\times_3	Xy	~		
	O	(0	(0)	100			
	1		(_		
					_		
Ŕ	7			3000			
					_		
	, Oj	IM	1000	1000			
					-		
•	1	2.1-	= 0.0	0001			
	7		_ 0	.0005	- 7		
	16	VOD	_			0-1	
	·				4		
		500	-)	0.5			
	\ (/	000					
	_	66					
	7	360		6.05	40		
		50	_	0.0	05		
		100	U -				
		1000)()	= (
		107	70 O	•			

Center limit theorem




No of sample =
$$40$$

[Sample Size >= 30 {
 $(n_1, n_2, n_3 - - - - x_{40})$
 $(\overline{x}_1, \overline{x}_1, \overline{x}_3 - - - \overline{x}_{40})$
Doutaset $(x_1, x_2, x_3 - - \overline{x}_{40})$

max no. of sample will head to normalize the data curve.

Population = 10000 N = 30 - 30 - 30 - 30 $\overline{X} = 12 - 15 - 16 - 11 - \cdots$



M = 1 F = 0

,			1	1	
	Cendr	Movies tal	G.	M.	
	M	S	1	0	
	Ļ	M	0	1	
	F	M	0	1	
	M	ξ	1	0	
	M	M	1	1	