

Parsley Grow Guide

For curly parsley, flat-leaf/Italian parsley, seedlings, containers, kitchen gardens, and small farm herb production.

Quick Rescue - Quick Numbers - Sunlight/Soil/Water - Feeding - Harvesting - Problems - Transplanting

Crop	Parsley / <i>Petroselinum crispum</i>
Common types	Curly parsley, flat-leaf parsley / Italian parsley, and root parsley.
Important note	Parsley is a biennial. It mainly grows leaves in year 1, then sends a flower stalk and sets seed in year 2.

ParaLife Trees

Planting a Life in Paradise

1. Quick Rescue Page: If You See This, Do This

What You See	Most Likely Cause	What To Do First
Seedling wilts after planting	Transplant shock, sun stress, dry roots	Shade 1-2 days, water gently, and keep soil evenly moist. Do not flood the roots.
Leaves yellow from bottom	Overwatering, old leaves, low nitrogen	Check drainage first. If soil is not soggy, feed lightly and remove yellow leaves.
Leaves turn pale green	Low nutrition, low light, or root stress	Improve light, add compost, and use a mild feed after the plant settles.
Brown crispy leaf edges	Heat stress, dry soil, fertilizer burn	Water evenly, give afternoon shade in hot weather, and avoid strong fertilizer.
Plant grows tall center stalk	Second-year flowering / bolting	Harvest leaves now. Flavor usually drops after flowering begins. Start new plants for best leaves.
Leaves have chewing holes	Caterpillars / parsleyworms / armyworms	Check early morning/evening and handpick. Use netting if pressure is high.
Tiny insects on new growth	Aphids, thrips, or leafhoppers	Spray with water first. Use mild soap spray every 3-5 days if needed.
White winding lines in leaves	Leafminers	Remove affected leaves. Keep weeds down. Use insect netting for new crops.
White powder on leaves	Powdery mildew	Improve airflow and spacing. Remove badly affected leaves. Avoid wet foliage late evening.

For any serious problem, send photos of the full plant, leaf close-up, leaf underside, stem base, soil/pot, and any insects to WhatsApp: 1 (876) 372-7415.

2. Quick Numbers Page

Item	Specific Range / Target
Botanical name	Petroselinum crispum
Crop type	Biennial herb, usually grown for leaves as an annual
Common edible types	Curly leaf, flat-leaf/Italian, and root parsley
Seed germination	14-28 days; parsley is slow to sprout
Seedling transplant age	5-8 weeks old, once strong enough to handle
Hardening-off period	5-7 days before full outdoor planting
Plant height	About 12-18 inches for leaf parsley; flower stalks can grow taller in year 2
Plant width	About 10-18 inches depending on variety and harvest style
Plant spacing for full plants	8-12 inches apart
Row spacing for small farmers	12-18 inches for hand harvest and airflow
Container depth	8-10 inches minimum; 10-12 inches better for steady growth
Sunlight	4-6+ hours daily; in hot conditions, morning sun plus afternoon shade is useful
Soil pH	6.0-7.0 preferred
First useful harvest after transplant	About 30-45 days after transplant if seedlings are established
From seed to mature harvest	About 70-90 days, depending on variety and conditions

Flat-leaf parsley is usually stronger-flavored and easier to chop. Curly parsley is often used for garnish and bunching. Both grow with the same basic care.

3. Parsley Type Identification

Type	How To Identify It	Use / Difference
Curly parsley	Tightly curled, frilly leaves; compact look	Good for garnish, bunching, and fresh use. Holds shape well.
Flat-leaf / Italian parsley	Flat, deeply cut leaves; looks similar to cilantro but usually thicker and less delicate	Often stronger flavor; good for cooking, sauces, seasoning, and chopping.
Root parsley	Leaves look parsley-like, but plant is grown for a thick white root	Less common for herb pots; needs deeper soil and more time.
Cilantro comparison	Cilantro leaves are usually softer and more rounded/lacy; smell is very different	Do not label cilantro as parsley. Smell the leaves if unsure.

For customer guides, title it "Parsley Grow Guide" and include both curly and flat-leaf parsley unless you are selling one named type only.

4. Sunlight, Soil, and Water

Need	Target / Customer Instruction
Sunlight	Give 4-6+ hours daily. In hot Jamaican conditions, morning sun and light afternoon shade help keep leaves tender.
Best location	Bright area with airflow. Avoid crowded, damp corners.
Soil	Loose, rich, well-draining soil with compost. Parsley prefers steady moisture but not swampy roots.
Soil pH	Aim for 6.0-7.0 when possible.
Watering	Keep evenly moist. Water when the top 1 inch of soil begins to dry.
Containers	Use drainage holes. Do not let pots sit in standing water.
Mulch	Use light mulch in hot beds. Keep mulch slightly away from the crown.

Parsley grows best when it does not repeatedly wilt. The goal is steady moisture, not constant wetness.

5. Feeding Guide

Stage	Timing	Feeding Direction
At planting	Day 0	Mix compost into the soil. Avoid fresh manure or strong fertilizer touching roots.
After settling	7-10 days after planting	If leaves are pale, apply a light liquid feed or mild balanced fertilizer.
Leaf production	Every 3-4 weeks if needed	Feed lightly for steady leaf growth. Do not overfeed.
Container plants	Every 2-4 weeks if growth slows	Containers lose nutrients faster; use gentle feeding and steady watering.
Before harvest	Final 3-5 days	Focus on clean water and healthy leaves, not heavy fertilizer.

Parsley is grown for leaves, so it needs enough nitrogen for green growth, but too much fertilizer can make weak soft growth and reduce quality.

6. Harvesting Guide

Harvest Type	When / How
First light harvest	When plants have several strong outer stems; often 30-45 days after transplant.
Regular harvest	Cut outer stalks close to the base. Leave the center growing point to keep producing.
Harvest frequency	Every 7-14 days once plant is established and growing strongly.
How much to remove	Do not remove more than about 1/3 of the plant at one cutting if you want regrowth.
Best time of day	Harvest early morning after leaves dry, or late evening after heat drops.
Fresh handling	Keep parsley shaded and cool after cutting. Do not leave bunches in hot sun.
End of plant cycle	When a tall flower stalk appears, leaf quality usually declines. Replace with new seedlings.

Best customer instruction: cut whole outer stems near the base, not just the leaf tips. This encourages cleaner regrowth from the center.

7. Leaf Color and Growth Problem Guide

Leaf / Growth Sign	Likely Cause	What To Do
Deep green, firm leaves	Healthy growth	Continue steady watering and regular outer-stem harvest.
Pale green leaves	Low nutrition, low light, or root stress	Improve light and feed lightly after checking drainage.
Yellow lower leaves	Age, wet soil, or low nitrogen	Remove yellow leaves. Check roots and reduce watering if soil is wet.
Brown crispy edges	Heat stress, dry soil, fertilizer burn	Water evenly, add light shade, and avoid strong fertilizer.
Soft collapsing leaves	Too much water, root disease, or heat stress	Check soil moisture and drainage. Remove rotten leaves.
Plant very slow after transplant	Root disturbance or normal slow start	Keep moist, shaded briefly, and wait 7-14 days for recovery.
Tall stalk with smaller leaves	Flowering/bolting	Harvest leaves now and start replacement plants.

8. Flower Stalks, Bolting, and Year 2 Growth

Parsley is a biennial. In the first year it focuses on leaves. In the second year it naturally sends up a tall flower stalk, flowers, makes seed, and then the plant declines.

What You See	Meaning	What To Do
Tall center stalk	Plant is entering flowering stage	Harvest leaves now; quality usually drops after flowering starts.
Small yellow-green flower clusters	Normal second-year flowering	Leave only if you want seed or beneficial insect flowers.
Leaves become smaller/tougher	Energy shifted to flowers/seed	Replace with new plants for fresh bunching.
Early stalk in young plant	Heat, stress, or root restriction	Use shade, steady water, and avoid root-bound seedlings next batch.

9. Small Farmer Notes

Topic	Recommendation
Planting batches	For steady supply, plant or transplant in batches every 3-4 weeks.
Spacing	Use 8-12 inches between plants and 12-18 inches between rows for hand harvest.
Bed setup	Use raised beds where soil holds water. Keep rows weeded for airflow.
Shade strategy	In hot periods, use afternoon shade or 30-40% shade cloth for tender leaves.
Harvest handling	Cut in the cool of the day. Keep bunches shaded, clean, and cool.
Market quality	Remove yellow, dry, insect-damaged, muddy, or diseased stems before bunching.

10. Home Garden Notes

Situation	Recommendation
2-6 plants	Plant 8-12 inches apart in a bright bed or container.
Container growing	Use a pot at least 8-10 inches deep with drainage holes.
Kitchen use	Harvest outer stems often instead of cutting the whole plant.
Hot patio	Give morning sun and shade after midday. Check water daily.
Continuous supply	Start new parsley every few months, because older plants eventually flower and decline.

11. Clean Handling and Seedling Sanitation

Practice	Why It Matters
Wash hands before handling seedlings	Reduces transfer of pests, disease organisms, and dirty residues.
Clean scissors/knife before harvest	Reduces disease spread and keeps fresh herbs cleaner.
Do not handle sick plants then healthy plants	Many plant problems move by hands, tools, splashing water, and insects.
Keep trays off dirty ground	Reduces soil splash, ants, slugs, and root disease risk.
Avoid smoking over seedling trays	Smoke, ash, and dirty hands are poor handling practice for fresh herbs.
Remove heavily diseased plants	Protects the rest of the crop and reduces pest/disease buildup.

Parsley is often eaten fresh, so clean handling matters: clean water, clean containers, clean tools, and clean hands.

12. Hardening Off Seedlings

Hardening off prepares parsley seedlings for direct sun, wind, and outdoor heat. This is useful for tray seedlings and small pots before they are planted out.

Day	Exposure
Day 1-2	Bright shade for 2-3 hours, then back to protected area.
Day 3-4	Morning sun for 2-4 hours, then bright shade.
Day 5-6	Longer morning sun plus bright afternoon shade. Keep cells moist.
Day 7	Ready to plant in early morning or late evening if seedling is not wilted.

Do not let tray cells dry out during hardening off. Small parsley seedlings recover slowly after severe wilt.

13. Transplanting From Trays or Small Pots

Step	Instruction
1	Water seedlings 1-2 hours before transplanting.
2	Prepare holes before removing seedlings from the tray/pot.
3	Remove the seedling gently from the bottom of the cell. Do not pull by the leaves.
4	Keep the root ball together as much as possible. Parsley can be slow after root disturbance.
5	Plant at the same soil depth as the tray/pot. Do not bury the crown deeply.
6	Press soil gently around the root ball and water immediately.
7	Shade lightly for 1-2 days if the weather is hot.
8	Check morning and evening for the first 3-5 days.

Best transplant timing: early morning before 8:30 a.m. or late evening after 4:30 p.m. Avoid midday transplanting.

14. Common Pests

Pest	Signs	Action
Aphids	Tiny soft insects on new growth; curling; sticky leaves	Spray off with water. Use mild soap spray every 3-5 days if needed. Control ants.
Caterpillars / parsleyworms	Chewed leaves; droppings; visible green/striped worms	Handpick early morning/evening. Use netting if pressure is high.
Leafminers	White winding tunnels inside leaves	Remove affected leaves. Keep weeds down. Use insect netting for new crops.
Thrips	Silvery marks, distorted tips, tiny insects in growing points	Remove damaged tips. Spray lightly if needed. Improve plant health.
Leafhoppers	Small jumping insects; yellowing/stunting; disease spread risk	Keep weeds down. Use netting for young crops where pressure is high.
Spider mites	Fine webbing, speckled dry leaves	Rinse undersides, avoid drought stress, repeat treatment every 3-5 days.

15. Common Diseases

Disease / Problem	Signs	Action
Damping off	Seedling collapses near soil line	Avoid overwatering. Improve airflow. Use clean media. Remove collapsed plants.
Root rot	Wilting with wet soil; yellowing; weak roots	Improve drainage. Reduce watering. Repot if container stays soggy.
Powdery mildew	White powdery coating on leaves	Increase airflow and spacing. Remove bad leaves. Avoid wet foliage late.
Leaf spots	Brown/black spots, often worse with wet leaves	Remove spotted leaves. Water soil, not leaves. Avoid crowding.
Crown rot	Soft rot at base/crown; plant collapses	Keep crown dry, improve drainage, remove badly affected plants.
Aster yellows / viral-like disorders	Yellowing, stunting, distorted growth or abnormal flowers	Remove badly affected plants. Control leafhoppers/aphids and weeds.

Prevention matters more than rescue: clean trays, good drainage, enough space, dry foliage, and early pest control are the best disease protection.

16. ParaLife Trees Support Page

Need help with your parsley plant?

Send clear photos of:

Photo Needed	Why
Full plant	Shows size, wilting, spacing, and growth habit.
Leaf close-up	Shows color, chewing, powder, spotting, or curling.
Leaf underside	Many pests hide under leaves.
Stem base and soil/pot	Shows root rot, damping off, drainage, and moisture issues.
Any insect or worm seen	Helps identify the pest and the safest next step.

WhatsApp: 1 (876) 372-7415

ParaLife Trees - Planting a Life in Paradise

Source basis: University of Delaware Cooperative Extension, University of Minnesota Extension, UC ANR Master Gardeners, Purdue Extension, RHS, UF/IFAS, and other extension-style herb production references.