



On-line Kronecker Product Structured Covariance Estimation with Riemannian geometry for t-distributed data

Florent Bouchard¹, Arnaud Breloy², **Ammar Mian**³, Guillaume Ginolhac³


¹: L2S, CentraleSupélec, CNRS, Univ. Paris Saclay

²: LEME, Univ. Paris Nanterre

³: LISTIC, Univ. Savoie Mont-Blanc



@Freepik



Outline

- ➊ Introduction
- ➋ Information geometry and recursive estimation
- ➌ Numerical results
- ➍ Conclusion
- ➎ References

Outline

- 1 **Introduction**
- 2 Information geometry and recursive estimation
- 3 Numerical results
- 4 Conclusion
- 5 References

Kronecker structure and heterogeneous clutter

Kronecker structure of data arises in numerous applications:

- MIMO : [YBO⁺04]
- MEG/EEG data : [dMHW02]
- Space Time Adaptive Processing: [GZH16]
- Synthetic Aperture Radar : [MOAG19]

Moreover, when resolution of data is high (in radar), the data is **heterogeneous** and modeled by heavy-tailed distributions.

The model

The data set $\{\mathbf{x}_i\}_{i=1}^n \in (\mathbb{R}^p)^n$ is assumed to contain independent and identically distributed vectors drawn from the multivariate real Student t -distribution with **unknown** scatter matrix Σ and **known** $d \in \mathbb{N}^*$ degrees of freedom. The model, denoted $\mathbf{x} \sim \mathbb{R}t_d(\mathbf{0}, \Sigma)$, implies that the probability density function of \mathbf{x} is of the form

$$f(\mathbf{x}) \propto |\Sigma|^{-1/2} \left(1 + \frac{\mathbf{x}^T \Sigma^{-1} \mathbf{x}}{d} \right)^{-(d+p)/2}. \quad (1)$$

Kronecker structure

The scatter matrix Σ is assumed to admit a Kronecker product structure, *i.e.*, $\Sigma = A \otimes B$, where $A \in sS_a^{++} = \{M \in S_a^{++} : |M| = 1\}$ and $B \in S_b^{++}$.

Estimation problem

Parameter of interest: $\theta = (A, B)$

Maximum likelihood estimation

$$\hat{\theta} = \underset{\theta \in (s\mathcal{S}_a^{++} \times \mathcal{S}_b^{++})}{\operatorname{argmin}} - \sum_{i=1}^n \log f(\mathbf{x}_i; \theta) \quad (2)$$

- Problem is non-convex in Euclidean sense but iterative algorithms exists [SBP16, MRB⁺21].
- On the other hand, considering the parameter space as the product manifold $\mathcal{M}_{a,b} = s\mathcal{S}_a^{++} \times \mathcal{S}_b^{++}$, it is geodesic-convex.

Problems to tackle

- High-dimensionality of Kronecker products is costly
- Online estimation


Solution: Use a Riemannian recursive framework [ZS19]

Stochastic gradient on manifolds allow to obtain fast and efficient estimation of the parameters. The scheme is as follows:

$$\theta_{i+1} = \exp_{\theta_i}(-t_i \text{grad} \ell_i(\theta_i)), \quad (3)$$

where $\ell_i(\theta) = -\log f(\mathbf{x}_i, \theta)$.


Recursive estimation illustration



Outline

- 1 Introduction
- 2 Information geometry and recursive estimation**
- 3 Numerical results
- 4 Conclusion
- 5 References

What we need



Information geometry

We can use geometry of product manifold without taking into account the statistical model but not as efficient[Ama99].
→ Derivation of Fisher Information metric.

Gradient

We also need the **Riemannian gradient** of the likelihood $\ell_i(\theta)$.

Derivation of metric

Proposition

Given $\theta \in \mathcal{M}_{a,b}$, ξ and $\eta \in T_\theta \mathcal{M}_{a,b}$, the Fisher information metric on $\mathcal{M}_{a,b}$ induced by the likelihood is

$$\langle \xi, \eta \rangle_\theta = \alpha b \operatorname{tr}(\mathbf{A}^{-1} \boldsymbol{\xi}_A \mathbf{A}^{-1} \boldsymbol{\eta}_A) + \alpha a \operatorname{tr}(\mathbf{B}^{-1} \boldsymbol{\xi}_B \mathbf{B}^{-1} \boldsymbol{\eta}_B) + (\alpha - 1) a^2 \operatorname{tr}(\mathbf{B}^{-1} \boldsymbol{\xi}_B) \operatorname{tr}(\mathbf{B}^{-1} \boldsymbol{\eta}_B), \quad (4)$$

where $\alpha = (d+p)/(d+p+1)$.

Proof: Derived from results of [BBG⁺21, Proposition 7] on mappings and Fisher information metric of elliptical distributions in [BGRB19].

Exponential mapping and retraction

the Riemannian exponential mapping at $\theta \in \mathcal{M}_{a,b}$ is defined for $\xi \in T_\theta \mathcal{M}_{a,b}$ as

$$\exp_\theta^{\mathcal{M}_{a,b}}(\xi) = (\mathbf{A} \exp(\mathbf{A}^{-1} \xi_A), \mathbf{B} \exp(\mathbf{B}^{-1} \xi_B)). \quad (5)$$

For better numerical cost and stability, it might be advantageous to prefer a second order approximation:

Retraction

$$R_\theta(\xi) = \left(\mathbf{A} + \xi_A + \frac{1}{2} \xi_A \mathbf{A}^{-1} \xi_A, \mathbf{B} + \xi_B + \frac{1}{2} \xi_B \mathbf{B}^{-1} \xi_B \right). \quad (6)$$

Riemannian gradient

Proposition

The Riemannian gradient of ℓ_i at $\theta \in \mathcal{M}_{a,b}$ according to the Fisher information metric is:

$$\text{grad } \ell_i(\theta) = \left(\frac{1}{\alpha b} P_A(\mathbf{A} \text{sym}(\nabla_A \ell_i(\theta)) \mathbf{A}), \frac{1}{\alpha a} \mathbf{B} \text{sym}(\nabla_B \ell_i(\theta)) \mathbf{B} - \frac{(\alpha - 1) \text{tr}(\mathbf{B} \nabla_B \ell_i(\theta))}{\alpha(\alpha + (\alpha - 1)p)} \mathbf{B} \right), \quad (7)$$

where $\text{sym}(\cdot)$ returns the symmetrical part of its argument; $P_A : \mathcal{S}_a \rightarrow T_A \mathcal{S}_a^{++}$ is the orthogonal projection map such that $P_A(\xi_A) = \xi_A - \frac{\text{tr}(\mathbf{A}^{-1} \xi_A)}{a} \mathbf{A}$; and $\nabla \ell_i(\theta) = (\nabla_A \ell_i(\theta), \nabla_B \ell_i(\theta))$ is the Euclidean gradient of ℓ_i at θ , defined as

$$\nabla_A \ell_i(\theta) = \frac{1}{2} \mathbf{A}^{-1} \left(b \mathbf{A} - \frac{d+p}{d+Q_i(\theta)} \mathbf{M}_i^T \mathbf{B}^{-1} \mathbf{M}_i \right) \mathbf{A}^{-1}, \quad \nabla_B \ell_i(\theta) = \frac{1}{2} \mathbf{B}^{-1} \left(a \mathbf{B} - \frac{d+p}{d+Q_i(\theta)} \mathbf{M}_i \mathbf{A}^{-1} \mathbf{M}_i^T \right) \mathbf{B}^{-1},$$

with \mathbf{M}_i , the $b \times a$ matrix such that $\text{vec}(\mathbf{M}_i) = \mathbf{x}_i$ and $Q_i(\theta) = \text{tr}(\mathbf{A}^{-1} \mathbf{M}_i^T \mathbf{B}^{-1} \mathbf{M}_i)$.

Recursive algorithm

Algorithm 1: Online estimation of θ

Result: Estimate $\theta = (\mathbf{A}, \mathbf{B})$

initialization with $\theta = (\mathbf{A}_0, \mathbf{B}_0)$;

for $i=1,\dots,n$ **do**

$\theta_i = R_{\theta_i} \left(-\frac{1}{i} \text{grad} \ell_i(\theta_i) \right)$

end

Outline

- 1 Introduction
- 2 Information geometry and recursive estimation
- 3 Numerical results**
- 4 Conclusion
- 5 References

Setup of Montecarlo i

100 sets $\{\mathbf{x}_i\}_{i=1}^n$ are drawn from the multivariate Student t -distribution with covariance Σ and $d \in \{3, 100\}$ degrees of freedom, where $n \in \llbracket 1, 500 \rrbracket$. To generate a Σ :

$$\begin{aligned}\Sigma &= \mathbf{A} \otimes \mathbf{B}, \\ \mathbf{A} &= \mathbf{U}_A \Lambda_A \mathbf{U}_A^T, \quad \mathbf{B} = \mathbf{U}_B \Lambda_B \mathbf{U}_B^T,\end{aligned}\tag{8}$$

where $a = b = 4$,

- \mathbf{U}_A and \mathbf{U}_B are random orthogonal matrices,
- Λ_A and Λ_B are diagonal matrices whose minimal and maximal elements are $1/\sqrt{c}$ and \sqrt{c} ($c = 10$ is the condition number with respect to inversion); their other elements are randomly drawn from the uniform distribution between $1/\sqrt{c}$ and \sqrt{c} ; the determinant of Λ_A is then normalized.

Setup of Montecarlo ii

For this experiment, we consider the following estimators:

- the classical **maximum-likelihood estimator** obtained with Riemannian gradient descent (GD). Optimization for this estimator is performed with manopt toolbox [BMAS14].
- the **online version** obtained through stochastic gradient descent (SGD) presented here.

Setup of Montecarlo iii

Both algorithms are initialized with $\theta_0 = (\mathbf{I}_a, \mathbf{I}_b)$.

In order to measure the performance, we consider an error measure for each component \mathbf{A} and \mathbf{B} , which are given by the usual Riemannian distances on ${}_sS_a^{++}$ and S_b^{++}

$$\begin{aligned}\text{err}(\hat{\mathbf{A}}) &= \|\log(\mathbf{A}^{-1/2} \hat{\mathbf{A}} \mathbf{A}^{-1/2})\|_2^2, \\ \text{err}(\hat{\mathbf{B}}) &= \|\log(\mathbf{B}^{-1/2} \hat{\mathbf{B}} \mathbf{B}^{-1/2})\|_2^2.\end{aligned}\tag{9}$$

Comparison between gradient descent and recursive estimation ($d = 3$)




Figure 1: Mean of error measures on A (left) and B (right) of the classical gradient descent method (GD) and its on-line counterpart (SGD) as functions of the number of samples n . $d = 3$.

Comparison between gradient descent and recursive estimation ($d = 100$)




Figure 2: Mean of error measures on A (left) and B (right) of the classical gradient descent method (GD) and its on-line counterpart (SGD) as functions of the number of samples n . $d = 100$.

Outline

- 1 Introduction
- 2 Information geometry and recursive estimation
- 3 Numerical results
- 4 Conclusion**
- 5 References

Conclusion

We have achieved:

- Information geometry on Kronecker products based on Fisher information metric of a Student-t distribution.
- Efficient online scheme for estimation.




Next:

- Extension to all elliptical distributions and deterministic compound-Gaussian distribution.
- Applications in STAP and SAR problems.

Outline

- 1 Introduction
- 2 Information geometry and recursive estimation
- 3 Numerical results
- 4 Conclusion
- 5 References**




References i

-  Shun-ichi Amari, *Natural gradient learning for over-and under-complete bases in ica*, Neural computation **11** (1999), no. 8, 1875–1883.
-  F. Bouchard, A. Breloy, G. Ginolhac, A. Renaux, and F. Pascal, *A Riemannian framework for low-rank structured elliptical models*, IEEE Transactions on Signal Processing **69** (2021), 1185–1199.
-  A. Breloy, G. Ginolhac, A. Renaux, and F. Bouchard, *Intrinsic Cramèr – Rao bounds for scatter and shape matrices estimation in CES distributions*, IEEE Signal Processing Letters **26** (2019), no. 2, 262–266.


References ii

-  N. Boumal, B. Mishra, P.-A. Absil, and R. Sepulchre, *Manopt, a Matlab toolbox for optimization on manifolds*, Journal of Machine Learning Research **15** (2014), 1455–1459.
-  J.C. de Munck, H.M. Huizenga, L.J. Waldorp, and R.A. Heethaar, *Estimating stationary dipoles from meg/eeg data contaminated with spatially and temporally correlated background noise*, IEEE Transactions on Signal Processing **50** (2002), no. 7, 1565–1572.
-  Kristjan Greenewald, Edmund Zelnio, and Alfred Hero, *Robust sar stap via kronecker decomposition*, 2016.
-  A. Mian, J. Ovarlez, A. M. Atto, and G. Ginolhac, IEEE Transactions on Geoscience and Remote Sensing **57** (2019), no. 6, 3919–3932.

References iii

-  Bruno Meriaux, Chengfang Ren, Arnaud Breloy, Mohammed Nabil El Korso, and Philippe Forster, *Efficient estimation of kronecker product of linear structured scatter matrices under t-distribution*, 2020 28th European Signal Processing Conference (EUSIPCO), IEEE, 2021, pp. 2418–2422.
-  Ying Sun, Prabhu Babu, and Daniel P Palomar, *Robust estimation of structured covariance matrix for heavy-tailed elliptical distributions*, IEEE Transactions on Signal Processing **64** (2016), no. 14, 3576–3590.
-  Kai Yu, M. Bengtsson, B. Ottersten, D. McNamara, P. Karlsson, and M. Beach, *Modeling of wide-band mimo radio channels based on nlos indoor measurements*, IEEE Transactions on Vehicular Technology **53** (2004), no. 3, 655–665.

References iv

 Jialun Zhou and Salem Said, *Fast, asymptotically efficient, recursive estimation in a Riemannian manifold*, Entropy **21** (2019), no. 10, 1021.