

Homework 7

Problem 1. Suppose X follows a geometric distribution,

$$P(X = k) = p(1 - p)^{k-1}$$

and assume an i.i.d sample of size n .

(b) Find the mle of p .

(c) Find the asymptotic variance of the mle.

(b) The log likelihood is given by

$$l(p) = \sum_{i=1}^n \log(p(1-p)^{k_i-1}) = \sum_{i=1}^n (\log(p) + (k_i - 1)\log(1-p)) = n \log(p) + \log(1-p) \sum_{i=1}^n (k_i - 1).$$

Then

$$l'(p) = \frac{n}{p} - \frac{1}{1-p} \sum_{i=1}^n (k_i - 1).$$

Setting this equal to 0 gives

$$\hat{p} = n \left(\sum_{i=1}^n k_i \right)^{-1}.$$

This is indeed a maximum because

$$l''(p) = -\frac{n}{p^2} - \frac{1}{(1-p)^2} \sum_{i=1}^n (k_i - 1)$$

and

$$l''(\hat{p}) = - \left(\sum_{i=1}^n k_i \right)^2 \left(\frac{1}{n} + \frac{1}{\sum_{i=1}^n (k_i - 1)} \right) < 0.$$

(c) Since \hat{p} is an average, the central limit theorem will tell us that $\text{Var}(\hat{p}) \approx \text{Var}(X)/n = p^2(1-p)/n$.

Problem 2. Consider an i.i.d. sample of random variables with density function

$$f(x | \sigma) = \frac{1}{2\sigma} \exp\left(-\frac{|x|}{\sigma}\right).$$

(b) Find the maximum likelihood estimate of σ .

(c) Find the asymptotic variance of the mle.

(b) The log likelihood is given by

$$l(\sigma) = \sum_{i=1}^n \log\left(\frac{1}{2\sigma} \exp\left(-\frac{|x_i|}{\sigma}\right)\right) = -\frac{1}{\sigma} \sum_{i=1}^n |x_i| - n \log(2\sigma).$$

Then

$$l'(\sigma) = -\frac{1}{\sigma^2} \sum_{i=1}^n |x_i| - \frac{n}{\sigma}.$$

Setting this equal to 0 gives

$$\hat{\sigma} = \frac{1}{n} \sum_{i=1}^n |x_i|.$$

This is indeed a maximum because

$$l''(\sigma) = \frac{n}{\sigma^2} - \frac{2}{\sigma^3} \sum_{i=1}^n |x_i|$$

and

$$l''(\hat{\sigma}) = \left(\sum_{i=1}^n |x_i| \right)^{-2} \left(\frac{1}{n} - 2 \right) < 0.$$

(c) We have

$$\begin{aligned} I(\sigma) &= -E \left(\frac{\partial^2}{\partial \sigma^2} \log \left(\frac{1}{2\sigma} \exp \left(-\frac{|x|}{\sigma} \right) \right) \right) \\ &= -E \left(\frac{\partial^2}{\partial \sigma^2} \left(-\frac{|x|}{\sigma} - \log(2\sigma) \right) \right) \\ &= -E \left(\frac{\partial}{\partial \sigma} \left(\frac{|x|}{\sigma^2} - \frac{1}{\sigma} \right) \right) \\ &= -E \left(\frac{1}{\sigma^2} - \frac{2|x|}{\sigma^3} \right) \\ &= - \left(\frac{1}{\sigma^2} - \frac{2E(|x|)}{\sigma^3} \right). \end{aligned}$$

Note that for this distribution, $|X|$ follows an exponential distribution with parameter $1/\sigma$. Thus we have

$$E(|X|) = \int_{-\infty}^{\infty} \frac{x}{2\sigma} e^{-|x|/\sigma} dx = \sigma.$$

Plugging this in gives $I(\sigma) = -(1/\sigma^2 - 2\sigma/\sigma^3) = 1/\sigma^2$. Thus $\text{Var}(\hat{\sigma}) = \sigma^2/n$

Problem 3. Suppose that X_1, X_2, \dots, X_n are i.i.d. random variables on the interval $[0, 1]$ with the density function

$$f(x | \alpha) = \frac{\Gamma(2\alpha)}{\Gamma(\alpha)^2} [x(1-x)]^{\alpha-1}$$

where $\alpha > 0$ is a parameter to be estimated from the sample. It can be shown that

$$\begin{aligned} E(X) &= \frac{1}{2} \\ \text{Var}(X) &= \frac{1}{4(2\alpha + 1)}. \end{aligned}$$

(c) What equation does the mle of α satisfy?

(d) What is the asymptotic variance of the mle?

(c) The log likelihood is given by

$$l(\alpha) = \sum_{i=1}^n \log \left(\frac{\Gamma(2\alpha)}{\Gamma(\alpha)^2} [x_i(1-x_i)]^{\alpha-1} \right) = n \log(\Gamma(2\alpha)) - 2n \log(\Gamma(\alpha)) + (\alpha-1) \sum_{i=1}^n \log(x_i(1-x_i))$$

Differentiating we have

$$l'(\alpha) = \frac{2n\Gamma'(2\alpha)}{\Gamma(2\alpha)} - \frac{2n\Gamma'(\alpha)}{\Gamma(\alpha)} + \sum_{i=1}^n \log(x_i(1-x_i)).$$

Then $\hat{\alpha}$ must satisfy $l'(\alpha) = 0$ where $l'(\alpha)$ is as above.

(d) We have

$$\begin{aligned}
I(\alpha) &= -E \left(\frac{\partial^2}{\partial \alpha^2} \log \left(\frac{\Gamma(2\alpha)}{\Gamma(\alpha)^2} [x(1-x)]^{\alpha-1} \right) \right) \\
&= -E \left(\frac{\partial^2}{\partial \alpha^2} (\log(\Gamma(2\alpha)) - 2\log(\Gamma(\alpha)) + (\alpha-1)\log(x(1-x))) \right) \\
&= -E \left(\frac{\partial}{\partial \alpha} \left(\frac{2\Gamma'(2\alpha)}{\Gamma(2\alpha)} - \frac{2\Gamma'(\alpha)}{\Gamma(\alpha)} + \log(x(1-x)) \right) \right) \\
&= -E \left(2 \frac{\Gamma(2\alpha)\Gamma''(2\alpha) - \Gamma'(2\alpha)^2}{\Gamma(2\alpha)^2} - 2 \frac{\Gamma(\alpha)\Gamma''(\alpha) - \Gamma'(\alpha)^2}{\Gamma(\alpha)^2} \right) \\
&= 2 \left(\frac{\Gamma(\alpha)\Gamma''(\alpha) - \Gamma'(\alpha)^2}{\Gamma(\alpha)^2} - \frac{\Gamma(2\alpha)\Gamma''(2\alpha) - \Gamma'(2\alpha)^2}{\Gamma(2\alpha)^2} \right)
\end{aligned}$$

and so

$$\text{Var}(\hat{\alpha}) = \frac{1}{2n} \left(\frac{\Gamma(\alpha)\Gamma''(\alpha) - 2\Gamma'(\alpha)}{\Gamma(\alpha)^2} - \frac{\Gamma(2\alpha)\Gamma''(2\alpha) - 2\Gamma'(2\alpha)}{\Gamma(2\alpha)^2} \right)^{-1}.$$

Problem 4. Suppose that X_1, X_2, \dots, X_n are i.i.d. with density function

$$f(x | \theta) = e^{-(x-\theta)}, \quad x \geq \theta$$

and $f(x | \theta) = 0$ otherwise.

(b) Find the mle of θ .

(b) The log likelihood is given by

$$l(\theta) = \sum_{i=1}^n \log \left(e^{-(x_i-\theta)} \right) = \sum_{i=1}^n (\theta - x_i).$$

Since we're subtracting x_i from θ in each term, we want θ to be as close to the x_i terms as possible. On the other hand, we want $\theta \leq x_i$ for each term, otherwise $f(x_i | \theta) = 0$ for some i and so the joint distribution function is 0. Therefore, we pick $\hat{\theta}$ to be the value of x_i which is less than or equal to all the other values. That is,

$$\hat{\theta} = \min\{x_i | 1 \leq i \leq n\}.$$

Problem 5. The Pareto distribution has been used in economics as a model for a density function with a slowly decaying tail:

$$f(x | x_0, \theta) = \theta x_0^\theta x^{-\theta-1}, \quad x \geq x_0, \quad \theta > 1.$$

Assume that $x_0 > 0$ is given and that X_1, X_2, \dots, X_n is an i.i.d. sample.

(b) Find the mle of θ .

(c) Find the asymptotic variance of the mle.

(b) The log likelihood is given by

$$l(\theta) = \sum_{i=1}^n \log(\theta x_0^\theta x_i^{-\theta-1}) = n \log(\theta) + n\theta \log(x_0) - (\theta+1) \sum_{i=1}^n \log(x_i).$$

Then

$$l'(\theta) = \frac{n}{\theta} + n \log(x_0) - \sum_{i=1}^n \log(x_i).$$

Setting this equal to 0 we have

$$\hat{\theta} = n \left(\sum_{i=1}^n \log(x_i) - n \log(x_0) \right)^{-1} = n \left(\sum_{i=1}^n \log \left(\frac{x_i}{x_0} \right) \right)^{-1}.$$

This is indeed a maximum because

$$l''(\theta) = -\frac{n}{\theta^2} < 0.$$

(c) We have

$$\begin{aligned} I(\theta) &= -E \left(\frac{\partial^2}{\partial \theta^2} \log(\theta x_0^\theta x^{-\theta-1}) \right) \\ &= -E \left(\frac{\partial^2}{\partial \theta^2} (\log(\theta) + \theta \log(x_0) - (\theta+1) \log(x)) \right) \\ &= -E \left(\frac{\partial}{\partial \theta} \left(\frac{1}{\theta} + \log(x_0) - \log(x) \right) \right) \\ &= -E \left(-\frac{1}{\theta^2} \right) \\ &= \frac{1}{\theta^2} \end{aligned}$$

and so $\text{Var}(\hat{\theta}) = \theta^2/n$.

Problem 6. Let X_1, \dots, X_n be an i.i.d. sample from a Rayleigh distribution with parameter $\theta > 0$;

$$f(x | \theta) = \frac{x}{\theta^2} e^{-x^2/(2\theta^2)}, \quad x \geq 0.$$

(b) Find the mle of θ .

(c) Find the asymptotic variance of the mle.

(b) The log likelihood is given by

$$l(\theta) = \sum_{i=1}^n \log \left(\frac{x_i}{\theta^2} e^{-x_i^2/(2\theta^2)} \right) = \sum_{i=1}^n \log(x_i) - \frac{1}{2\theta^2} \sum_{i=1}^n x_i^2 - 2n \log(\theta).$$

Then

$$l'(\theta) = \frac{1}{\theta^3} \sum_{i=1}^n x_i^2 - \frac{2n}{\theta}.$$

Setting this equal to 0 we have

$$\hat{\theta} = \pm \frac{1}{\sqrt{2n}} \sqrt{\sum_{i=1}^n x_i^2}.$$

Then we have

$$l''(\theta) = \frac{2n}{\theta^2} - \frac{3}{\theta^4} \sum_{i=1}^n x_i^2.$$

Putting in either the positive or negative square root gives

$$l''(\hat{\theta}) = -8n^2 \left(\sum_{i=1}^n x_i^2 \right)^{-1} < 0.$$

Thus both values of $\hat{\theta}$ are maximums.

(c) We have

$$\begin{aligned}
 I(\theta) &= -E \left(\frac{\partial^2}{\partial \theta^2} \log \left(\frac{x}{\theta^2} e^{-x^2/(2\theta^2)} \right) \right) \\
 &= -E \left(\frac{\partial^2}{\partial \theta^2} (\log(x) - \frac{x^2}{2\theta^2} - 2 \log(\theta)) \right) \\
 &= -E \left(\frac{\partial}{\partial \theta} \left(\frac{x^2}{\theta^3} - \frac{2}{\theta} \right) \right) \\
 &= -E \left(\frac{2}{\theta^2} - \frac{3x^2}{\theta^4} \right) \\
 &= \frac{3E(x^2)}{\theta^4} - \frac{2}{\theta^2}.
 \end{aligned}$$

We now have

$$E(X^2) = \int_0^\infty \frac{x^3}{\theta^2} e^{-x^2/(2\theta^2)} dx = e^{-x^2/(2\theta^2)} (2\theta^2 + x^2) \Big|_0^\infty = 2\theta^2.$$

Plugging this in gives us

$$I(\theta) = \frac{6\theta^2}{\theta^4} - \frac{2}{\theta^2} = \frac{4}{\theta^2}.$$

Thus $\text{Var}(\hat{\theta}) = \theta^2/4n$.

Problem 7. Let X_1, \dots, X_n be i.i.d. random variables with the density function

$$f(x | \theta) = (\theta + 1)x^\theta, \quad 0 \leq x \leq 1.$$

(b) Find the mle of θ .

(c) Find the asymptotic variance of the mle.

(b) The log likelihood is given by

$$l(\theta) = \sum_{i=1}^n \log((\theta + 1)x_i^\theta) = n \log(\theta + 1) + \theta \sum_{i=1}^n \log(x_i).$$

Then

$$l'(\theta) = \frac{n}{\theta + 1} + \sum_{i=1}^n \log(x_i).$$

Setting this equal to 0 we have

$$\hat{\theta} = -1 - n \left(\sum_{i=1}^n \log(x_i) \right)^{-1}.$$

This is indeed a maximum because

$$l''(\theta) = -\frac{n}{(\theta + 1)^2} < 0.$$

(c) We have

$$\begin{aligned} I(\sigma) &= -E \left(\frac{\partial^2}{\partial \theta^2} \log((\theta + 1)x^\theta) \right) \\ &= -E \left(\frac{\partial^2}{\partial \theta^2} (\log(\theta + 1) + \theta \log(x)) \right) \\ &= -E \left(\frac{\partial}{\partial \theta} \left(\frac{1}{\theta + 1} + \log(x) \right) \right) \\ &= -E \left(-\frac{1}{(\theta + 1)^2} \right) \\ &= \frac{1}{(\theta + 1)^2} \end{aligned}$$

so $\text{Var}(\hat{\theta}) = (\theta + 1)^2/n$.