The Schwarzschild-Tangherlini metric from scattering amplitudes in various dimensions

Stavros Mougiakakos^a and Pierre Vanhove^{a,b,c}

- ^aInstitut de Physique Théorique,
 - Université Paris Saclay, CNRS, F-91191 Gif-sur-Yvette, France
- ^bNational Research University Higher School of Economics, Russian Federation
- ^c Theoretical Physics Department, CERN, 1211 Geneva 23, Switzerland

ABSTRACT: We derive the static Schwarzschild-Tangherlini metric by extracting the classical contributions from the multi-loop vertex functions of a graviton emitted from a massive scalar field. At each loop orders the classical contribution is proportional to a unique master integral given by the massless sunset integral. By computing the scattering amplitudes up to three-loop order in general dimension, we explicitly derive the expansion of the metric up to the fourth post-Minkowskian order $O(G_N^4)$ in four, five and six dimensions. There are ultraviolet divergences that are cancelled with the introduction of higher-derivative non-minimal couplings. The standard Schwarzschild-Tangherlini is recovered by absorbing their effects by an appropriate coordinate transformation induced from the de Donder gauge condition.

Contents

| 1 | Introduction | 2 |
|--------------|---|------------|
| 2 | The Schwarzschild-Tangherlini metric from scalar field am | plitudes 3 |
| | 2.1 The classical contribution of the amplitude | 4 |
| | 2.2 The master integrals for the classical limit | 8 |
| 3 | The metric perturbation from graviton emission | 9 |
| | 3.1 Tree-level amplitude | 10 |
| | 3.2 One-loop amplitude | 11 |
| | 3.3 Two-loop amplitude | 13 |
| | 3.4 Three-loop amplitude | 16 |
| 4 | Non-minimal couplings and renormalised metric | 18 |
| | 4.1 Tree-level insertions | 20 |
| | 4.2 One-loop insertions | 22 |
| | 4.3 Two-loop insertions | 23 |
| | 4.4 The renormalised metric in four dimensions | 24 |
| | 4.5 The renormalised metric in five dimensions | 26 |
| | 4.6 The renormalised stress-tensor in six dimensions | 30 |
| 5 | The Schwarzschild-Tangherlini metric in de Donder gauge in four, five | |
| | and six dimensions | 30 |
| | 5.1 The metric in the de Donder gauge in four dimensions | 31 |
| | 5.2 The metric in the de Donder gauge in five dimensions | 33 |
| | 5.3 The metric in the de Donder gauge in six dimensions | 34 |
| 6 | Recovering the Schwarzschild-Tangherlini metric from the amplitude com- | |
| | putations | 35 |
| | 6.1 The Schwarzschild metric in four dimensions | 35 |
| | 6.2 The Schwarzschild-Tangherlini metric in five dimensions | 36 |
| | 6.3 The Schwarzschild-Tangherlini metric in six dimensions | 37 |
| 7 | Discussion | 37 |
| \mathbf{A} | Fourier transforms | 38 |
| В | Vertices and Propagators | 39 |

1 Introduction

General relativity is a theory for the action of gravity in space and time. The dynamics of the gravitational field is constrained by the Einstein's classical field equations. They are tensorial non-linear equations, because of the self-interaction of the gravitational field, notoriously difficult to solve. It is therefore important to develop efficient methods for studying gravity in various regimes.

General relativity can be embedded in quantum theory where the gravitational force results from the exchange of a quantized massless spin-2 graviton field [1–5]. One can then consider the Einstein-Hilbert term as the first term of a low-energy effective action containing an infinite number of higher derivative operators [6].

The classical limit $\hbar \to 0$ has been studied by Duff in [7] where he showed how to reproduce the classical Schwarzschild metric in four dimensions from quantum tree graphs up to the second order $O(G_N^2)$ in Newton's constant.

The relation between the quantum theory of gravity and the classical Einstein's theory of general relativity has received a new interpretation with the understanding [8–13] that an appropriate (and subtle) $\hbar \to 0$ limit of quantum multi-loop scattering gravitational amplitudes lead to higher G_N -order classical gravity contributions. Considering the importance of such approach for the evaluation of the post-Minkowskian expansion for the gravitational two-body scattering [14–20], we use the procedure given in [12] for extracting the classical contributions from the multi-loop vertex function of a graviton emission from a massive scalar field to recover the Schwarzschild-Tangherlini metric in various dimensions. The scattering amplitude approach works in general dimensions [21–24] and gives the opportunity to explore general relativity in higher-dimensions [25, 26]. At tree-level and one-loop our results agree with the general dimension results in [21, 24]. We show how to reconstruct the metric up to the fourth order $O(G_N^4)$ in Newton's constant by evaluating the scattering amplitudes up to three-loop orders.

Using the procedure designed in [12] we argue, in section 2.1, that the classical contribution at l-loop order is given by the two-point l-loop massless sunset graphs. We verify this explicitly evaluating the classical limit of the quantum scattering amplitudes up to three-loop order.

The scattering amplitudes develop ultraviolet divergences. In section 4, we show how to recover the finite static Schwarzschild-Tangherlini metric by the addition of non-minimal couplings given schematically by (see (4.2) for a precise expression)

$$\delta^{(n)} S^{\text{ct.}} \sim (G_N m)^{\frac{2n}{d-2}} \int d^{d+1} x \sqrt{-g} \nabla^{2(n-1)} \mathcal{R}_{\mu\nu} \partial^{\mu} \phi \partial^{\nu} \phi. \tag{1.1}$$

In four dimensions the non-minimal couplings $\delta^{(1)}S^{\text{ct.}}$ have been introduced in [27] for the analysis up to the third post-Minkowskian order in the context of the world-line formalism. The relation between the world-line formalism and the amplitude approach is detailed in [20]. Higher-derivative couplings with $n \geq 2$ would be needed in four dimensions from the fifth post-Minkowskian order, but they appear at lowest order in higher dimensions. Indeed, we show that in five dimensions one needs to consider higher dimensional of non-minimal couplings $\delta^{(2)}S^{\text{ct.}}$ at the third post-Minkowskian order and $\delta^{(3)}S^{\text{ct.}}$ at the fourth

post-Minkowskian. Interestingly, the metric components are finite in space-time dimensions greater or equal to six, although the stress-tensor develops ultraviolet divergences from one-loop order in odd dimensions and from two-loop order in even dimensions. These divergences are cancelled by the non-minimal couplings $\delta^{(n)}S^{\text{ct.}}$. Actually, we expect that an all order computation in perturbation will require an infinite set of such non-minimal couplings.

We show that the effects of the non-minimal couplings can be reabsorbed by a coordinate transformation, and they do not affect the Schwarzschild-Tangherlini space-time geometry. Since we work in the fixed gauge de Donder gauge, we give the coordinate transformation for extracting the classical space-time metric from the scattering amplitudes in that gauge. Although general relativity is coordinate system invariant, our analysis shows that there is a preferred coordinate system when extracting the classical geometry from scattering amplitudes in the de Donder gauge. The lowest-order n = 1 non-minimal couplings have been shown to arise from the gauge fixing in [20, 24, 28]. We will not address the question of the gauge dependence, but we remark that the choice of coordinate system (or gauge) can be critical for finding solution to Einstein's equations [29].

Since "black hole formation is a robust prediction of the general theory of relativity" [30], it is satisfying to be able to embed such classical solutions in the new understanding of the relation between general relativity and the quantum theory of gravity.

The paper is organised as follows. In section 2 we setup the connection between the perturbation expansion vertex function for the emission a graviton from a massive scalar field and the post-Minkowskian expansion of the static metric in d+1 dimensions. In section 2.1 we show that the classical contribution from the multi-loop amplitudes is given by the massless sunset multi-loop integrals in d dimensions. In section 2.2 we evaluate the master integrals. In section 3 we derive the metric component up to the order $O(G_N^4)$ by computing the relevant amplitudes up to three-loop order in d+1 dimensions. In section 4 we compute the non-minimal couplings required for cancelling the ultraviolet divergences in the amplitude computation. In section 5 we solve the Einstein's equations in four (d=3), five (d=4) and six (d=5) dimensions in the de Donder gauge, and we show in section 6 how these results match the results derived from the amplitude computations. In section 7 we give an interpretation of the results in this paper. The appendix A contains formulæ for the Fourier transforms used in the text, and appendix B the vertices for the scattering amplitude computations.

2 The Schwarzschild-Tangherlini metric from scalar field amplitudes

The Schwarzschild metric is obtained by the gravitational scattering of a scalar field of mass m

$$S = \int d^{d+1}x \sqrt{-g} \left(\frac{R}{16\pi G_N} + \frac{1}{2} g^{\mu\nu} \partial_{\mu} \phi \partial_{\nu} \phi - \frac{1}{2} m^2 \phi^2 \right). \tag{2.1}$$

For further reference Newton's constant has length dimensions $[G_N] = (length)^{d-1}$, the scalar field has dimension $[\phi] = (length)^{1-d}$ and the mass $[m] = (length)^{-1}$. We work with

the mostly negative signature $(+, -, \cdots, -)$ metric.

The graviton emission from a scalar particle of mass $p_1^2 = p_2^2 = m^2$ is given by the three-point vertex function

At each loop order we extract the *l*-loop contribution to the transition density of the stress-energy tensor $\langle T_{\mu\nu}(q^2)\rangle = \sum_{l\geq 0} \langle T_{\mu\nu}^{(l)}(q^2)\rangle$

$$i\mathcal{M}_{3}^{(l)}(p_{1},q) = -\frac{i\sqrt{32\pi G_{N}}}{2} \langle T^{(l)\mu\nu}(q^{2})\rangle \epsilon_{\mu\nu}$$
 (2.3)

where $\epsilon^{\mu\nu}$ is the polarisation of the graviton with momentum $q=p_1-p_2$ is the momentum transfer.

The scattering amplitude computation is not done in the harmonic gauge coordinates $g^{\mu\nu}\Gamma^{\lambda}_{\mu\nu}(g) = 0$ but in the de Donder gauge coordinate system [2, 19, 21, 24, 27]

$$\eta^{\mu\nu}\Gamma^{\lambda}_{\mu\nu}(g) = \eta^{\mu\nu}g^{\lambda\rho}\left(\frac{\partial g_{\rho\mu}}{\partial x^{\nu}} + \frac{\partial g_{\rho\nu}}{\partial x^{\mu}} - \frac{\partial g_{\mu\nu}}{\partial x^{\rho}}\right) = 0, \qquad (2.4)$$

the metric perturbations $g_{\mu\nu} = \eta_{\mu\nu} + \sum_{n\geq 1} h_{\mu\nu}^{(n)}$ satisfy¹

$$\frac{\partial}{\partial x^{\lambda}} h_{\nu}^{\lambda(n)} - \frac{1}{2} \frac{\partial}{\partial x^{\nu}} h^{(n)} = 0.$$
 (2.5)

The de Donder gauge relation between the metric perturbation and the stress-energy tensor reads

$$h_{\mu\nu}^{(l+1)}(\vec{x}) = -16\pi G_N \int \frac{d^d \vec{q}}{(2\pi)^d} e^{i\vec{q}\cdot\vec{x}} \frac{1}{\vec{q}^2} \left(\langle T_{\mu\nu}^{(l)} \rangle^{\text{class.}}(q^2) - \frac{1}{d-1} \eta_{\mu\nu} \langle T^{(l)} \rangle^{\text{class.}}(q^2) \right) . \quad (2.6)$$

In this relation enters the classical contribution at l loop order $\langle T_{\mu\nu}^{(l)} \rangle^{\text{class.}}(q^2)$ defined by the classical limit of the quantum scattering amplitude [10, 12, 13]. From now, we are dropping the super-script class and just use the notation $\langle T_{\mu\nu}^{(l)} \rangle (q^2)$ for the classical contribution.

2.1 The classical contribution of the amplitude

In this section we derive the generic form of the classical contribution of the gravity amplitudes (2.2) in the static limit where $q = (0, \vec{q})$ and $\vec{q}^2 \ll m^2$. The classical limit is obtained by taking $\hbar \to 0$ with the momentum transfer q/\hbar held fixed [13].

¹The harmonic gauge linearized at the first order in perturbation gives (2.5) with n = 1. The higher-order expansions of the harmonic gauge differ from these conditions.

At the l-loop order we have to consider the graphs

The classical contribution emerges as a particular $\hbar \to 0$ limit of the amplitude in [8, 10–13]. The classical limit results in cutting the massive lines, projecting on the contribution from localised sources at different positions in space [12, 31, 32], pictorially represented by shaded blobs

$$\mathcal{M}_{3}^{(l) \text{ class.}}(p_{1},q) = \left.\begin{array}{c} & & & \\$$

In this process one keeps only the leading q^2 contribution from the multi-graviton tree-level amplitudes. The quantum tree-level graphs that were considered in [7] arise from the classical limit of the scattering amplitude up to two-loop order. In the rest of this section, we derive the generic features of the classical limit to all orders in perturbation. We then explicitly evaluate the classical limit up to three-loop order in perturbation.

The quantum amplitude in (2.7) is an l+2 gravitons amplitude with l+1 gravitons attached to the massive scalar line

(2.10)

with the momentum conservation condition $\ell_1 + \cdots + \ell_{l+1} = q = p_1 - p_2$ and the vertex for emitting a graviton from a scalar field²

$$\tau^{\mu\nu}(p_1, p_2) = p_1^{\mu} p_2^{\nu} + p_1^{\nu} p_2^{\mu} + \frac{1}{2} \eta^{\mu\nu} (p_1 - p_2)^2.$$
 (2.11)

This line is attached to an l+2 tree-level graviton amplitude

$$\mathcal{M}^{\mu_1\nu_1,\dots,\mu_{l+1}\nu_{l+1}}(\ell_1,\dots,\ell_{l+1},q) = \underbrace{\qquad \qquad \qquad }_{q}$$
 (2.12)

We have to sum over all the permutation of the graviton lines attached to the scalar lines. Because the gravity amplitude is invariant under the action of the permutation of the graviton lines we have

$$i\mathcal{M}_{3}^{(l)}(p_{1},q) = \frac{1}{\sqrt{4E_{1}E_{2}}} \int \prod_{n=1}^{l} \frac{d^{d+1}\ell_{n}}{(2\pi)^{D}} \left(\sum_{\sigma \in \mathfrak{S}_{l+1}} \mathcal{L}_{\mu_{1}\nu_{1},\dots,\mu_{l+1}\nu_{l+1}}(p_{1},p_{2},\ell_{\sigma(1)},\dots,\ell_{\sigma(l+1)}) \right) \times \prod_{i=1}^{l+1} \frac{i\mathcal{P}^{\mu_{i}\nu_{i},\rho_{i}\sigma_{i}}}{\ell_{i}^{2} + i\epsilon} \mathcal{M}_{\rho_{1}\sigma_{1},\dots,\rho_{l+1}\sigma_{l+1}}(\ell_{1},\dots,\ell_{l+1},q) \quad (2.13)$$

where \mathfrak{S}_{l+1} is the group of permutation of l+1 elements. In the static limit the vertex (2.11) becomes

$$\tau_{\mu\nu}(p_1, p_1 - \ell) \simeq -2m^2 \delta_{\mu}^0 \delta_{\nu}^0,$$
(2.14)

therefore the scalar line approximates to

$$\mathcal{L}(p_1, p_2, \ell_1, \dots, \ell_{l+1}) \simeq \frac{\prod_{i=1}^{l+1} i \sqrt{32\pi G_N} m^2 \delta_{\mu_i}^0 \delta_{\nu_i}^0}{\prod_{i=1}^{l} \left((p_1 - \sum_{j=1}^{i} \ell_j)^2 - m^2 + i\epsilon \right)}.$$
 (2.15)

In the static limit $(p_1 - L)^2 - m^2 + i\epsilon = L^2 - 2p_1 \cdot L + i\epsilon \simeq L_0^2 - \vec{L}^2 - 2mL_0 + i\epsilon$. In the limit where the mass m is large compared to the graviton loop momenta $|L| \ll m$ we have

$$L_0^2 - \vec{L}^2 - 2mL_0 + i\epsilon = \left(L_0 - m - \sqrt{\vec{L}^2 + m^2 - i\epsilon}\right) \left(L_0 - m + \sqrt{\vec{L}^2 + m^2 - i\epsilon}\right)$$

$$\simeq \left(L_0 - 2m - \frac{\vec{L}^2}{2m} + i\epsilon\right) \left(L_0 + \frac{\vec{L}^2}{2m} - i\epsilon\right) \simeq -2m(L_0 - i\epsilon) . \quad (2.16)$$

Therefore we have

$$\mathcal{L}(p_1, p_2, \ell_1, \dots, \ell_{l+1}) \simeq i\sqrt{32\pi G_N} m^2 \delta_{\mu_{l+1}}^0 \delta_{\nu_{l+1}}^0 \prod_{i=1}^l \frac{-i2\sqrt{2\pi G_N} m \delta_{\mu_i}^0 \delta_{\nu_i}^0}{\sum_{j=1}^i \ell_j^0 - i\epsilon}.$$
 (2.17)

²The vertices are given in appendix B. We have stripped of a factor $i\sqrt{8\pi G_N}$ from their normalisation.

Using momentum conservation $\ell_1 + \cdots + \ell_{l+1} = p_1 - p_2$ and that in the static limit $p_1^0 - p_2^0 \simeq 0$ we have

$$\mathcal{L}(p_1, p_2, \ell_1, \dots, \ell_{l+1}) \simeq 2mi\epsilon \prod_{i=1}^{l+1} \frac{-i2\sqrt{2\pi G_N} m \delta_{\mu_i}^0 \delta_{\nu_i}^0}{\sum_{j=1}^i \ell_j^0 - i\epsilon}.$$
 (2.18)

Using the identity³

$$\sum_{\sigma \in \mathfrak{S}_{l+1}} \prod_{i=1}^{l+1} \frac{1}{\sum_{j=1}^{i} x_{\sigma(j)}} = \prod_{i=1}^{l+1} \frac{1}{x_i}.$$
 (2.21)

In the limit $\epsilon \to 0$ the expression vanishes unless some of the ℓ_j^0 vanish at the same time. This means that one needs to pick the residues at $\ell_j^0 = i\epsilon$ for j = 1, ..., l to have a non vanishing answer. This implies that the amplitude (2.13) reduces to

$$i\mathcal{M}_{3}^{(l)}(p_{1},q) \simeq -i^{l} \left(2\sqrt{2\pi G_{N}}m\right)^{l+1} \times \int \prod_{n=1}^{l} \frac{d^{d}\vec{\ell}_{n}}{(2\pi)^{d}} \prod_{i=1}^{l+1} \frac{\mathcal{P}^{00,\rho_{i}\sigma_{i}}}{\prod_{i=1}^{l+1}(\ell_{i}^{2}+i\epsilon)} \mathcal{M}_{\rho_{1}\sigma_{1},\dots,\rho_{l+1}\sigma_{l+1}}(\ell_{1},\dots,\ell_{l+1},q)\Big|_{\ell_{i}^{0}=0}$$
(2.22)

with $\ell_1 + \cdots + \ell_{l+1} = q$. We recall that

$$\mathcal{P}^{00,\rho\sigma} = \delta_0^{\rho} \delta_0^{\sigma} - \frac{\eta^{\rho\sigma}}{D-2}. \tag{2.23}$$

The amplitude (2.22) corresponds to the graph where the scalar line has been collapsed to a point

$$\mathcal{M}_3^{(l)}(p_1,q)\simeq$$
 tree (2.24)

In the static with $q=(0,\vec{q}), |q|\ll m$, the l+2-tree level gravitons amplitude has the leading behaviour

$$\prod_{n=1}^{l+1} \mathcal{P}^{00,\rho_i \sigma_i} \mathcal{M}_{\rho_i \sigma_i, \dots, \rho_{l+1} \sigma_{l+1}}(\ell_1, \dots, \ell_{l+1}, q) \propto \sqrt{G_N}^l q^2, \qquad (2.25)$$

$$\Sigma(l+1) = \sum_{\sigma \in \mathfrak{S}_{l+1}} \prod_{i=1}^{l+1} \frac{1}{\sum_{j=1}^{i} x_{\sigma(j)}} = \frac{1}{x_1 + \dots + x_{l+1}} \sum_{i=1}^{l+1} \sum_{\sigma \in \mathfrak{S}_{l}} \prod_{i=1}^{l} \frac{1}{\sum_{j=1}^{i} \hat{x}_{\sigma(j)}}$$
(2.19)

where $\sigma(n+1) = i$ and the $\{\hat{x}_1, \dots, \hat{x}_l\} = \{x_1, \dots, x_{l+1}\} \setminus \{x_i\}$. By recursion hypothesis we can use the expression for $\Sigma(l)$

$$\Sigma(l+1) = \frac{1}{x_1 + \dots + x_{l+1}} \sum_{i=1}^{l+1} \prod_{i=1}^{l} \frac{1}{\hat{x}_i} = \frac{1}{x_1 + \dots + x_{l+1}} \sum_{i=1}^{l+1} x_i \prod_{i=1}^{l+1} \frac{1}{x_i} = \prod_{i=1}^{l+1} \frac{1}{x_i}.$$
 (2.20)

This was proven in the appendix of [33]. We give here an alternative proof using recursion. For l=1 we have $\Sigma(2) = \frac{1}{x_1(x_1+x_2)} + \frac{1}{x_2(x_1+x_2)} = \frac{1}{x_1x_2}$. Assuming that (2.21) is true at the order l, then at the order l+1 we have

and higher powers of \vec{q}^2 contribute to higher powers of \hbar and are sub-leading quantum corrections (see section 3.1 for more about this).

Therefore, the classical contribution to the stress-tensor in (2.3) is given by⁴

$$\langle T_{\mu\nu}^{(l)} \rangle = \pi^l (G_N m)^l m \left(c_1^{(l)}(d) \delta_\mu^0 \delta_\nu^0 + c_2^{(l)}(d) \left(\frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) \right) J_{(l)}(q^2) , \qquad (2.26)$$

where $c_1^{(l)}(d)$ and $c_2^{(l)}(d)$ are rational functions of the dimension d and $J_{(n)}(q^2)$ is the massless n-loop sunset graph

$$J_{(n)}(\vec{q}^2) = q \longrightarrow q \longrightarrow q = \int \frac{\vec{q}^2}{\prod_{i=1}^n \vec{l}_i^2 (\vec{l}_1 + \dots + \vec{l}_n + \vec{q})^2} \prod_{i=1}^n \frac{d^d \vec{l}_i}{(2\pi)^d}.$$
(2.27)

2.2 The master integrals for the classical limit

The master integrals (2.27) can be evaluated straightforwardly with the parametric representation of the n-loop sunset in D dimensions (see [36])

$$J_{(n)}(\vec{q}^2) = \frac{(\vec{q}^2)^{\frac{n(d-2)}{2}}}{(4\pi)^{\frac{nd}{2}}} \Gamma\left(n+1-\frac{nd}{2}\right) \int_{x_i \ge 0} \left(\frac{1}{x_1} + \dots + \frac{1}{x_n} + 1\right)^{\frac{(n+1)(2-d)}{2}} \prod_{i=1}^n \frac{dx_i}{x_i^{\frac{d}{2}}}$$
(2.28)

since the first Symanzik polynomial is $U_{n+1} = \left(\sum_{i=1}^{n+1} \frac{1}{x_i}\right) \left(\prod_{i=1}^{n+1} x_i\right)$ and the second Symanzik polynomial is $F_{n+1} = -q^2 x_1 \cdots x_{n+1} = \vec{q}^2 x_1 \cdots x_{n+1}$. Changing variables to $y_i = 1/x_i$ we have

$$J_{(n)}(\vec{q}^2) = \frac{(\vec{q}^2)^{\frac{n(d-2)}{2}}}{(4\pi)^{\frac{nd}{2}}} \Gamma\left(n+1-\frac{nd}{2}\right) \int_{y_i \ge 0} (y_1 + \dots + y_n + 1)^{\frac{(n+1)(2-d)}{2}} \prod_{i=1}^n \frac{dy_i}{y_i^{\frac{d-d}{2}}}. \quad (2.29)$$

Using the expression for Euler's beta-function

$$\int_0^\infty (x+a)^\alpha \frac{dx}{x^{1-\beta}} = a^{\alpha+\beta} \frac{\Gamma(-\beta-\alpha)\Gamma(\beta)}{\Gamma(-\alpha)},\tag{2.30}$$

the master integral is readily evaluated to be

$$J_{(n)}(\vec{q}^2) = \frac{(\vec{q}^2)^{\frac{n(d-2)}{2}}}{(4\pi)^{\frac{nd}{2}}} \frac{\Gamma\left(n+1-\frac{nd}{2}\right)\Gamma\left(\frac{d-2}{2}\right)^{n+1}}{\Gamma\left(\frac{(n+1)(d-2)}{2}\right)}.$$
 (2.31)

The master integrals develop ultraviolet poles at loop orders, inducing divergences in the stress-energy tensor. We will show in section 4 how to renormalise these divergences with the introduction of higher-derivative couplings.

⁴We have checked this explicitly to three-loop order using the LiteRed code [34, 35].

3 The metric perturbation from graviton emission

Using the relation (2.6) between the metric perturbation and using the expression (2.26) for the stress-energy tensor in d-dimension in the static limit we have

$$h_{\mu\nu}^{(l+1)}(\vec{q}) = -8\left(c_1^{(l)}(d)(2\delta_{\mu}^0\delta_{\nu}^0 - \eta_{\mu\nu}) + c_2^{(l)}(d)\left(2\frac{q_{\mu}q_{\nu}}{q^2} + (d-2)\eta_{\mu\nu}\right)\right) \times \frac{(\pi G_N m)^{l+1}J_{(l)}(\vec{q}^2)}{\vec{q}^2}.$$
(3.1)

The static space-time components are obtained by computing the Fourier transform in d dimensions

$$h_{\mu\nu}^{(l+1)}(\vec{x}) = \int_{\mathbb{R}^d} h_{\mu\nu}^{(l+1)}(\vec{q}) e^{i\vec{q}\cdot\vec{x}} \frac{d^d\vec{q}}{(2\pi)^d}.$$
 (3.2)

Using the Fourier transformations given in appendix A, and setting $r = |\vec{x}|$, the Fourier transform of the master integrals are given by

$$\int_{\mathbb{R}^d} \frac{J_{(l)}(\vec{q}^2)}{\vec{q}^2} e^{i\vec{q}\cdot\vec{x}} \frac{d^d \vec{q}}{(2\pi)^d} = \left(\frac{\Gamma\left(\frac{d-2}{2}\right)}{4\pi^{\frac{d}{2}}} \frac{1}{r^{d-2}}\right)^{l+1}$$
(3.3)

which is finite to all loop orders. The ultraviolet divergences in the momentum space representation in (2.31) has been cancelled by the Fourier transform.⁵

The tensorial Fourier transform

$$\int_{\mathbb{R}^d} \frac{q_i q_j}{\vec{q}^2} \frac{J_{(l)}(\vec{q}^2)}{\vec{q}^2} e^{i\vec{q}\cdot\vec{x}} \frac{d^d \vec{q}}{(2\pi)^d} = \left(\frac{\Gamma\left(\frac{d-2}{2}\right)}{4\pi^{\frac{d}{2}}} \frac{1}{r^{d-2}}\right)^{l+1} \frac{1}{2 - l(d-2)} \left(-\delta_{ij} + (l+1)(d-2)\frac{x_i x_j}{r}\right). \tag{3.4}$$

diverges for l = 1 and d = 4 and for l = 2 and d = 3, and are otherwise finite.

By spherical symmetry we parameterise the metric in d+1 dimensions

$$ds^{2} = h_{0}(r,d)dt^{2} - h_{1}(r,d)d\vec{x}^{2} - h_{2}(r,d)\frac{(\vec{x} \cdot d\vec{x})^{2}}{\vec{x}^{2}},$$
(3.5)

so that

$$h_i(\vec{x}) = h_i^{(0)} + \sum_{l>1} h_i^{(l)}(\vec{x}), \qquad (3.6)$$

with $h_i^{(0)} = 1, 1, 0$ for i = 0, 1, 2, the post-Minkowskian expansion of the metric components

$$h_0^{(l+1)}(r,d) = -\frac{16}{d-1} \left((d-2)c_1^{(l)}(d) + c_2^{(l)}(d) \right) \left(\frac{\rho(r,d)}{4} \right)^{l+1},$$

$$h_1^{(l+1)}(r,d) = \frac{16}{d-1} \left(c_1^{(l)}(d) - \left(1 + \frac{d-1}{2 - l(d-2)} \right) c_2^{(l)}(d) \right) \left(\frac{\rho(r,d)}{4} \right)^{l+1},$$

$$h_2^{(l+1)}(r,d) = 16 \frac{(d-2)(l+1)}{2 - l(d-2)} c_2^{(l)}(d) \left(\frac{\rho(r,d)}{4} \right)^{l+1}.$$

$$(3.7)$$

⁵This fact had been noticed by L. Planté in his PhD thesis [31].

We have introduced the radial parameter

$$\rho(r,d) = \frac{\Gamma\left(\frac{d-2}{2}\right)}{\pi^{\frac{d-2}{2}}} \frac{G_N m}{r^{d-2}},$$
(3.8)

which is our post-Minkowskian expansion parameter. Recall that in d+1 dimensions the length dimension of $[G_N m] = (length)^{d-2}$ and $\rho(r,d)$ is dimensionless.

The metric component present poles in four dimensions (d=3) from two-loop order and in five dimensions (d=4) from one-loop order. Such divergences will be removed by the contribution from the non-minimal coupling contributions in section 4.

3.1 Tree-level amplitude

At tree-level, the only contributing diagram is

$$\mathcal{M}_3^{(0)}(p_1,q) = \bigvee_{p_1} q \tag{3.9}$$

is the emission of a graviton from the scattering of two massive scalars of momenta p_1 and p_2 and $p_1^2 = p_2^2 = m^2$ with momentum transfert $q = p_1 - p_2$. The scattering amplitude is given by the 2-scalar-1-graviton vertex $\tau^{\mu\nu}(p_1, p_2)$ in (B.2)

$$i\mathcal{M}_{3}^{(0)}(p_{1},q) = -\frac{i\sqrt{32\pi G_{N}}}{2\sqrt{4E_{1}E_{2}}}\epsilon^{\mu\nu}\tau_{\mu\nu} = -\frac{i\sqrt{32\pi G_{N}}}{2}\epsilon^{\mu\nu}\left(p_{1\mu}p_{2\nu} + p_{2\mu}p_{1\nu} - \eta_{\mu\nu}(p_{1}\cdot p_{2} - m^{2})\right).$$
(3.10)

Using that $P = (p_1 + p_2)/2$ and $q = p_1 - p_2$ we have that

$$i\mathcal{M}_{3}^{(0)}(p_{1},q) = -\frac{i\sqrt{32\pi G_{N}}}{2\sqrt{4E_{1}E_{2}}}\epsilon^{\mu\nu}\left(2P_{\mu}P_{\nu} - \frac{1}{2}(q_{\mu}q_{\nu} - \eta_{\mu\nu}q^{2})\right). \tag{3.11}$$

In the static limit $q = p_1 - p_2 \simeq (0, \vec{q}), E_1 \simeq E_2 \simeq m$ and $|\vec{q}| \ll m$ we have

$$\langle T_{\mu\nu}^{(0)}(q^2)\rangle \simeq m\delta_{\mu}^0\delta_{\nu}^0 + \left(\frac{q_iq_j}{\bar{z}_{q^2}}\eta_{\mu}^i\eta_{\nu}^j + \frac{1}{2}\eta_{\mu\nu}\right)\bar{q}^2.$$
 (3.12)

The \vec{q}^2 term in this expression is the contact term which has a higher power of \hbar and does not contribute to the classical limit [12, 37]. The coefficients of the classical contribution to the stress-tensor at tree-level are given by

$$c_1^{(0)}(d) = 1,$$

 $c_2^{(0)}(d) = 0.$ (3.13)

From this we deduce the metric components in d+1 dimensions using (3.7)

$$h_0^{(1)}(r,d) = -4\frac{d-2}{d-1}\rho(r,d),$$


$$h_1^{(1)}(r,d) = \frac{4}{d-1}\rho(r,d),$$

$$h_2^{(1)}(r,d) = 0,$$
(3.14)

where $\rho(r,d)$ is defined in (3.8). This reproduces the expression given in [21, 24].

3.2 One-loop amplitude

At one-loop the only contributing diagram to the classical limit is



from which we extract the one-loop contribution to the stress-energy tensor in d+1 dimensions

$$T^{(1)} \mu\nu(q^2) = \frac{i8\pi G_N}{\sqrt{4E_1E_2}} \int \frac{d^{d+1}l}{(2\pi)^D} \frac{\tau^{\sigma\rho}(p_1, l+p_1)\tau_{(3)\sigma\rho,\kappa\delta}^{\mu\nu}(l,q)\tau^{\kappa\delta}(p_2, l+p_1)}{(l^2+i\epsilon)((l+q)^2+i\epsilon)((l+p_1)^2-m^2+i\epsilon)},$$
(3.16)

where $\tau_{(3) \pi\rho,\sigma\tau}^{\mu\nu}(p_1,p_2)$ is the three graviton vertex and $\tau^{\mu\nu}(p_1,p_2)$ the vertex for the emission of a graviton from two scalars with momenta p_1 and p_2 . We refer to appendix B for definitions and normalisation of our vertices.

In the static limit, $\bar{q}^2 \ll m^2$, the classical contribution coming from the two scalars to one-graviton vertex is

$$\tau_{\alpha\beta} \approx 2m^2 \delta_{\alpha}^0 \delta_{\beta}^0, \tag{3.17}$$

using that $p_1^2 = p_2^2 = m^2$. This gives for the stress-energy tensor

$$T^{(1)\mu\nu}(q^2) = i16\pi G_N m^3 \int \frac{d^{d+1}l}{(2\pi)^D} \frac{\tau_{(3)00,00}^{\mu\nu}(l,q)}{(l^2+i\epsilon)((l+q)^2+i\epsilon)((l+p_1)^2-m^2+i\epsilon)}.$$
 (3.18)

At this point, we want to focus on the computation of the classical contribution at the static limit. Thus, we will employ a trick, which will prove useful for higher loops. We symmetrize the diagram

$$T^{(1) \mu\nu}(q^2) = i8\pi G_N m^3 \int \frac{d^{d+1}l}{(2\pi)^D} \frac{\tau_{(3)00,00}^{\mu\nu}(l,q)}{(l^2 + i\epsilon)((l+q)^2 + i\epsilon)} \times \left[\frac{1}{(l+p_1)^2 - m^2 + i\epsilon} + \frac{1}{(l-p_2)^2 - m^2 + i\epsilon} \right]. \quad (3.19)$$

In the approximation $l^2 \ll m^2$ we have $(l+p_i)^2-m^2=l^2+2l\cdot p_1=l^2+2l_0E-\vec{l}\cdot \vec{q}\simeq l_0^2+2ml_0$ and the amplitude reduces at leading order

$$T^{(1) \mu\nu}(q^2) \simeq i8\pi G_N m^3 \int \frac{d^{d+1}l}{(2\pi)^D} \frac{\tau_{(3)00,00}^{\mu\nu}(l,q)}{(l^2+i\epsilon)((l+q)^2+i\epsilon)} \times \left[\frac{1}{l_0^2+2ml_0+i\epsilon} + \frac{1}{l_0^2-2ml_0+i\epsilon}\right]. \quad (3.20)$$

It is obvious that at $\mathcal{O}(\epsilon^0)$ order we get a zero contribution at leading order in 1/m, since $l_0 \ll m$. Thus, we can compute the leading contribution of the integral over l_0 via Cauchy's theorem, by taking the residue $2ml_0 = i\epsilon$ and closing the contour of integration in the upper half-plane⁶

$$T^{(1)\mu\nu}(q^2) = 4\pi G_N m^2 \int \frac{d^d \vec{l}}{(2\pi)^d} \frac{\tau_{(3)00,00}^{\mu\nu}(l,q)}{(\vec{l}^2 - i\epsilon)((\vec{l} + \vec{q})^2 - i\epsilon)} \bigg|_{l_0 = 0}, \tag{3.21}$$

with

$$\tau_{(3)00,00}^{\mu\nu}(l,q) = \frac{1}{d-1} \left((d-2) \left(l^{\mu} l^{\nu} + (l+q)^{\mu} (l+q)^{\nu} + q^{\mu} q^{\nu} + \frac{3}{2} \eta^{\mu\nu} \vec{q}^{2} \right) - 2(d-2) \left(\vec{l_{1}}^{2} + (\vec{l_{1}} + \vec{q})^{2} \right) \left(\delta_{0}^{\mu} \delta_{0}^{\mu} - \frac{\eta^{\mu\nu}}{4} \right) - 2(d-3) \vec{q}^{2} \delta_{0}^{\mu} \delta_{0}^{\mu} \right).$$

$$(3.22)$$

The component of the stress-tensor are proportional to the one-loop master integral $J_{(1)}(\vec{q}^2)$ as expected from the general discussion of section 2.2

$$\langle T_{\mu\nu}^{(1)} \rangle = \pi G_N m^2 \left(c_1^{(1)}(d) \delta_\mu^0 \delta_\nu^0 + c_2^{(1)}(d) \left(\frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) \right) J_{(1)}(q^2) , \qquad (3.23)$$

with the master integral

$$J_{(1)}(q^2) = \frac{\Gamma\left(\frac{4-d}{2}\right)\Gamma\left(\frac{d-2}{2}\right)^2}{2^d \pi^{\frac{d}{2}}\Gamma(d-2)} \left(\vec{q}^2\right)^{\frac{d-2}{2}}, \tag{3.24}$$

and the coefficients

$$c_1^{(1)}(d) = -\frac{2(4d^2 - 15d + 10)}{(d-1)^2},$$

$$c_2^{(1)}(d) = -\frac{2(d-2)(3d-2)}{(d-1)^2}.$$
(3.25)

3.2.1 The one-loop contribution to the metric components

Using (3.7) we get for the metric components in d+1 dimensions

$$h_0^{(2)}(r,d) = \frac{8(d-2)^2}{(d-1)^2} \rho(r,d)^2,$$

$$h_1^{(2)}(r,d) = -\frac{4(2d^2 - 9d + 14)}{(d-4)(d-1)^2} \rho(r,d)^2,$$

⁶One could have taken the residue at $2ml_0 = -i\epsilon$ and closing the contour in the lower half-plane with the same result.

$$h_2^{(2)}(r,d) = \frac{4(d-2)^2(3d-2)}{(d-4)(d-1)^2}\rho(r,d)^2,$$
(3.26)

where $\rho(r,d)$ is defined in (3.8).


This reproduces the expression given in [21] and the expression in [24, eq. (22)] for $\alpha = 0$.

3.3 Two-loop amplitude

The diagrams contributing to the classical corrections at third post-Minkowskian order of the metric at the two-loop graphs

$$i\mathcal{M}_{3}^{(2)}(p_1, q) = -\sqrt{32\pi G_N} T^{(2)\mu\nu} \epsilon_{\mu\nu},$$
 (3.27)

there are four contributions



3.3.1 The diagrams (a), (b), (c)

The sum of the contributions from the diagrams (a), (b), (c) after appropriate labelling of the momenta, can be expressed as

$$\sum_{i=a}^{c} T_{(i)}^{(2)\mu\nu} = -\frac{16G_{N}^{2}\pi^{2}}{m} \int \prod_{n=1}^{3} \frac{d^{d+1}l_{n}}{(2\pi)^{2d}} \delta(l_{1} + l_{2} + l_{3} + q)$$

$$\times \frac{\tau^{\gamma\delta}(p_{1}, l_{1} + p_{1})\tau^{\sigma\tau}(l_{1} + p_{1}, -l_{2} + p_{1})\tau^{\iota\theta}(l_{2} - p_{2}, -p_{2})\tau^{\phi\chi}_{(3)\iota\theta,\sigma\tau}(-l_{2}, l_{1} + q) \cdot \mathcal{P}^{\alpha\beta}_{\phi\chi} \cdot \tau^{\mu\nu}_{(3)\alpha\beta,\gamma\delta}(l_{1} + q, q)}{l_{1}^{2}l_{2}^{2}l_{3}^{2}(l_{1} + q)^{2}}$$

$$\times \left(\frac{1}{(l_{1} + p_{1})^{2} - m^{2}} \frac{1}{(l_{2} - p_{2})^{2} - m^{2}} + \frac{1}{(l_{3} + p_{1})^{2} - m^{2}} \frac{1}{(l_{1} - p_{2})^{2} - m^{2}} + \frac{1}{(l_{3} + p_{1})^{2} - m^{2}} \frac{1}{(l_{2} - p_{2})^{2} - m^{2}}\right). \quad (3.28)$$

Using the approximate form of the two scalars one graviton vertex in (3.17) and $(l_1+p_1)^2 - m^2 \approx 2ml_1^0$ and taking the residue $2ml_i^0 = i\epsilon$, since for the rest of the residues we get a zero contribution at order $\mathcal{O}(\epsilon^0)$, we get

$$\sum_{i=a}^{c} T_{(i)}^{(2)\,\mu\nu} = 32\pi^2 G_N^2 m^3 \int \prod_{n=1}^2 \frac{d^{d+1} l_n}{(2\pi)^{2d}} \frac{\tau_{(3)\alpha\beta,00}^{\mu\nu}(l_1+q,q) \cdot \mathcal{P}_{\phi\chi}^{\alpha\beta} \cdot \tau_{(3)\,00,00}^{\phi\chi}(-l_2,l_1+q)}{(\vec{l_1})^2 (\vec{l_2})^2 (\vec{l_3})^2 (\vec{l_1}+\vec{q})^2} \Bigg|_{\substack{l_1^0 = l_2^0 = 0 \\ (3.29)}},$$

with

$$\eta_{\mu\nu}\tau^{\mu\nu}_{(3)\phi\chi,00}(l_1+q,q) = \left(l^{\mu}l^{\nu} - (l+q)^{\mu}(l+q)^{\nu} - q^{\mu}q^{\nu}\right) - \frac{3}{2}\eta^{\mu\nu}\vec{q}^2\left(\eta^{\mu\nu} - (d-1)\delta_0^{\mu}\delta_0^{\nu}\right) + \frac{\eta^{\mu\nu}}{2}\left(\vec{l_1}^2 - (\vec{l_1} + \vec{q})^2\right) - \frac{5-d}{2}\delta_0^{\mu}\delta_0^{\nu}\left(\vec{l_1}^2 + (\vec{l_1} + \vec{q})^2\right), \quad (3.30)$$

and

$$\delta_{\mu}^{0}\delta_{\nu}^{0}\tau_{(3)\phi\chi,00}^{\mu\nu}(l_{1}+q,q) = \frac{1}{d-1} \left((d-3) \left((l+q)^{\mu}(l+q)^{\nu} + q^{\mu}q^{\nu} \right) + (d-1) \left(l_{1}^{\mu}l_{1}^{\nu} - \frac{\vec{l_{1}}^{2}}{2} (3\delta_{0}^{\mu}\delta_{0}^{\nu} - \eta^{\mu\nu}) \right) + \frac{\eta^{\mu\nu} - \delta_{0}^{\mu}\delta_{0}^{\nu}}{2} \left(\vec{q}^{2}(d-5) + (3d-7)(\vec{l_{1}} + \vec{q})^{2} \right) \right), \quad (3.31)$$

and

$$\tau_{(3)00,00}^{\mu\nu}(l,q) = \frac{1}{d-1} \left((d-2) \left(l^{\mu} l^{\nu} + (l+q)^{\mu} (l+q)^{\nu} + q^{\mu} q^{\nu} + \frac{3}{2} \eta^{\mu\nu} \bar{q}^{2} \right) - 2(d-2) \left(\vec{l_{1}}^{2} + (\vec{l_{1}} + \bar{q})^{2} \right) \left(\delta_{0}^{\mu} \delta_{0}^{\mu} - \frac{\eta^{\mu\nu}}{4} \right) - 2(d-3) \bar{q}^{2} \delta_{0}^{\mu} \delta_{0}^{\mu} \right).$$
(3.32)

Using the LiteRed code [34, 35] in d dimensions, we find that all the contributions are proportional to the master integral as expected from the general discussion of section 2.2

$$J_{(2)}(\vec{q}) = \int \prod_{i=1}^{2} \frac{d^{d}\vec{l}_{i}}{(2\pi)^{d}} \frac{\vec{q}^{2}}{\prod_{i=1}^{2} \vec{l}_{i}^{2} (\vec{l}_{1} + \vec{l}_{2} + \vec{q})^{2}}$$

$$= -\frac{\vec{q}^{2}}{32\pi^{2} (d-3)} - \left(-3 + \gamma_{E} - \log(4\pi) + \log(\vec{q}^{2})\right) \vec{q}^{2} + O(d-3), \quad (3.33)$$

where $\gamma_E = 0.57721 \cdots$ is the Euler-Mascheroni constant [38].

We find for the 00-component

$$\sum_{i=1}^{c} T_{(i)}^{(2)\,00} = \frac{32\pi^2 G_N^2 m^3}{3} \frac{6d^3 - 45d^2 + 134d - 160}{(d-4)(d-1)^2} J_{(2)}(\vec{q}^2), \qquad (3.34)$$

and for the trace part

$$\sum_{i=a}^{c} T_{(i)}^{(2)\mu\nu} \eta_{\mu\nu} = -\frac{32\pi^2 G_N^2 m^3}{3} \frac{10d^3 - 63d^2 + 123d - 86}{(d-1)^2} J_{(2)}(\vec{q}^2). \tag{3.35}$$

3.3.2 The diagrams (d)

The diagram (d) after symmetrisation over the massive scalar legs reads

$$T_{(d)}^{(2)\mu\nu} = -\frac{32G_N^2\pi^2}{3m} \int \prod_{n=1}^3 \frac{d^{d+1}l_n}{(2\pi)^{2d}} \frac{\delta(l_1+l_2+l_3+q)}{l_1^2 l_2^2 l_3^2} \left(\frac{1}{(l_1+p_1)^2 - m^2 + i\epsilon} \frac{1}{(l_2-p_2)^2 - m^2 + i\epsilon} + \frac{1}{(l_3+p_1)^2 - m^2 + i\epsilon} \frac{1}{(l_1-p_2)^2 - m^2 + i\epsilon} + \frac{1}{(l_3+p_1)^2 - m^2 + i\epsilon} \frac{1}{(l_2-p_2)^2 - m^2 + i\epsilon} \right)$$

$$\times \tau^{\gamma\delta}(p_1, l_1 + p_1)\tau^{\sigma\tau}(l_1 + p_1, -l_2 + p_1)\tau^{\iota\theta}(l_2 - p_2, -p_2)\tau^{\mu\nu}_{(4)\gamma\delta,\sigma\tau,\iota\theta}(q, l_1, l_2, l_3), \quad (3.36)$$

and leads to the contribution

$$T_{(d)}^{(2)\mu\nu} = -\frac{64\pi^2 G_N^2 m^3}{3} \int \prod_{n=1}^2 \frac{d^{d+1} l_n}{(2\pi)^d} \frac{\tau_{(4)\,00,00,00}^{\mu\nu}(q,l_1,l_2,-l_1-l_2-q)}{(\vec{l_1})^2 (\vec{l_2})^2 (\vec{l_1}+\vec{l_2}+\vec{q})^2} \Bigg|_{\substack{l_1^0 = l_2^0 = 0}}$$
(3.37)

with the vertex

$$\tau_{(4)\,00,00,00}^{\mu\nu}(q,l_1,l_2,l_3) = \frac{1}{(d-1)^2} \left(\vec{q}^2 \frac{\delta_0^{\mu} \delta_0^{\nu}}{2} (7d^2 - 45d + 70) - \vec{q}^2 \frac{\eta^{\mu\nu}}{2} (d-2)(6d-23) \right) + (d-2) \left((9-2d)q^{\mu}q^{\nu} + (7-2d)\left(l_1^{\mu}l_1^{\nu} + l_2^{\mu}l_2^{\nu} + l_3^{\mu}l_3^{\nu}\right) \right) + \frac{d-2}{2} \left(\vec{l_1}^2 + \vec{l_2}^2 + \vec{l_3}^2 \right) \left(\delta_0^{\mu} \delta_0^{\nu} (7d-23) - \eta^{\mu\nu} (2d-9) \right) \right). \quad (3.38)$$

Evaluating these integral we find, for the 00-component

$$T_{(d)}^{(2)\,00} = -\frac{32\pi^2 G_N^2 m^3}{3} \frac{(4-d)(6-d)}{(d-1)^2} J_{(2)}(\vec{q}^2) , \qquad (3.39)$$

and for the trace part

$$T_{(d)}^{(2)\mu\nu}\eta_{\mu\nu} = \frac{64\pi^2 G_N^2 m^3}{3} \frac{3d^3 - 20d^2 + 41d - 30}{(d-1)^2} J_{(2)}(\vec{q}^2). \tag{3.40}$$

3.3.3 The two-loop contribution to the metric components

Summing up all the contributions the two-loop stress-tensor is given by

$$\langle T_{\mu\nu}^{(2)} \rangle = \pi^2 G_N^2 m^3 \left(c_1^{(2)}(d) \delta_\mu^0 \delta_\nu^0 + c_2^{(2)}(d) \left(\frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) \right) J_{(2)}(q^2) , \tag{3.41}$$

with the coefficients given by

$$c_1^{(2)}(d) = \frac{32}{3(d-4)(d-1)^3} \left(9d^4 - 70d^3 + 203d^2 - 254d + 104\right),$$

$$c_2^{(2)}(d) = \frac{64(d-2)}{3(d-4)(d-1)^3} \left(2d^3 - 13d^2 + 25d - 10\right),$$
(3.42)

and the expression for the master integral

$$J_{(2)}(\vec{q}^2) = \frac{\Gamma(3-d)\Gamma\left(\frac{d-2}{2}\right)^3}{(4\pi)^d\Gamma\left(\frac{3(d-2)}{2}\right)} \left(\vec{q}^2\right)^{d-2} . \tag{3.43}$$

From which we extract the metric components using the relations (3.7) (using the definition of $\rho(r,d)$ in (3.8))

$$h_0^{(3)}(r,d) = -\frac{8(3d-7)(d-2)^3}{(d-4)(d-1)^3}\rho(r,d)^3,$$

$$h_1^{(3)}(r,d) = \frac{8(7d^4 - 63d^3 + 214d^2 - 334d + 212)}{3(d-3)(d-4)(d-1)^3} \rho(r,d)^3,$$


$$h_2^{(3)}(r,d) = -\frac{8(d-2)^2(2d^3 - 13d^2 + 25d - 10)}{(d-3)(d-4)(d-1)^3} \rho(r,d)^3.$$
(3.44)

3.4 Three-loop amplitude

The diagrams contributing to the classical corrections at third post-Minkowskian order of the metric at the two-loop graphs

$$i\mathcal{M}_{3}^{(3)}(p_{1},q) = -\sqrt{32\pi G_{N}}T^{(3)\mu\nu}\epsilon_{\mu\nu},$$
(3.45)

where the three-loop stress-tensor is given by five distinct diagrams



As before, we permute the internal momenta such that by taking the residue at $2ml_i^0 = i\epsilon$ from the massive propagators, we extract the non-analytic terms which contribute to the classical metric in the static limit. After taking the residues and including the symmetry factors

$$\begin{split} T_{(a)}^{(3)\,\mu\nu} &= 64\pi^3 G_N^3 m^4 \int \prod_{n=1}^3 \frac{d^d \vec{l}_n}{(2\pi)^d} \frac{\tau_{(3)\pi\rho,\sigma\tau}^{\mu\nu}(l_1+l_2,q)\tau_{(3)}^{\pi\rho}(-l_1,l_1+l_2)\tau_{(3)}^{\sigma\tau}(-l_3,l_3+l_4)}{(\vec{l}_1)^2(\vec{l}_2)^2(\vec{l}_3)^2(\vec{l}_4)^2(\vec{l}_1+\vec{l}_2)^2(\vec{l}_3+\vec{l}_4)^2} \bigg|_{l_1^0=l_2^0=l_3^0=0}, \\ T_{(b)}^{(3)\,\mu\nu} &= 256\pi^3 G_N^3 m^4 \int \prod_{n=1}^3 \frac{d^d \vec{l}_n}{(2\pi)^d} \frac{\tau_{(3)\sigma\tau,00}^{\mu\nu}(l_1+q,q)\tau_{(3)}^{\pi\rho}(-l_3,l_3+l_4)\tau_{(3)00,\pi\rho}^{\sigma\tau}(-l_2,l_1+q)}{(\vec{l}_1)^2(\vec{l}_2)^2(\vec{l}_3)^2(\vec{l}_4)^2(\vec{l}_1+\vec{q})^2(\vec{l}_3+\vec{l}_4)^2} \bigg|_{l_1^0=l_2^0=l_3^0=0}, \\ T_{(c)}^{(3)\,\mu\nu} &= -\frac{512\pi^3 G_N^3 m^4}{3} \int \prod_{n=1}^3 \frac{d^d \vec{l}_n}{(2\pi)^d} \frac{\tau_{(3)\alpha\beta,00}^{\mu\nu}(l_1+q,q)\tau_{(4)00,00,00}^{\alpha\beta}(l_1+q,l_2,l_3,l_4)}{(\vec{l}_1)^2(\vec{l}_2)^2(\vec{l}_3)^2(\vec{l}_4)^2(\vec{l}_1+\vec{q})^2} \bigg|_{l_1^0=l_2^0=l_3^0=0}, \\ T_{(d)}^{(3)\,\mu\nu} &= -256\pi^3 G_N^3 m^4 \int \prod_{n=1}^3 \frac{d^d \vec{l}_n}{(2\pi)^d} \frac{\tau_{(3)}^{\gamma\delta}(-l_3,l_3+l_4)\tau_{(4)\gamma\delta,00,00}^{\mu\nu}(q,l_1,l_2,l_3+l_4)}{(\vec{l}_1)^2(\vec{l}_2)^2(\vec{l}_3)^2(\vec{l}_4)^2(\vec{l}_3+\vec{l}_4)^2} \bigg|_{l_1^0=l_2^0=l_3^0=0}, \\ T_{(e)}^{(3)\,\mu\nu} &= \frac{256\pi^3 G_N^3 m^4}{3} \int \prod_{n=1}^3 \frac{d^d \vec{l}_n}{(2\pi)^d} \frac{\tau_{(5)00,00,00,00}^{\mu\nu}(q,l_1,l_2,l_3,l_4)}{(\vec{l}_1)^2(\vec{l}_2)^2(\vec{l}_3)^2(\vec{l}_4)^2} \bigg|_{l_1^0=l_2^0=l_3^0=0}, \\ T_{(e)}^{(3)\,\mu\nu} &= \frac{256\pi^3 G_N^3 m^4}{3} \int \prod_{n=1}^3 \frac{d^d \vec{l}_n}{(2\pi)^d} \frac{\tau_{(5)00,00,00,00}^{\mu\nu}(q,l_1,l_2,l_3,l_4)}{(\vec{l}_1)^2(\vec{l}_2)^2(\vec{l}_3)^2(\vec{l}_4)^2} \bigg|_{l_1^0=l_2^0=l_3^0=0}, \\ T_{(e)}^{(3)\,\mu\nu} &= \frac{256\pi^3 G_N^3 m^4}{3} \int \prod_{n=1}^3 \frac{d^d \vec{l}_n}{(2\pi)^d} \frac{\tau_{(5)00,00,00,00}^{\mu\nu}(q,l_1,l_2,l_3,l_4)}{(\vec{l}_1)^2(\vec{l}_2)^2(\vec{l}_3)^2(\vec{l}_4)^2} \bigg|_{l_1^0=l_2^0=l_3^0=0}, \\ T_{(e)}^{(3)\,\mu\nu} &= \frac{256\pi^3 G_N^3 m^4}{3} \int \prod_{n=1}^3 \frac{d^d \vec{l}_n}{(2\pi)^d} \frac{\tau_{(5)00,00,00,00}^{\mu\nu}(q,l_1,l_2,l_3,l_4)}{(\vec{l}_1)^2(\vec{l}_2)^2(\vec{l}_3)^2(\vec{l}_4)^2} \bigg|_{l_1^0=l_2^0=l_3^0=0}, \\ T_{(e)}^{(3)\,\mu\nu} &= \frac{256\pi^3 G_N^3 m^4}{3} \int \prod_{n=1}^3 \frac{d^d \vec{l}_n}{(2\pi)^d} \frac{\tau_{(5)00,00,00,00}^{\mu\nu}(q,l_1,l_2,l_3,l_4)}{(\vec{l}_1)^2(\vec{l}_2)^2(\vec{l}_3)^2(\vec{l}_4)^2} \bigg|_{l_1^0=l_2^0=l_3^0=0}, \\ T_{(e)}^{(3)\,\mu\nu} &= \frac{256\pi^3 G_N^3 m^4}{3} \int \prod_{n=1}^3 \frac{d^d \vec{l}_n}{(2\pi)^d} \frac{\tau_{(5)00,00,00,00}^{\mu\nu$$

with the five-graviton vertex contribution

$$\tau_{(5)\ 00,00,00,00}^{\mu\nu}(k_{1},k_{2},k_{3},k_{4},k_{5}) := \tilde{\tau}_{(5)\ \alpha\beta,\gamma\delta,\epsilon\eta,\kappa\lambda}^{\mu\nu}(k_{1},k_{2},k_{3},k_{4},k_{5}) \mathcal{P}_{00}^{\alpha\beta} \mathcal{P}_{00}^{\gamma\delta} \mathcal{P}_{00}^{\epsilon\eta} P_{00}^{\kappa\lambda} \\
= \frac{1}{4(d-1)^{3}} \left(4\delta_{\mu}^{0} \delta_{\nu}^{0} \left(4(2d^{3}-18d^{2}+57d-61)k_{1}^{2} + (d-2)(8d^{2}-47d+79) \sum_{i=2}^{5} k_{i}^{2} \right) \\
- (d-2)\eta_{\mu\nu} \left((29d^{2}-191d+362)k_{1}^{2} + (7d^{2}-61d+142) \sum_{i=2}^{5} k_{i}^{2} \right) \\
+ 2(d-2) \left((11d^{2}-73d+150)k_{1\mu}k_{1\nu} + (7d^{2}-53d+102)(k_{2\mu}k_{2\nu} + k_{3\mu}k_{3\nu} + k_{4\mu}k_{4\nu} + k_{5\mu}k_{5\nu}) \right) \right). \tag{3.47}$$

where the vertex $\tau^{\mu\nu}_{(5) \alpha\beta,\gamma\delta,\epsilon\eta,\kappa\lambda}(k_1,k_2,k_3,k_4,k_5)$ has been derived using the results of [39]. The integral reduction is done using the LiteRed code [34, 35] in d dimensions. In agreement with the general analysis of section 2.2, we find that the classical contribution is proportional to the single master integral

$$J_{(3)}(\vec{q}^2) = \int \frac{d^d \vec{l}_1 d^d \vec{l}_2 d^d \vec{l}_3}{(2\pi)^{3d}} \frac{\vec{q}^2}{\vec{l}_1^2 \vec{l}_2^2 \vec{l}_3^2 (\vec{l}_1 + \vec{l}_2 + \vec{l}_3 + \vec{q})^2}.$$
 (3.48)

3.4.1 The $\mu = \nu = 0$ component

$$\begin{split} T_{(a)}^{(3)\,00} &= -\frac{32\pi^3 G_N^3 m^4}{3} \frac{3d^5 - 169d^4 + 1378d^3 - 4592d^2 + 7256d - 4752}{3} J_{(3)}(\vec{q}^2), \\ T_{(b)}^{(3)\,00} &= -\frac{128\pi^3 G_N^3 m^4}{3} \frac{68d^6 - 1003d^5 + 6211d^4 - 20820d^3 + 40020d^2 - 41584d + 17824}{3} J_{(3)}(\vec{q}^2), \\ T_{(c)}^{(3)\,00} &= \frac{64\pi^3 G_N^3 m^4}{3} \frac{37d^5 - 502d^4 + 2731d^3 - 7486d^2 + 10164d - 5256}{3} J_{(3)}(\vec{q}^2), \\ T_{(d)}^{(3)\,00} &= \frac{32\pi^3 G_N^3 m^4}{3} \frac{53d^4 - 615d^3 + 2690d^2 - 5572d + 4840}{(d-4)(d-1)^3} J_{(3)}(\vec{q}^2), \\ T_{(e)}^{(3)\,00} &= 64\pi^3 G_N^3 m^4 \frac{(6-d)(d^2 - 7d + 14)}{(d-1)^3} J_{(3)}(\vec{q}^2). \end{split}$$

3.4.2 Contraction with $\eta_{\mu\nu}$

$$\begin{split} T_{(a)}^{(3)}{}^{\mu\nu}\eta_{\mu\nu} &= \frac{32\pi^3G_N^3m^4}{3}\frac{85d^6 - 1126d^5 + 6307d^4 - 19114d^3 + 32944d^2 - 30472d + 11952}{3}J_{(3)}(\vec{q}^2), \\ T_{(b)}^{(3)}{}^{\mu\nu}\eta_{\mu\nu} &= \frac{128\pi^3G_N^3m^4}{3}\frac{168d^6 - 2231d^5 + 12319d^4 - 35796d^3 + 57396d^2 - 48304d + 16736}{3}J_{(3)}(\vec{q}^2), \\ T_{(c)}^{(3)}{}^{\mu\nu}\eta_{\mu\nu} &= -\frac{64\pi^3G_N^3m^4}{3}\frac{147d^6 - 1801d^5 + 8727d^4 - 21555d^3 + 28942d^2 - 20148d + 5688}{3}J_{(3)}(\vec{q}^2), \\ T_{(d)}^{(3)}{}^{\mu\nu}\eta_{\mu\nu} &= -\frac{32\pi^3G_N^3m^4}{3}\frac{179d^5 - 2146d^4 + 10305d^3 - 24614d^2 + 28972d - 13704}{3}J_{(3)}(\vec{q}^2), \end{split}$$

$$T_{(e)}^{(3)\mu\nu}\eta_{\mu\nu} = \frac{64\pi^3 G_N^3 m^4}{3} \frac{29d^4 - 274d^3 + 973d^2 - 1484d + 852}{(d-1)^3} J_{(3)}(\bar{q}^2).$$
(3.50)

3.4.3 The classical three-loop contribution to the stress-tensor

Summing up all the contributions we get for the three-loop stress-tensor

$$\langle T_{\mu\nu}^{(3)} \rangle = \pi^3 G_N^3 m^4 \left(c_1^{(3)}(d) \delta_\mu^0 \delta_\nu^0 + c_2^{(3)}(d) \left(\frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) \right) J_{(3)}(q^2) , \qquad (3.51)$$

with the master integral

$$J_{(3)}(q^2) = \frac{\Gamma\left(\frac{8-3d}{2}\right)\Gamma\left(\frac{d-2}{2}\right)^4}{8^d \pi^{\frac{3d}{2}}\Gamma(2(d-2))} |\vec{q}|^{3(d-2)}, \tag{3.52}$$

and the three-loop coefficients are given by

$$c_1^{(3)}(d) = -\frac{64}{3(d-3)(d-4)^2(d-1)^4} \times \left(56d^7 - 889d^6 + 5868d^5 - 20907d^4 + 43434d^3 - 52498d^2 + 33888d - 8760\right),$$

$$c_2^{(3)}(d) = -\frac{64}{3(d-3)(d-4)^2(d-1)^4} \times \left(45d^7 - 670d^6 + 4167d^5 - 14016d^4 + 27430d^3 - 30916d^2 + 18104d - 3952\right).$$

$$(3.53)$$

Using the relations (3.7) we obtained the three-loop contribution to the metric from the classical stress-tensor in (4.40) (using the notation for ρ in (3.8))

$$h_0^{(4)}(r,d) = \frac{16(d-2)^3(14d^3 - 85d^2 + 165d - 106)}{3(d-3)(d-4)(d-1)^4} \rho(r,d)^4,$$

$$h_1^{(4)}(r,d) = -\frac{8(39d^7 - 691d^6 + 5155d^5 - 21077d^4 + 51216d^3 - 74346d^2 + 60168d - 21208)}{3(d-3)(d-4)^2(d-1)^4(3d-8)} \rho(r,d)^4,$$

$$h_2^{(4)}(r,d) = \frac{16(d-2)^2(45d^6 - 580d^5 + 3007d^4 - 8002d^3 + 11426d^2 - 8064d + 1976)}{3(d-3)(d-4)^2(d-1)^4(3d-8)} \rho(r,d)^4.$$

$$(3.54)$$

4 Non-minimal couplings and renormalised metric

The stress-tensor and the metric components have ultraviolet divergences. These divergences can be removed by the addition of the non-minimal couplings made from the powers of the covariant derivative ∇_{μ} acting on a single power of the Riemann tensor and its contractions. The Bianchi identity on the Riemann tensor $\nabla_{\mu}R_{\nu\rho\sigma\lambda} + \nabla_{\nu}R_{\rho\mu\sigma\lambda} + \nabla_{\rho}R_{\mu\nu\sigma\lambda} = 0$, implies that

$$\nabla_{\mu}R^{\mu}{}_{\rho\sigma\lambda} = \nabla_{\sigma}R_{\rho\lambda} - \nabla_{\lambda}R_{\rho\sigma}, \qquad \nabla_{\mu}R^{\mu}{}_{\nu} = \frac{1}{2}\nabla_{\nu}R. \tag{4.1}$$

The counter-terms are powers of covariant derivative acting on a single power of the Ricci tensor and Ricci scalar. Therefore the counter-terms are given by the following non-minimal couplings

$$\delta^{(n)} S^{\text{ct.}} = (G_N m)^{\frac{2n}{d-2}} \int d^{d+1} x \sqrt{-g} \Big(\alpha^{(n)} (d) (\nabla^2)^{n-1} R \partial_{\mu} \phi \partial^{\mu} \phi + \Big(\beta_0^{(n)} (d) \nabla_{\mu} \nabla_{\nu} (\nabla^2)^{n-2} R + \beta_1^{(n)} (d) (\nabla^2)^{n-1} R_{\mu\nu} \Big) \partial^{\mu} \phi \partial^{\nu} \phi \Big).$$
(4.2)

where $\alpha^{(n)}(d)$, $\beta_0^{(n)}(d)$ and $\beta_1^{(n)}(d)$ are dimensionless coefficients depending on the spacetime dimension. The power of $G_N m$ is determined by dimensional analysis, and give the correct order of G_N in all dimensions. The first non-minimal coupling with n=1 is given by

$$\delta^{(1)}S^{\text{ct.}} = (G_N m)^{\frac{2}{d-2}} \int d^{d+1}x \sqrt{-g} \left(\alpha^{(1)}(d)R\partial_\mu\phi\partial^\mu\phi + \beta^{(1)}(d)R^{\mu\nu}\partial_\mu\phi\partial_\nu\phi\right). \tag{4.3}$$

This non-minimal coupling has been introduced in [27] in four dimensions and [24] in five dimensions. We will see that up to three-loop order the renormalisation of the static metric component only require the counter-term $\alpha^{(1)}(d)R\partial_{\mu}\phi\partial^{\mu}\phi$, whereas both couplings are needed for the cancellation of the stress-tensor divergences. This coupling is induced by harmonic gauge condition [20, 24] and the value of its coefficient depends on the choice of gauge. In our gauge, the de Donder gauge, this corresponds to $\alpha = 0$ in the work of [24] and $\xi = \frac{1}{4}$ in the work of [20]. Since we are working in fixed gauge we will not discuss further the gauge dependence of the higher-order non-minimal coupling coefficients, but we expect that the gauge dependence of these coefficients will be an extension of the discussion in [24, app. B].

The power of the Newton constant in (4.3) is an integer only in four dimensions with d=3 and five dimensions d=4. Therefore this counter-term will not appear in dimensions $D \ge 6$.

In four dimensions, from five-loop order, or the sixth post-Minkowskian order $O(G_N^6)$, one expects that higher derivative non-minimal couplings will be needed to get finite stress-tensor components. In dimensions five and six, the higher-derivative non-minimal couplings arise at lower loop order.

In five dimensions one needs to consider higher-derivative non-minimal couplings $\delta^{(n)}S^{\text{ct.}}$ with $n \geq 2$ for removing the divergences in the stress-tensor. The non-minimal coupling at this order is then given by

$$\delta^{(2)}S^{ct.} = (G_N m)^{\frac{4}{d-2}} \int d^{d+1}x \sqrt{-g} \Big(\alpha^{(2)}(d) \square R \partial_\mu \phi \partial^\mu \phi + \Big(\beta_0^{(2)}(d) \nabla_\mu \nabla_\nu R + \beta_1^{(2)}(d) \square R_{\mu\nu} \Big) \partial^\mu \phi \partial^\nu \phi \Big). \quad (4.4)$$

We will need the non-minimal coupling

$$\delta^{(3)}S^{ct.} = (G_N m)^{\frac{6}{d-2}} \int d^{d+1}x \sqrt{-g} \Big(\alpha^{(3)}(d)(\nabla^2)^2 R \partial_\mu \phi \partial^\mu \phi + \Big(\beta_0^{(3)}(d) \nabla_\mu \nabla_\nu \nabla^2 R + \beta_1^{(3)}(d)(\nabla^2)^2 R_{\mu\nu} \Big) \partial^\mu \phi \partial^\nu \phi \Big), \quad (4.5)$$

for removing the two-loop divergence in the stress-tensor in six (d = 5) dimensions and the three-loop divergence in five (d = 4) dimensions. In five dimensions (d = 4) the metric,

up to G_N^4 , is renormalised using only the n=1 and the metric is finite to all order in six dimensions (d=5).

The higher-order non-minimal couplings $\delta^{(n)}S^{\text{ct.}}$ with $n \geq 2$ will not contribute to the classical limit when inserted into graphs with loops, because they contribute to higher powers in the momentum transfer \vec{q} , and are sub-leading with respect to the classical contributions. Their tree-level insertions will contribute to the renormalisation of the stress-tensor but thanks to the properties of the Fourier transform they will not contribute to the metric components.

4.1 Tree-level insertions

We give the contribution of the insertions of the non-minimal counter-terms with n = 1 in (4.3), with n = 2 in (4.4) and with n = 3 in (4.5) in the tree-level graph.

4.1.1 Insertion of $\delta^{(1)}S^{\text{ct.}}$

The insertion of the non-minimal couplings $\delta^{(1)}S^{\text{ct.}}$ in (4.3) into the tree-level diagram

$$\delta^{(1)}\mathcal{M}^{(0)}(p_1,q) = \begin{array}{c} p_2 \\ p_1 \\ p_1 \end{array}$$
(4.6)

leads to the stress-tensor contribution in d+1 dimensions

$$\delta^{(1)}\langle T_{\mu\nu}^{(0)}\rangle = -\bar{q}^2 (G_N m)^{\frac{2}{d-2}} m \left(-\beta^{(1)}(d) \delta_{\mu}^0 \delta_{\nu}^0 + 2\alpha^{(1)}(d) \left(\frac{q_{\mu} q_{\nu}}{q^2} - \eta_{\mu\nu}\right)\right), \tag{4.7}$$

and using (2.6) this contributes to the metric components

$$\delta^{(1)} h_0^{(1)}(r,d) = 0, \tag{4.8}$$

$$\delta^{(1)} h_1^{(1)}(r,d) = \frac{16\alpha^{(1)}(d)\Gamma\left(\frac{d}{2}\right)}{\pi^{\frac{d-2}{2}}} \left(\frac{(G_N m)^{\frac{1}{d-2}}}{r}\right)^d,$$

$$\delta^{(1)}h_2^{(1)}(r,d) = -\frac{32\alpha^{(1)}(d)\Gamma\left(\frac{d+2}{2}\right)}{\pi^{\frac{d-2}{2}}} \left(\frac{(G_N m)^{\frac{1}{d-2}}}{r}\right)^d. \tag{4.9}$$

Thanks to the properties of the Fourier transformation (see appendix A) only the coefficient $\alpha(d)$ contributes to static metric perturbation.

4.1.2 Insertion of $\delta^{(2)}S^{\text{ct.}}$

The insertion of the non-minimal couplings $\delta^{(2)}S^{\text{ct.}}$ in (4.4) into the tree-level diagram

$$\delta^{(2)}\mathcal{M}^{(0)}(p_1,q) = \begin{cases} p_2 \\ p_2 \\ p_3 \end{cases}$$

$$(4.10)$$

leads to the stress-tensor condition in d+1 dimensions

$$\delta^{(2)}\langle T_{\mu\nu}^{(0)}\rangle = |\vec{q}|^4 (G_N m)^{\frac{4}{d-2}} m \left(-\beta_1^{(2)}(d) \delta_\mu^0 \delta_\nu^0 + 2 \left(\alpha^{(2)}(d) + \frac{1}{2} \beta_0^{(2)}(d) \right) \left(\frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) \right). \tag{4.11}$$

Because of the vanishing of the Fourier transforms

$$\int_{\mathbb{R}^d} |\vec{q}|^2 e^{i\vec{q}\cdot\vec{x}} \frac{d^d\vec{q}}{(2\pi)^d} = 0, \qquad \int_{\mathbb{R}^d} \frac{q_i q_j}{|\vec{q}|^2} |\vec{q}|^2 e^{i\vec{q}\cdot\vec{x}} \frac{d^d\vec{q}}{(2\pi)^d} = 0, \tag{4.12}$$

this extra contribution to the stress-tensor does not affect the metric components

$$\delta^{(2)}h_0^{(1)}(r,d) = 0, (4.13)$$

$$\delta^{(2)}h_1^{(1)}(r,d) = 0.$$

$$\delta^{(2)}h_2^{(1)}(r,d) = 0. (4.14)$$

4.1.3 Insertion of $\delta^{(3)}S^{\text{ct.}}$

The insertion of the non-minimal couplings $\delta^{(3)}S^{\text{ct.}}$ in (4.5) into the tree-level diagram

$$\delta^{(3)}\mathcal{M}^{(0)}(p_1,q) = \begin{cases} p_2 \\ q \\ p_3 \end{cases} \tag{4.15}$$

leads to the stress-tensor condition in six dimensions (d = 5)

$$\delta^{(3)} \langle T_{\mu\nu}^{(0)} \rangle = -|\vec{q}|^6 (G_N m)^{\frac{6}{d-2}} m \left(-\beta_1^{(3)}(d) \delta_\mu^0 \delta_\nu^0 + 2 \left(\alpha^{(3)}(d) + \frac{1}{4} \beta_0^{(3)}(d) \right) \left(\frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) \right). \tag{4.16}$$

Because of the vanishing of the Fourier transforms

$$\int_{\mathbb{R}^d} |\vec{q}|^4 e^{i\vec{q}\cdot\vec{x}} \frac{d^d\vec{q}}{(2\pi)^d} = 0, \qquad \int_{\mathbb{R}^d} \frac{q_i q_j}{|\vec{q}|^2} |\vec{q}|^4 e^{i\vec{q}\cdot\vec{x}} \frac{d^d\vec{q}}{(2\pi)^d} = 0, \tag{4.17}$$

this extra contribution to the stress-tensor does not affect the metric components

$$\delta^{(2)}h_0^{(1)}(r,d) = 0, (4.18)$$

$$\delta^{(2)}h_1^{(1)}(r,d) = 0,$$


$$\delta^{(2)}h_2^{(1)}(r,d) = 0. (4.19)$$

4.2 One-loop insertions

We give the contribution of the insertions of the counter-terms (4.2) with n = 1 in (4.3) in the one-loop graph.

4.2.1 Insertion of $\delta^{(1)}S^{\text{ct.}}$

The insertion of the non-minimal coupling in (4.3) in the one-loop graph



leads to the stress-tensor contribution

$$\delta^{(1)} \langle T_{\mu\nu}^{(1)} \rangle = 32i\alpha^{(1)}(d)\pi (G_N m)^{\frac{d}{d-2}} m^2$$

$$\times \int \frac{d^d \vec{l}}{(2\pi)^d} \frac{\tau_{10\ \alpha\beta,\gamma\delta}^{\mu\nu}(l,q) \mathcal{P}_{00}^{\alpha\beta} l^{\gamma} l^{\delta}}{l^2 (l+q)^2} \left[\frac{1}{(l+p_1)^2 - m^2 + i\epsilon} + \frac{1}{(l-p_2)^2 - m^2 + i\epsilon} \right]$$

$$= 8\pi \alpha^{(1)}(d) (G_N m)^{\frac{d}{d-2}} m \vec{q}^2 \frac{d-2}{(d-1)^2} \left(d\delta_{\mu}^0 \delta_{\nu}^0 + \frac{q_{\mu} q_{\nu}}{a^2} - \eta_{\mu\nu} \right) J_{(1)}(\vec{q}^2) . \tag{4.21}$$

where we used that

$$\eta_{\mu\nu}\tau_{10\ 00,\gamma\delta}^{\mu\nu}(l,q)l^{\gamma}l^{\delta} = \frac{\vec{l_1}^2}{2} \left(\vec{q}^2 + (\vec{l_1} + \vec{q})^2 - \vec{l_1}^2\right),\tag{4.22}$$

and

$$\delta^{0}_{\mu}\delta^{0}_{\nu}\tau^{\mu\nu}_{10\ 00,\gamma\delta}(l,q)l^{\gamma}l^{\delta} = \frac{1}{2(d-1)} \left((d-2)\vec{q}^{4} + (d-2)(\vec{l_{1}} + \vec{q})^{2} \left((\vec{l_{1}} + \vec{q})^{2} - 2\vec{q}^{2} \right) - \vec{l_{1}}^{4} - (d-3)\vec{l_{1}}^{2} \left((\vec{l_{1}} + \vec{q})^{2} + \vec{q}^{2} \right) \right). \tag{4.23}$$

Using the Fourier transforms

$$\int_{\mathbb{R}^d} J_{(1)}(\vec{q}^2) e^{i\vec{q}\cdot\vec{x}} \frac{d^d \vec{q}}{(2\pi)^d} = -\frac{\Gamma\left(\frac{d}{2}\right)^2}{2\pi^d r^{2(d-1)}},$$

$$\int_{\mathbb{R}^d} \frac{q_i q_j}{\vec{q}^2} J_{(1)}(\vec{q}^2) e^{i\vec{q} \cdot \vec{x}} \frac{d^d \vec{q}}{(2\pi)^d} = \frac{\Gamma\left(\frac{d-2}{2}\right) \Gamma\left(\frac{d}{2}\right)}{4\pi^d r^{2(d-1)}} \left(\delta_{ij} - 2(d-1)\frac{x_i x_j}{r}\right) . \tag{4.24}$$

and the relation between the stress-tensor and the metric components in (2.6) we obtain the following contribution to the metric components

$$\delta^{(1)}h_0^{(2)}(r,d) = 64\alpha^{(1)}(d)\frac{(d-2)\Gamma(\frac{d}{2})^2}{(d-1)\pi^{d-2}} \left(\frac{(G_N m)^{\frac{1}{d-2}}}{r}\right)^{2(d-1)},$$


$$\delta^{(1)}h_1^{(2)}(r,d) = -64\alpha^{(1)}(d)\frac{\Gamma(\frac{d}{2})^2}{(d-1)\pi^{d-2}} \left(\frac{(G_N m)^{\frac{1}{d-2}}}{r}\right)^{2(d-1)},$$

$$\delta^{(1)}h_2^{(2)}(r,d) = 128\alpha^{(1)}(d)\frac{\Gamma(\frac{d}{2})^2}{(d-1)\pi^{d-2}} \left(\frac{(G_N m)^{\frac{1}{d-2}}}{r}\right)^{2(d-1)}.$$

$$(4.25)$$

4.2.2 Two insertions of $\delta^{(1)}S^{\text{ct.}}$

Two insertions of the non-minimal coupling $\delta^{(1)}S^{\text{ct.}}$ in (4.3) in the one-loop graph



leads to the stress-tensor contribution

$$(\delta^{1})^{2} \langle T_{\mu\nu}^{(1)} \rangle = \frac{2(\alpha^{(1)}(d))^{2} (G_{N}m)^{\frac{d+2}{d-2}} \pi m \vec{q}^{4}}{d-1} \left(\delta_{\mu}^{0} \delta_{\nu}^{0} - (d-2) \left(\frac{q_{\mu} q_{\nu}}{q^{2}} - \eta_{\mu\nu} \right) \right) J_{(1)}(\vec{q}^{2}), \quad (4.27)$$

and the metric contributions

$$(\delta^{1})^{2}h_{0}^{(2)}(r,d) = 0,$$

$$(\delta^{1})^{2}h_{1}^{(2)}(r,d) = \frac{64(\alpha^{(1)}(d))^{2}}{\pi^{d-2}}\Gamma\left(\frac{d}{2}\right)^{2}\left(\frac{(G_{N}m)^{\frac{1}{d-2}}}{r}\right)^{2d},$$

$$(\delta^{1})^{2}h_{2}^{(2)} = \frac{64d(d-2)(\alpha^{(1)}(d))^{2}}{\pi^{d-2}}\Gamma\left(\frac{d}{2}\right)^{2}\left(\frac{(G_{N}m)^{\frac{1}{d-2}}}{r}\right)^{2d}.$$

$$(4.28)$$

4.3 Two-loop insertions

For the insertion of the non-minimal coupling $\delta^{(1)}S^{\text{ct.}}$ in (4.3) in the two-loop graph one needs to sum over all the contributions in table 1. The classical limit of the sum of all these graphs lead to the following contribution to the stress-tensor




Table 1. Insertion of the non-minimal coupling in the two-loop graph

$$\delta^{(1)}\langle T_{\mu\nu}^{(2)}\rangle = -\frac{128\pi^2(d-2)\alpha^{(1)}(d)}{3(d-4)(3d-4)(d-1)^2}(G_N m)^{\frac{2(d-1)}{d-2}} m \bar{q}^2 \left((3d^3 - 19d^2 + 28d - 10)\delta_{\mu}^0 \delta_{\nu}^0 + (3d^3 - 15d^2 + 18d - 4)\left(\frac{q_{\mu}q_{\nu}}{q^2} - \eta_{\mu\nu}\right) \right) J_{(2)}(\bar{q}^2), \quad (4.29)$$

which leads to the following contributions to the metric components

$$\delta^{(1)}h_0^{(3)}(r,d) = -\frac{512\alpha^{(1)}(d)}{d-1} \frac{\Gamma(\frac{d}{2})^3}{\pi^{\frac{3}{2}(d-2)}} \left(\frac{(G_N m)^{\frac{1}{d-2}}}{r}\right)^{3d-4}, \tag{4.30}$$

$$\delta^{(1)}h_1^{(3)}(r,d) = \frac{256\alpha^{(1)}(d)\left(3d^3 - 23d^2 + 46d - 28\right)}{(d-4)(d-2)(d-1)^2(3d-4)} \frac{\Gamma(\frac{d}{2})^3}{\pi^{\frac{3}{2}(d-2)}} \left(\frac{(G_N m)^{\frac{1}{d-2}}}{r}\right)^{3d-4},$$

$$\delta^{(1)}h_2^{(3)}(r,d) = -\frac{256\alpha^{(1)}(d)\left(3d^3 - 15d^2 + 18d - 4\right)}{(d-4)(d-2)(d-1)^2} \frac{\Gamma(\frac{d}{2})^3}{\pi^{\frac{3}{2}(d-2)}} \left(\frac{(G_N m)^{\frac{1}{d-2}}}{r}\right)^{3d-4}.$$

4.4 The renormalised metric in four dimensions

The metric components have ultraviolet poles in four dimensions from two-loop order. We show how the addition of the non-minimal couplings leads to finite renormalised metric components.

4.4.1 The two-loop renormalisation

The two-loop metric components in (3.44) have a divergence in four dimensions (d=3)

$$\begin{split} h_0^{(3)}(r,d) &= O(1), \\ h_1^{(3)}(r,d) &= -\frac{2}{3(d-3)} \left(\frac{G_N m}{r}\right)^3 + O(1), \end{split}$$

$$h_2^{(3)}(r,d) = \frac{2}{d-3} \left(\frac{G_N m}{r}\right)^3 + O(1).$$
 (4.31)

This divergence is cancelled by adding the metric contribution from the non-minimal coupling in (4.8)

$$h_i^{\text{renor. (3)}}(r,d) := h_i^{(3)}(r,d) + \delta^{(1)}h_i^{(1)}(r,d), \qquad i = 0, 1, 2$$
 (4.32)

and setting the $\alpha^{(1)}(d)$ coefficient to be

$$\alpha^{(1)}(d) = \frac{1}{12(d-3)} + a^{(1)}(3) - \frac{\log(2)}{6} + O(d-3). \tag{4.33}$$

The resulting renormalised two-loop metric reads

$$h_0^{\text{renor. (3)}}(r,d) = 2\left(\frac{G_N m}{r}\right)^3 + O(d-3),$$

$$h_1^{\text{renor. (3)}}(r,d) = \frac{4}{3}\left(-\frac{1}{2} + 6a^{(1)}(3) + \log\left(\frac{rC_E}{G_N m}\right)\right)\left(\frac{G_N m}{r}\right)^3 + O(d-3),$$

$$h_2^{\text{renor. (3)}}(r,d) = 4\left(\frac{1}{3} - 6a^{(1)}(3) - \log\left(\frac{rC_E}{G_N m}\right)\right)\left(\frac{G_N m}{r}\right)^3 + O(d-3). \tag{4.34}$$

where we have introduced the following combination of the Euler-Mascheroni constant [38] and π

$$C_E := \sqrt{\pi}e^{\frac{\gamma_E}{2}}. (4.35)$$

The divergence in the two-loop stress-tensor in (3.41)

$$\langle T_{\mu\nu}^{(2)} \rangle = \frac{G_N^2 \vec{q}^2 m^3}{6(d-3)} \left(2\delta_{\mu}^0 \delta_{\nu}^0 + \left(\frac{q_{\mu}q_{\nu}}{q^2} - \eta_{\mu\nu} \right) \right) + O(1) , \qquad (4.36)$$

is cancelled by adding the contribution in (4.7) from the non-minimal coupling with the following choice of $\beta^{(1)}(d)$ coefficient

$$\beta^{(1)}(d) = -\frac{1}{3(d-3)} + O(1). \tag{4.37}$$

Notice that this computation does not determine the finite part of the $\alpha^{(1)}(d)$ and $\beta^{(1)}(d)$. They are free scales in the logarithms. We will show in section 6 that this freedom is totally reabsorbed in the change of coordinate and the Schwarzschild-Tangherlini metric does not have any ambiguity.

4.4.2 The three-loop renormalisation

The three-loop metric components in (3.54) have a divergence in four dimensions (d = 3) given by

$$h_0^{(4)}(r,d) = -\frac{2}{3(d-3)} \left(\frac{G_N m}{r}\right)^4 + O(1),$$

$$h_1^{(4)}(r,d) = \frac{2}{3(d-3)} \left(\frac{G_N m}{r}\right)^4 + O(1),$$

$$h_2^{(4)}(r,d) = -\frac{4}{3(d-3)} \left(\frac{G_N m}{r}\right)^4 + O(1),$$
(4.38)

Adding to this contribution the (4.25) from the insertion of the non-minimal couplings at one-loop, and using the value of $\alpha^{(1)}(d)$ determined in (4.33), we obtain the renormalised three-loop metric

$$h_0^{\text{renorm.(4)}}(r) = \left(-\frac{32}{3} + 8a^{(1)}(3) + \frac{4}{3}\log\left(\frac{rC_E}{G_N m}\right)\right) \left(\frac{G_N m}{r}\right)^4 + O(d-3),$$

$$h_1^{\text{renorm.(4)}}(r) = \left(10 - 8a^{(1)}(3) - \frac{4}{3}\log\left(\frac{rC_E}{G_N m}\right)\right) \left(\frac{G_N m}{r}\right)^4 + O(d-3),$$

$$h_2^{\text{renorm.(4)}}(r) = \left(-\frac{86}{3} + 16a^{(1)}(3) + \frac{8}{3}\log\left(\frac{rC_E}{G_N m}\right)\right) \left(\frac{G_N m}{r}\right)^4 + O(d-3). \tag{4.39}$$

The classical three-loop contribution to the stress-tensor has an ultraviolet divergence

$$\langle T_{\mu\nu}^{(3)}(\vec{q})\rangle = -\frac{\pi G_N^3 m^4 |\vec{q}|^{\frac{3}{2}}}{48(d-3)} \left(3\delta_{\mu}^0 \delta_{\nu}^0 + \left(\frac{q_{\mu}q_{\nu}}{q^2} - \eta_{\mu\nu} \right) \right) + O(1), \qquad (4.40)$$

this divergence is cancelled by the addition of the contribution in (4.21) from the non-minimal coupling and the choice of $\alpha^{(1)}(d)$ in (4.33).

4.5 The renormalised metric in five dimensions

The metric components have ultraviolet divergences in five dimensions from one-loop order. We show how the addition of the non-minimal couplings leads to finite renormalised metric components.

4.5.1 The one-loop renormalisation

The metric components in (3.26) have a divergence in five dimension (d=4) given by

$$h_0^{(2)}(r,d) = O(1),$$

$$h_1^{(2)}(r,d) = -\frac{40}{9(d-4)} \left(\frac{G_N m}{\pi r^2}\right)^2 + O(1),$$

$$h_2^{(2)}(r,d) = \frac{160}{9(d-4)} \left(\frac{G_N m}{\pi r^2}\right)^2 + O(1).$$
(4.41)

The divergences in the metric components (4.41) are cancelled for the choice

$$\alpha^{(1)}(d) = \frac{5}{18\pi(d-4)} + a^{(1)}(5) + O(d-4), \qquad (4.42)$$

so that the renormalised metric components

$$h_i^{\text{renor. (2)}}(r,d) := h_i^{(2)}(r,d) + \delta^{(1)}h_i^{(1)}(r,d), \qquad i = 0, 1, 2,$$
 (4.43)

have a finite expansion near d=4

$$h_0^{\text{renor. (2)}}(r,d) = \frac{32}{9} \left(\frac{G_N m}{\pi r^2}\right)^2 + O(d-4), \tag{4.44}$$

$$h_1^{\text{renor. (2)}}(r,d) = \frac{20}{9} \left(\frac{14}{15} + \frac{36a^{(1)}(5)\pi}{5} + \log\left(\frac{r^2 C_E^2}{G_N m}\right)\right) \left(\frac{G_N m}{\pi r^2}\right)^2 + O(d-4),$$

$$h_2^{\text{renor(2)}}(r,d) = -\frac{80}{9} \left(\frac{7}{30} + \frac{36a^{(1)}(5)\pi}{5} + \log\left(\frac{r^2 C_E^2}{G_N m}\right)\right) \left(\frac{G_N m}{\pi r^2}\right)^2 + O(d-4).$$

where C_E is defined in (4.35).

Thanks to the properties of the Fourier transform, only the coefficient $\alpha^{(1)}(d)$ enters the counter-term contribution to the metric component. To determine as well the coefficient $\beta^{(1)}(d)$ in (4.3) one needs to look at the divergences of the stress-tensor

$$\langle T_{\mu\nu}^{(1)} \rangle = \frac{G_N m^2 \bar{q}^2}{18\pi (d-4)} \left(7\delta_{\mu}^0 \delta_{\nu}^0 + 10 \left(\frac{q_{\mu} q_{\nu}}{q^2} - \eta_{\mu\nu} \right) \right) + O(1)$$
 (4.45)

The cancellation of the pole fixes the pole part of $\beta^{(1)}(d)$ near five dimensions

$$\beta^{(1)}(d) = -\frac{7}{18\pi(d-4)} + O(1). \tag{4.46}$$

4.5.2 The two-loop renormalisation

The two-loop metric components in (3.44) have a divergence in five dimensions (d=4)

$$h_0^{(3)}(r,d) = -\frac{320}{27(d-4)} \left(\frac{G_N m}{\pi r^2}\right)^3 + O(1),$$

$$h_1^{(3)}(r,d) = \frac{160}{27(d-4)} \left(\frac{G_N m}{\pi r^2}\right)^3 + O(1),$$

$$h_2^{(3)}(r,d) = -\frac{320}{27(d-4)} \left(\frac{G_N m}{\pi r^2}\right)^3 + O(1).$$
(4.47)

The divergences in the metric components (3.44) are cancelled for the choice made at one-loop in (4.42), so that the renormalised metric components

$$h_i^{\text{renor. (3)}}(r,d) := h_i^{(3)}(r,d) + \delta^{(1)}h_i^{(2)}(r,d), \qquad i = 0, 1, 2,$$
 (4.48)

have a finite expansion near d=4

$$h_0^{\text{renor. (3)}}(r,d) = \frac{160}{27} \left(\frac{2}{15} + \frac{36a^{(1)}(5)\pi}{5} + \log\left(\frac{r^2C_E^2}{G_Nm}\right) \right) \left(\frac{G_Nm}{\pi r^2}\right)^3 + O(d-4),$$

$$h_1^{\text{renor. (3)}}(r,d) = -\frac{80}{27} \left(\frac{7}{15} + \frac{36a^{(1)}(5)\pi}{5} + \log\left(\frac{r^2C_E^2}{G_Nm}\right) \right) \left(\frac{G_Nm}{\pi r^2}\right)^3 + O(d-4),$$

$$h_2^{\text{renor(3)}}(r,d) = \frac{160}{27} \left(-\frac{1}{15} + \frac{36a^{(1)}(5)\pi}{5} + \log\left(\frac{r^2C_E^2}{G_Nm}\right) \right) \left(\frac{G_Nm}{\pi r^2}\right)^3 + O(d-4). (4.49)$$

The two-loop stress-tensor in (3.41) is not finite in d=4 as it diverges like

$$\langle T_{\mu\nu}^{(2)} \rangle = \frac{5G_N^2 m^3 |\vec{q}|^4}{162\pi^2 (d-4)^2} \left(4\delta_\mu^0 \delta_\nu^0 + \frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) + \frac{5G^2 m^3 |\vec{q}|^4}{162\pi^2 (d-4)} \left(\left(4\log\left(\frac{\vec{q}^2}{4\pi}\right) + 4\gamma_E - \frac{183}{20} \right) \delta_\mu^0 \delta_\nu^0 + \left(\log\left(\frac{\vec{q}^2}{4\pi}\right) + \gamma_E - \frac{41}{20} \right) \left(\frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) \right) + O(1) \,. \tag{4.50}$$

The addition of the counter-term in (4.21) from the non-minimal couplings in (4.3) is not enough for making the stress-tensor finite in d=4

$$\langle T_{\mu\nu}^{(2)} \rangle + \delta^{(1)} \langle T_{\mu\nu}^{(1)} \rangle = -\frac{5G_N^2 m^3 |\vec{q}|^4}{162\pi^2 (d-4)^2} \left(4\delta_\mu^0 \delta_\nu^0 + \frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) + \frac{5G^2 m^3 |\vec{q}|^4}{162\pi^2 (d-4)} \left(\left(4\log \left(G_N m \right) - \frac{144\pi a^{(1)}(5)}{5} - \frac{109}{60} \right) \delta_\mu^0 \delta_\nu^0 \right. + \left. \left(\left(\log \left(G_N m \right) + \frac{17}{60} - \frac{36}{5}\pi a^{(1)}(5) \right) \left(\frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) \right) \right) + O(1) . \quad (4.51)$$

We need to consider the addition of the counter-term from the insertion of $\delta^{(2)}S^{\text{ct.}}$ evaluated in section 4.1.2 with the values of the coefficient near d=4

$$\beta_1^{(2)}(d) = \frac{1}{\pi^2} \left(\frac{10}{81(d-4)^2} + \frac{109 + 1728\pi a^{(1)}(5)}{1944(d-4)} + a^{(2)}(5) + O(d-4) \right),$$

$$\alpha^{(2)}(d) + \frac{1}{2} \beta_0^{(2)}(d) = -\frac{1}{2\pi^2} \left(\frac{5}{162(d-4)^2} + \frac{432\pi a^{(1)}(5) - 17}{1944(d-4)} + b^{(2)}(5) + O(d-4) \right),$$

$$(4.52)$$

plugged in (4.11) cancel the divergences in (4.51)

$$\langle T_{\mu\nu}^{(2)} \rangle + \delta^{(1)} \langle T_{\mu\nu}^{(1)} \rangle + \delta^{(2)} \langle T_{\mu\nu}^{(0)} \rangle = O(1).$$
 (4.53)

4.5.3 The three-loop renormalisation

The three-loop metric components in (3.54) have a divergence in five dimensions (d=4)

$$h_0^{(4)}(r,d) = \frac{1280}{27(d-4)} \left(\frac{G_N m}{\pi r^2}\right)^4 + O(1),$$

$$h_1^{(4)}(r,d) = \left(\frac{400}{81(d-4)^2} - \frac{20\left(101 + 120\log\left(r^2 C_E^2\right)\right)}{243(d-4)}\right) \left(\frac{G_N m}{\pi r^2}\right)^4 + O(1),$$

$$h_2^{(4)}(r,d) = \left(\frac{3200}{81(d-4)^2} + \frac{160\left(187 - 120\log\left(r^2 C_E^2\right)\right)}{243}\right) \left(\frac{G_N m}{\pi r^2}\right)^4 + O(1). \quad (4.54)$$

The divergences in the metric components (3.54) are cancelled for the choice made at one-loop in (4.42), so that the renormalised metric components

$$h_i^{\text{renor. }(4)}(r,d) := h_i^{(4)}(r,d) + \delta^{(1)}h_i^{(3)}(r,d) + (\delta^1)^2h_i^{(2)}(r,d), \qquad i = 0,1,2\,, \tag{4.55}$$

have a finite expansion near d=4

$$h_0^{\text{renor. (4)}}(r,d) = -\frac{128}{243} \left(23 + 324a^{(1)}(5)\pi + 45\log\left(\frac{r^2C_E^2}{G_Nm}\right) \right) \left(\frac{G_Nm}{\pi r^2}\right)^4 + O(d-4),$$

$$h_1^{\text{renor. (4)}}(r,d) = \frac{100}{81} \left(\left(\frac{36a^{(1)}(5)\pi}{5} + \log\left(\frac{r^2C_E^2}{G_Nm}\right) \right) \left(\frac{161}{30} + \frac{36}{5}a^{(1)}(5)\pi + \log\left(\frac{r^2C_E^2}{G_Nm}\right) \right) + \frac{7085}{1800} \right) \times \left(\frac{G_Nm}{\pi r^2}\right)^4 + O(d-4),$$

$$h_2^{\text{renor(4)}}(r,d) = -\frac{800}{81} \left(\left(\frac{36a^{(1)}(5)\pi}{5} + \log\left(\frac{r^2C_E^2}{G_Nm}\right) \right) \left(\frac{41}{15} - \frac{36}{5}a^{(1)}(5)\pi - \log\left(\frac{r^2C_E^2}{G_Nm}\right) \right) + \frac{2381}{900} \right) \times \left(\frac{G_Nm}{\pi r^2}\right)^4 + O(d-4).$$

$$(4.56)$$

The three-loop stress-tensor in (3.51) is not finite in d=4 as it diverges like

$$\begin{split} \langle T_{\mu\nu}^{(3)} \rangle &= \frac{25 G_N^3 m^4 |\vec{q}|^6}{5832 \pi^3 (d-4)^3} \left(-\frac{1}{2} \delta_\mu^0 \delta_\nu^0 + \frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) \\ &\quad + \frac{25 G_N^3 m^4 |\vec{q}|^6}{3888 \pi^3 (d-4)^2} \left(-\frac{1}{2} \left(\log \left(\frac{\vec{q}^2}{4\pi} \right) + \gamma_E - \frac{41}{6} \right) \delta_\mu^0 \delta_\nu^0 \right. \\ &\quad + \left(\log \left(\frac{\vec{q}^2}{4\pi} \right) + \gamma_E - \frac{17}{10} \right) \left(\frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) \right) \\ &\quad + \frac{225 G_N^3 m^4 |\vec{q}|^6}{839808 \pi^3 (d-4)} \left(\frac{1}{2} \left(\frac{70939}{450} + \pi^2 - 18 \left(\log \left(\frac{\vec{q}^2}{4\pi} \right) + \gamma_E - \frac{41}{6} \right)^2 \right) \delta_\mu^0 \delta_\nu^0 \\ &\quad + \left(\frac{4769}{450} - \pi^2 + 18 \left(\log \left(\frac{\vec{q}^2}{4\pi} \right) + \gamma_E - \frac{17}{10} \right)^2 \right) \left(\frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) \right) + O(1) \,. \end{split}$$

The addition of the counter-terms in $(\delta^1)^2 \langle T_{\mu\nu}^{(1)} \rangle$ in (4.27), and $\delta^{(1)} \langle T_{\mu\nu}^{(2)} \rangle$ in (4.29) from the non-minimal couplings in (4.3) is not enough for making the stress-tensor finite in d=4

$$\begin{split} \langle T_{\mu\nu}^{(3)} \rangle + (\delta^1)^2 \langle T_{\mu\nu}^{(1)} \rangle + \delta^{(1)} \langle T_{\mu\nu}^{(2)} \rangle &= \frac{25 G_N^3 m^4 |\vec{q}|^6}{5832 \pi^3 (d-4)^3} \left(-\frac{1}{2} \delta_\mu^0 \delta_\nu^0 + \frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) \\ &+ \frac{25 G_N^3 m^4 |\vec{q}|^6}{3888 \pi^3 (d-4)^2} \left(-\frac{1}{2} \left(\frac{25}{12} + \frac{36}{5} a^{(1)} (5) \pi - \log \left(G_N m \right) \right) \delta_\mu^0 \delta_\nu^0 \right. \\ &+ \left(\frac{1}{60} + \frac{36}{5} a^{(1)} (5) \pi - \log \left(G_N m \right) \right) \left(\frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu} \right) \right) \\ &- \frac{25 G_N^3 m^4 |\vec{q}|^6}{5184 \pi^3 (d-4)} \times \\ &\times \left(\left(\frac{27487}{48600} + a^{(1)} (5) \pi \left(1 + \frac{288}{25} a^{(1)} (5) \pi \right) - \frac{\log \left(G_N m \right)}{3} \left(\frac{7}{2} + \log \left(G_N m \right) + \frac{72}{5} a^{(1)} (5) \pi \right) \right) \delta_\mu^0 \delta_\nu^0 \end{split}$$

$$+\left(\frac{6749}{16200} - \frac{6a^{(1)}(5)\pi}{25}\left(1 + 144a^{(1)}(5)\pi\right) - \log\left(G_N m\right)\left(\frac{19}{30} + \log\left(G_N m\right) - \frac{72}{5}a^{(1)}(5)\pi\right)\right)\left(\frac{q_\mu q_\nu}{q^2} - \eta_{\mu\nu}\right)\right) + O(1). \quad (4.58)$$

We need to consider the addition of the counter-term from the insertion of $\delta^{(3)}S^{\text{ct.}}$ evaluated in section 4.1.3 with the values of the coefficient near d=4

$$\beta_1^{(3)}(d) = \frac{25}{11664\pi^3(d-4)^3} + \frac{5(432\pi a^{(1)}(5) + 125)}{93312\pi^3(d-4)^2} + \frac{559872(\pi a^{(1)}(5))^2 + 486000\pi a^{(1)}(5) + 27487}{6718464\pi^3(d-4)} + O(1),$$

$$\alpha^{(3)}(d) + \frac{1}{4}\beta_0^{(3)}(d) = \frac{25}{11664\pi^3(d-4)^3} + \frac{2160\pi a^{(1)}(5) + 5}{93312\pi^3(d-4)^2} + \frac{559872(\pi a^{(1)}(5))^2 + 3888\pi a^{(1)}(5) - 6749}{6718464\pi^3(d-4)} + O(1),$$

$$(4.59)$$

plugged in (4.11) cancel the divergences in (4.51)

$$\langle T_{\mu\nu}^{(2)} \rangle + \delta^{(1)} \langle T_{\mu\nu}^{(2)} \rangle + (\delta^{(1)} \langle T_{\mu\nu}^{(2)} \rangle + \delta^{(3)} \langle T_{\mu\nu}^{(0)} \rangle = O(1). \tag{4.60}$$

4.6 The renormalised stress-tensor in six dimensions

In six dimensions, the metric component are finite to all order in perturbation but the two-loop stress-tensor in (3.41) presents an ultraviolet divergence in six dimensions (d = 5)

$$\langle T_{\mu\nu}^{(2)} \rangle = -\frac{G_N^2 m^3 |\vec{q}|^6}{40320\pi^2 (d-5)} \left(49 \delta_{\mu}^0 \delta_{\nu}^0 + 15 \left(\frac{q_{\mu} q_{\nu}}{q^2} - \eta_{\mu\nu} \right) \right) + O(1) , \qquad (4.61)$$

which is cancelled by the addition of the insertion of the non-minimal coupling $\delta^{(3)}S^{\text{ct.}}$ at tree-level in (4.16) with the choice of the coefficients

$$\alpha^{(3)}(d) + \frac{1}{4}\beta_0^{(3)}(d) = -\frac{15}{80640\pi^2(d-5)} + O(1),$$

$$\beta_1^{(3)}(d) = -\frac{49}{40320\pi^2(d-5)} + O(1). \tag{4.62}$$

5 The Schwarzschild-Tangherlini metric in de Donder gauge in four, five and six dimensions

The Schwarzschild-Tangherlini [40] space-time metric in d+1 dimensions is given by the Tangherlini solution, using $\rho(r,d)$ defined in (3.8),⁷

$$ds_{\text{Schw}}^2 = \left(1 - 4\frac{d-2}{d-1}\rho(r,d)\right)dt^2 - d\vec{x}^2 - \frac{4\frac{d-2}{d-1}\rho(r,d)}{1 - 4\frac{d-2}{d-1}\rho(r,d)} \frac{(\vec{x} \cdot d\vec{x})^2}{r^2}.$$
 (5.2)

$$ds^{2} = \left(1 - \frac{\mu}{r^{d-2}}\right)dt^{2} - \frac{dr^{2}}{1 - \frac{\mu}{r^{d-1}}} - r^{2}d\Omega_{d-1}$$
(5.1)

with $\mu = \frac{16\pi G_N m}{(d-1)\Omega_{d-1}}$ and $\Omega_{d-1} = \frac{2\pi^{\frac{d}{2}}}{\Gamma(\frac{d}{2})}$ is the area of the unit (d-1)-sphere.

⁷In spherical coordinate the metric reads

As explained in section 2 the amplitude computation selects the de Donder gauge in (2.4). We make the coordinate transformation $(t, \vec{x}) \to (t, f(r)\vec{x})$ so that the Schwarzschild metric reads

$$ds^{2} = h_{0}(r)dt^{2} - h_{1}(r)d\vec{x}^{2} - h_{2}(r)\frac{(\vec{x} \cdot d\vec{x})^{2}}{r^{2}},$$
(5.3)

with $r = |\vec{x}|$ and

$$h_0(r) := 1 - 4\frac{d-2}{d-1} \frac{\rho(r,d)}{f(r)^{d-2}},$$

$$h_1(r) := f(r)^2,$$

$$h_2(r) := -f(r)^2 - f(r)^{d-2} \frac{(f(r) + r\frac{df(r)}{dr})^2}{f(r)^{d-2} - 4\frac{d-2}{r}\rho(r,d)}.$$
(5.4)

The de Donder gauge condition (2.4) then reads

$$2(d-1)h_2(r) = r\frac{d}{dr}\left(h_0(r) + (d-2)h_1(r) - h_2(r)\right).$$
(5.5)

We will be solving the de Donder gauge condition (2.4) in four dimensions (d = 3), five dimensions (d = 4) and six dimensions (d = 5), using the post-Minkowskian expansion

$$f(r) = 1 + \sum_{n>1} f_n(r)\rho(r,d)^n$$
(5.6)

with the condition at each order that

$$\lim_{r \to +\infty} f_n(r)/r^n = 0. \tag{5.7}$$

5.1 The metric in the de Donder gauge in four dimensions

The de Donder gauge condition (2.4) in d=3 reads

$$4h_2(r) = r\frac{d}{dr} \left(h_0(r) + h_1(r) - h_2(r) \right) , \qquad (5.8)$$

supplemented with the asymptotic boundary condition

$$\lim_{r \to \infty} f(r) = 1. \tag{5.9}$$

This differential equation implies either that f(r) = C/r, which does not satisfy the boundary condition (5.9), or f(r) satisfies the differential equation, with $x = G_N m/r$

$$xf(x)^{3}(2x - f(x))\frac{d^{2}f(x)}{dx^{2}} + (xf(x))^{2}\left(\frac{df(x)}{dx}\right)^{2} + 2f(x)^{3}(f(x) - 3x)\frac{df(x)}{dx} - 3(f(x))^{4} + 8(f(x))^{3}x + (f(x))^{2} - 4f(x)x + 4x^{2} = 0.$$
(5.10)

We solve the equation (5.10) using a series expansion in $G_N m$ using (5.6) and the boundary condition (5.7). The result to the order $(G_N m)^7$ is given by

$$f(r) = 1 + \frac{G_N m}{r} + 2\left(\frac{G_N m}{r}\right)^2 + \frac{2}{3}\log\left(\frac{rC_3}{G_N m}\right)\left(\frac{G_N m}{r}\right)^3 + \left(\frac{2}{3} - \frac{4}{3}\log\left(\frac{rC_3}{G_N m}\right)\right)\left(\frac{G_N m}{r}\right)^4 + \left(-\frac{21}{25} + \frac{32}{15}\log\left(\frac{rC_3}{G_N m}\right)\right)\left(\frac{G_N m}{r}\right)^5 + \left(\frac{112}{75} - \frac{28}{15}\log\left(\frac{rC_3}{G_N m}\right)\right)\left(\frac{G_N m}{r}\right)^6 + \left(\frac{50023}{34300} + \frac{1139}{2205}\log\left(\frac{rC_3}{G_N m}\right) + \frac{2}{7}\log\left(\frac{rC_3}{G_N m}\right)^2\right)\left(\frac{G_N m}{r}\right)^7 + O(G_N^8). \quad (5.11)$$

This solution is finite and has $\log(r)$ terms from the order G_N^3 . The solution has a single constant of integration C_3 associated with the scale of the logarithm.

5.1.1 The metric perturbation

In d=3 we derive components of the metric in perturbation by plugging the expression for f(r) in (5.11) in (5.4).

We obtain for the time component

$$h_0^{\text{dD}}(r) = 1 - 2\frac{G_N m}{r} + 2\left(\frac{G_N m}{r}\right)^2 + 2\left(\frac{G_N m}{r}\right)^3 + \left(\frac{4}{3}\log\left(\frac{rC_3}{G_N m}\right) - 6\right)\left(\frac{G_N m}{r}\right)^4 + \left(-\frac{16}{3}\log\left(\frac{rC_3}{G_N m}\right) + \frac{10}{3}\right)\left(\frac{G_N m}{r}\right)^5 + \left(\frac{124}{15}\log\left(\frac{rC_3}{G_N m}\right) + \frac{424}{75}\right)\left(\frac{G_N m}{r}\right)^6 + \left(-\frac{8}{9}\log\left(\frac{rC_3}{G_N m}\right)^2 + \frac{16}{15}\log\left(\frac{rC_3}{G_N m}\right) - \frac{674}{75}\right)\left(\frac{G_N m}{r}\right)^7 + O(G_N^8), \quad (5.12)$$

and for the spatial components

$$h_1^{\text{dD}}(r) = 1 + 2\frac{G_N m}{r} + 5\left(\frac{G_N m}{r}\right)^2 + \left(\frac{4}{3}\log\left(\frac{rC_3}{G_N m}\right) + 4\right)\left(\frac{G_N m}{r}\right)^3 + \left(-\frac{4}{3}\log\left(\frac{rC_3}{G_N m}\right) + \frac{16}{3}\right)\left(\frac{G_N m}{r}\right)^4 + \left(\frac{64}{15}\log\left(\frac{rC_3}{G_N m}\right) - \frac{26}{75}\right)\left(\frac{G_N m}{r}\right)^5 + \left(\frac{4}{9}\log\left(\frac{rC_3}{G_N m}\right)^2 - \frac{24}{5}\log\left(\frac{rC_3}{G_N m}\right) + \frac{298}{75}\right)\left(\frac{G_N m}{r}\right)^6 + O(G_N^7), \quad (5.13)$$

and

$$h_2^{\text{dD}}(r) = -7\left(\frac{G_N m}{r}\right)^2 - \left(4\log\left(\frac{rC_3}{G_N m}\right) + \frac{38}{3}\right)\left(\frac{G_N m}{r}\right)^3 + \left(\frac{8}{3}\log\left(\frac{rC_3}{G_N m}\right) - \frac{58}{3}\right)\left(\frac{G_N m}{r}\right)^4 - \left(\frac{16}{3}\log\left(\frac{rC_3}{G_N m}\right) - \frac{32}{3}\right)\left(\frac{G_N m}{r}\right)^5 + \left(\frac{4}{3}\log\left(\frac{rC_3}{G_N m}\right)^2 + \frac{508}{45}\log\left(\frac{rC_3}{G_N m}\right) + \frac{7378}{225}\right)\left(\frac{G_N m}{r}\right)^6 + O(G_N^7). \quad (5.14)$$

Notice the appearance of the $\log(r)^2$ at the sixth post-Minkowskian order, G_N^6 , in the spatial components of the metric. This is one order less than the appearance in the time

component. The same phenomenon happens for the $\log(r)$ contribution which appears one order earlier in the spatial component than in the time component.

5.2 The metric in the de Donder gauge in five dimensions

The de Donder gauge condition (2.4) in d=4 reads

$$6h_2(r) = r\frac{d}{dr} \left(h_0(r) + 2h_1(r) - h_2(r) \right) , \qquad (5.15)$$

supplemented with the asymptotic boundary condition

$$\lim_{r \to \infty} f(r) = 1. \tag{5.16}$$

This differential equation implies either that f(r) = C/r, which does not satisfy the boundary condition (5.16), or f(r) satisfies the differential equation, setting $x = G_N m/(\pi r^2)$

$$xf(x)^{5} \left(8x - 3f(x)^{2}\right) \frac{d^{2}f(x)}{dx^{2}} + 8f(x)^{4}x^{2} \left(\frac{df(x)}{dx}\right)^{2} + f(x)^{5} \left(3f(x)^{2} - 16x\right) \frac{df(x)}{dx}$$
$$-4f(x)^{6} + (16x + 2)f(x)^{4} - \frac{32}{3}xf(x)^{2} + \frac{128x^{2}}{9} = 0. \quad (5.17)$$

We solve the equation (5.17) using a series expansion in $G_N m$ using (5.6) and the boundary condition (5.7). The result to the order $(G_N m)^7$ is given by

$$f(r) = 1 + \frac{2}{3} \frac{G_N m}{\pi r^2} + \frac{10}{9} \log \left(\frac{r^2 C_2}{G_N m}\right) \left(\frac{G_N m}{\pi r^2}\right)^2 - \frac{4}{81} \left(-8 + 45 \log \left(\frac{r^2 C_2}{G_N m}\right)\right) \left(\frac{G_N m}{\pi r^2}\right)^3 + \frac{67 + 3780 \log \left(\frac{r^2 C_2}{G_N m}\right)}{972} \left(\frac{G_N m}{\pi r^2}\right)^4 - \frac{32963 + 156420 \log \left(\frac{r^2 C_2}{G_N m}\right) - 43200 \log \left(\frac{r^2 C_2}{G_N m}\right)^2}{21870} \left(\frac{G_N m}{\pi r^2}\right)^5 + \frac{409303 + 1620270 \log \left(\frac{r^2 C_2}{G_N m}\right) - 1087200 \log \left(\frac{r^2 C_2}{G_N m}\right)^2}{131220} \left(\frac{G_N m}{\pi r^2}\right)^6 - \frac{11148022313 + 37508666370 \log \left(\frac{r^2 C_2}{G_N m}\right) - 64367301600 \log \left(\frac{r^2 C_2}{G_N m}\right)^2}{2362944150} \left(\frac{G_N m}{\pi r^2}\right)^7 - \frac{4939200000}{2362944150} \log \left(\frac{r^2 C_2}{G_N m}\right)^3 \left(\frac{G_N m}{\pi r^2}\right)^7 + O(G_N^8). \quad (5.18)$$

Again there is a single constant of integration C_2 arising as the scale of the $\log(r)$ arising from the G_N^2 order.

5.2.1 The metric perturbation

In d=4 we derive components of the metric in perturbation by plugging the expression for f(r) in (5.18) in (5.4).

We obtain for the time component

$$h_0^{\text{dD}}(r) = 1 - \frac{8}{3} \frac{G_N m}{\pi r^2} + \frac{32}{9} \left(\frac{G_N m}{\pi r^2}\right)^2 + \frac{32\left(-3 + 5\log\left(\frac{r^2 C_2}{G_N m}\right)\right)}{27} \left(\frac{G_N m}{\pi r^2}\right)^3 - \frac{640\left(-2 + 9\log\left(\frac{r^2 C_2}{G_N m}\right)\right)}{243} \left(\frac{G_N m}{\pi r^2}\right)^4 + O(G_N^5), \quad (5.19)$$

and for the spatial components

$$h_1^{\text{dD}}(r) = 1 + \frac{4}{3} \frac{G_N m}{\pi r^2} + \frac{4\left(1 + 5\log\left(\frac{r^2 C_2}{G_N m}\right)\right)}{9} \left(\frac{G_N m}{\pi r^2}\right)^2 + \frac{\left(64 - 240\log\left(\frac{r^2 C_2}{G_N m}\right)\right)}{81} \left(\frac{G_N m}{\pi r^2}\right)^3 + \frac{\left(323 + 2340\log\left(\frac{r^2 C_2}{G_N m}\right) + 600\log^2\left(\frac{r^2 C_2}{G_N m}\right)\right)}{486} \left(\frac{G_N m}{\pi r^2}\right)^4 + O(G_N^5), \quad (5.20)$$

and

$$h_2^{\text{dD}}(r) = \frac{40\left(1 - 2\log\left(\frac{r^2C_2}{G_Nm}\right)\right)}{9} \left(\frac{G_Nm}{\pi r^2}\right)^2 + \frac{32\left(-4 + 5\log\left(\frac{r^2C_2}{G_Nm}\right)\right)}{27} \left(\frac{G_Nm}{\pi r^2}\right)^3 + \frac{8\left(-31 - 1260\log\left(\frac{r^2C_2}{G_Nm}\right) + 300\log\left(\frac{r^2C_2}{G_Nm}\right)^2\right)}{243} \left(\frac{G_Nm}{\pi r^2}\right)^4 + O(G_N^5). \quad (5.21)$$

5.3 The metric in the de Donder gauge in six dimensions

The de Donder gauge condition (2.4) in d = 5 reads

$$8h_2(r) = r\frac{d}{dr}\left(h_0(r) + 3h_1(r) - h_2(r)\right), \qquad (5.22)$$

supplemented with the asymptotic boundary condition

$$\lim_{r \to \infty} f(r) = 1. (5.23)$$

This differential equation implies either that f(r) = C/r, which does not satisfy the boundary condition (5.23), or f(r) satisfies the differential equation with $x = G_N m/(\pi r^3)$

$$xf(x)^{7} \left(6x - 4f(x)^{3}\right) \frac{d^{2} f(x)}{dx^{2}} + 9f(x)^{6} x^{2} \left(\frac{df(x)}{dx}\right)^{2} + f(x)^{7} \left(\frac{8}{3} f(x)^{3} - 10x\right) \frac{df(x)}{dx} - \frac{5}{3} f(x)^{8} + f(x)^{6} + 4xf(x)^{5} - 3xf(x)^{3} + \frac{9x^{2}}{4} = 0.$$
 (5.24)

We solve the equation (5.24) using a series expansion in G_N using (5.6) and the boundary condition (5.7). Asking for an expression with only integer powers of G_N , the result to the order G_N^7 is given by

$$f(r) = 1 + \frac{G_N m}{4\pi r^3} - \frac{5}{8} \left(\frac{G_N m}{\pi r^3}\right)^2 + \frac{2}{3} \left(\frac{G_N m}{\pi r^3}\right)^3 - \frac{775}{1344} \left(\frac{G_N m}{\pi r^3}\right)^4 + \frac{545977}{537600} \left(\frac{G_N m}{\pi r^3}\right)^5 - \frac{15194099}{10483200} \left(\frac{G_N m}{\pi r^3}\right)^6 + \frac{4421000509}{1878589440} \left(\frac{G_N m}{\pi r^3}\right)^7 + O(G_N^8) \,. \quad (5.25)$$

The expression is uniquely determined and finite.

5.3.1 The metric perturbation

In d = 5 we derive components of the metric in perturbation by plugging the expression for f(r) in (5.25) in (5.4).

We obtain for the metric components

$$h_0^{\text{dD}}(r) = 1 - \frac{3G_N m}{2\pi r^3} + \frac{9}{8} \left(\frac{G_N m}{\pi r^3}\right)^2 - \frac{27}{8} \left(\frac{G_N m}{\pi r^3}\right)^3 + \frac{387}{64} \left(\frac{G_N m}{\pi r^3}\right)^4 + O(G_N^5),$$

$$h_1^{\text{dD}}(r) = 1 + \frac{G_N m}{2\pi r^3} - \frac{19}{16} \left(\frac{G_N m}{\pi r^3}\right)^2 + \frac{49}{48} \left(\frac{G_N m}{\pi r^3}\right)^3 - \frac{577}{1344} \left(\frac{G_N m}{\pi r^3}\right)^4 + O(G_N^5),$$

$$h_2^{\text{dD}}(r) = \frac{117}{16} \left(\frac{G_N m}{\pi r^3}\right)^2 - \frac{45}{16} \left(\frac{G_N m}{\pi r^3}\right)^3 + \frac{1599}{112} \left(\frac{G_N m}{\pi r^3}\right)^4 + O(G_N^5).$$
(5.26)

6 Recovering the Schwarzschild-Tangherlini metric from the amplitude computations

In this section we show how the amplitude computations match the Schwarzschild-Tangherlini metric in four, five and six dimensions in the de Donder gauge of the previous section.

6.1 The Schwarzschild metric in four dimensions

6.1.1 The first post-Minkowskian contribution $O(G_N)$

Setting d = 3 in the expressions for the metric perturbation from the tree-level amplitude in (3.14) matches the de Donder gauge first post-Minkowskian order in four dimension (d = 3) in (5.12)–(5.14).

6.1.2 The second post-Minkowskian contribution $O(G_N^2)$

At the order G_N^2 , setting d=3 in the metric perturbation from the one-loop amplitude in (3.26) matches the metric in the de Donder gauge in four dimensions (d=3) in (5.12)–(5.14).

6.1.3 The third post-Minkowskian contributions $O(G_N^3)$

At this order the components of the metric in the de Donder gauge in four dimensions (d=3) from (5.12)—(5.14) match the metric components from the renormalised two-loop amplitude computation in (4.34) for the value of the constant of integration

$$\log C_3 = \log C_E - \frac{7}{2} + 6a^{(1)}(3), \qquad (6.1)$$

where C_E is given in (4.35).

With this identification we recover the results of [27] for the renormalisation of the metric divergences and the coordinate change from the de Donder gauge to the harmonic gauge from the world-line approach.

Substituting this value of C_3 in the solution (5.11) completely determines the solution to the de Donder gauge in four dimensions and the coordinate change in (5.11) to the Schwarzschild metric in (5.3) in four dimensions. The parameter $a^{(1)}(3)$ is a free parameter, which corresponds to the running coupling in [27].

6.1.4 The fourth post-Minkowskian contribution $O(G_N^4)$

At the fourth post-Minkowskian order, we get again a diverging metric from the amplitude computation. This finite component metric in the de Donder gauge in four dimensions (d=3) in (5.12)—(5.14) using the value of the constant of integration C_3 determined in (6.1) give

$$h_0^{\text{dD}(4)} = \left(-\frac{32}{3} + 8a^{(1)}(3) + \frac{4}{3}\log\left(\frac{rC_E}{G_N m}\right)\right) \left(\frac{G_N m}{r}\right)^4,$$

$$h_1^{\text{dD}(4)} = \left(10 - 8a^{(1)}(3) - \frac{4}{3}\log\left(\frac{rC_E}{G_N m}\right)\right) \left(\frac{G_N m}{r}\right)^4,$$

$$h_2^{\text{dD}(4)} = \left(-\frac{86}{3} + 16a^{(1)}(3) + \frac{8}{3}\log\left(\frac{rC_E}{G_N m}\right)\right) \left(\frac{G_N m}{r}\right)^4.$$
(6.2)

This matches exactly the renormalised metric components from the three-loop amplitude computation obtained in (4.39) with d=3.

6.2 The Schwarzschild-Tangherlini metric in five dimensions

6.2.1 The first post-Minkowskian contribution $O(G_N)$

Setting d = 4 in the expressions for the metric perturbation from the tree-level amplitude in (3.14) matches the de Donder gauge first post-Minkowskian order in five dimensions (d = 4) in (5.19)–(5.21).

6.2.2 The second post-Minkowskian contribution $O(G_N^2)$

The renormalised one-loop computation in (4.44) matches the expression at order $O(G_N^2)$ from the de Donder gauge in (5.19)—(5.21) for the choice of the constant of integration

$$\log C_2 = \frac{11}{15} + 2\log C_E + \frac{36\pi}{5}a^{(1)}(5). \tag{6.3}$$

Again there is a free parameter $a^{(1)}(5)$ which can be associated with a running coupling constant.

6.2.3 The third post-Minkowskian contributions $O(G_N^3)$

At this order in perturbation, the two-loop amplitude computation had divergences that had to be renormalized to give (4.49). This matches exactly the finite component metric in the de Donder gauge in five dimensions (d = 4) in (5.19)—(5.21), using the value of the constant of integration C_2 determined in (6.3), given by

$$h_0^{\text{dD(3)}} = \frac{160}{27} \left(\frac{2}{15} + \frac{36a^{(1)}(5)\pi}{5} + \log\left(\frac{r^2 C_E^2}{G_N m}\right) \right) \left(\frac{G_N m}{\pi r^2}\right)^3 + O(d - 4), \tag{6.4}$$

$$h_1^{\text{dD(3)}} = -\frac{80}{27} \left(\frac{7}{15} + \frac{36a^{(1)}(5)\pi}{5} + \log\left(\frac{r^2 C_E^2}{G_N m}\right) \right) \left(\frac{G_N m}{\pi r^2}\right)^3 + O(d - 4),$$

$$h_2^{\text{dD(3)}} = \frac{160}{27} \left(-\frac{1}{15} + \frac{36a^{(1)}(5)\pi}{5} + \log\left(\frac{r^2 C_E^2}{G_N m}\right) \right) \left(\frac{G_N m}{\pi r^2}\right)^3 + O(d - 4).$$

6.2.4 The fourth post-Minkowskian contribution $O(G_N^4)$

The three-loop amplitude computation diverges and the finite metric component at the fourth post-Minkowskian order was obtained after normalisation in (4.56). This matches exactly, the finite component metric in the de Donder gauge in five dimensions (d = 4) in (5.19)—(5.21), using the value of the constant of integration C_2 determined in (6.3), given by

$$h_0^{\text{dD}(4)} = -\frac{128}{243} \left(23 + 324a^{(1)}(5)\pi + 45\log\left(\frac{r^2C_E^2}{G_Nm}\right) \right) \left(\frac{G_Nm}{\pi r^2}\right)^4 + O(d-4),$$

$$h_1^{\text{dD}(4)} = \left(\frac{7085 + 69552\pi a^{(1)}(5) + 93312(\pi a^{(1)}(5))^2}{1458} + \frac{10}{243}(161 + 432\pi a^{(1)}(5))\log\left(\frac{r^2C_E^2}{G_Nm}\right) + \frac{100}{81}\log\left(\frac{r^2C_E^2}{G_Nm}\right)^2\right) \left(\frac{G_Nm}{\pi r^2}\right)^4 + O(d-4),$$

$$h_2^{\text{dD}(4)} = \left(\frac{-19048 - 141696\pi a^{(1)}(5)373248(\pi a^{(1)}(5))^2}{729} + \frac{160}{243}(-41 + 216\pi a^{(1)}(5))\log\left(\frac{r^2C_E^2}{G_Nm}\right) + \frac{800}{81}\log\left(\frac{r^2C_E^2}{G_Nm}\right)^2\right) \left(\frac{G_Nm}{\pi r^2}\right)^4 + O(d-4).$$

$$(6.5)$$

6.3 The Schwarzschild-Tangherlini metric in six dimensions

The metric components in six dimensions (d = 5) are finite. They are given up to the order $O(G_N^4)$ in (5.26) and are reproduced by the sum of the contributions of the tree-level amplitude in (3.14), one-loop amplitude in (3.26), two-loop amplitude in (3.44) and three-loop amplitude in (3.54) and setting d = 5 in these expressions.

7 Discussion

General relativity can be considered in space-times of various dimensions. It is therefore important to validate our current understanding of the connection between scattering amplitudes and classical general relativity in general dimensions [22, 23]

We have shown how to reconstruct the classical Schwarzschild-Tangherlini metric from scattering amplitudes in four, five and six dimensions. We have extracted the classical contribution as defined in [12] from the vertex function for the emission of a graviton from a massive scalar field. For such a static metric, the classical contribution is obtained by taking appropriate residues on the time components of the loop momenta. These residues project the quantum scattering amplitude on contribution similar to the quantum tree graphs considered in [7], by cutting the massive propagators.

The amplitudes develop ultraviolet divergences which are renormalised by introducing higher-derivative non-minimal couplings in (4.2). The non-minimal coupling removes the ultraviolet divergences in the stress-tensor and the metric components. For the static solution the higher $n \geq 2$ non-minimal coupling only contribute from insertions in tree-level graphs. Interestingly, in six dimensions the metric components are finite but the stress-tensor has ultraviolet divergences. These divergences are removed by adding counter-terms

from non-minimal couplings. These counter-terms do not induce any contribution to the metric components. From the presence of ultraviolet poles in the master integrals $J_{(l)}(\vec{q}^2)$ in (2.31), we conclude that in all dimensions one needs to introduce an infinite set of higher-derivative non-minimal operators for removing the ultraviolet divergences from the scattering amplitude. These counter-terms do not affect the space-time geometry because their effect is reabsorbed by the change of coordinate from the de Donder coordinate system to the Schwarzschild-Tangherlini coordinate system.

The scattering amplitude approach presented in this work can be applied to any effective field theory of gravity coupled to matter fields. The amplitudes computations, being performed in general dimensions, lead to results that have an analytic dependence on the space-time dimensions. As black-hole solutions develop non trivial properties in general dimensions [25, 26], it is interesting to apply the method of this paper to other black-hole metrics. The Kerr-Newman and Reissner-Nordström metric in four dimensions have been obtained in [9, 41–47] by considering tree-level and one-loop vertex function of the emission of the graviton from a massive particle of spin s. The higher order post-Minkowskian contributions should be obtained from higher-loop amplitudes in a direct application of the methods used in this work.

Acknowledgments

We would like to thank Emil Bjerrum-Bohr, Poul Damgaard, Paolo di Vecchia, Ludovic Planté for discussions and comments. The research of P. Vanhove has received funding from the ANR grant "Amplitudes" ANR-17- CE31-0001-01, and is partially supported by Laboratory of Mirror Symmetry NRU HSE, RF Government grant, ag. N° 14.641.31.0001. P.V. is grateful to the I.H.E.S. for allowing to use their computer resources.

A Fourier transforms

Here we collect the Fourier integrals used to calculate the long range corrections to the energy momentum tensor and the metric.

The Fourier transform form momentum space to direct space

$$\mathcal{F}(\alpha, d) = \int_{\mathbb{R}^d} \frac{1}{|q|^{\alpha}} e^{i\vec{q}\cdot\vec{x}} \frac{d^d\vec{q}}{(2\pi)^d} = \frac{1}{(4\pi)^{\frac{d}{2}}} \frac{\Gamma\left(\frac{d-\alpha}{2}\right)}{\Gamma\left(\frac{\alpha}{2}\right)} \left(\frac{2}{|\vec{x}|}\right)^{d-\alpha}. \tag{A.1}$$

Using that

$$\partial_{x^i}\partial_{x^j}(\vec{x}^2)^\alpha = 2\alpha(\vec{x}^2)^{\alpha-1}\left(\delta_{ij} + 2(\alpha - 1)\frac{x_i x_j}{\vec{x}^2}\right),\tag{A.2}$$

we have that

$$\mathcal{F}_{ij}(\alpha, d) := \int_{\mathbb{R}^d} \frac{q_i q_j}{|\vec{q}|^{\alpha+2}} e^{i\vec{q}\cdot\vec{x}} \frac{d^d \vec{q}}{(2\pi)^d} = \mathcal{F}(\alpha, d) \left(\frac{1}{\alpha} \delta_{ij} + \frac{\alpha - d}{\alpha} \frac{x_i x_j}{\vec{x}^2} \right) . \tag{A.3}$$

We have in particular that

$$\mathcal{F}(0,d) = 0, \qquad \mathcal{F}_{ij}(0,d) = \frac{\Gamma\left(\frac{d}{2}\right)}{2\pi^{\frac{d}{2}}|\vec{x}|^d} \left(\delta_{ij} - d\frac{x_i x_j}{\vec{x}^2}\right). \tag{A.4}$$

B Vertices and Propagators

We will here list the Feynman rules which are employed in our calculation. For the derivation of these forms, see [3–5, 9, 48–50]. Our convention differs from these work by having all incoming momenta. We have stripped off factors of $i\sqrt{8\pi G_N}$ from the vertices and made them explicit in the amplitudes.

- The massive scalar propagator is $\frac{i}{q^2 m^2 + i\varepsilon}$.
- The graviton propagator in de Donder gauge can be written in the form $\frac{i\mathcal{P}^{\alpha\beta,\gamma\delta}}{q^2+i\varepsilon}$ where $\mathcal{P}^{\alpha\beta,\gamma\delta}$ is defined by

$$\mathcal{P}^{\mu\nu,\rho\sigma} = \frac{1}{2} \left(\eta^{\mu\rho} \eta^{\nu\sigma} + \eta^{\mu\sigma} \eta^{\nu\rho} - \frac{2}{D-2} \eta^{\mu\nu} \eta^{\rho\sigma} \right)$$
 (B.1)

• The 2-scalar-1-graviton vertex $\tau_1^{\mu\nu}(p_1, p_2)$ is

$$\tau^{\mu\nu}(p_1, p_2) = p_1^{\mu} p_2^{\nu} + p_1^{\nu} p_2^{\mu} + \frac{1}{2} \eta^{\mu\nu} (p_1 - p_2)^2.$$
 (B.2)

• The three-graviton vertex has been derived in [49], where $k + q + \pi = 0$,

$$\begin{split} \tau_{(3)}{}^{\mu\nu}_{\alpha\beta,\gamma\delta}(k,q) &= -\left(\mathcal{P}_{\alpha\beta\gamma\delta}\bigg[k^{\mu}k^{\nu} + \pi^{\mu}\pi^{\nu} + q^{\mu}q^{\nu} - \frac{3}{2}\eta^{\mu\nu}q^{2}\bigg] \\ &+ 2q_{\lambda}q_{\sigma}\bigg[I_{\alpha\beta}{}^{\sigma\lambda}I_{\gamma\delta}{}^{\mu\nu} + I_{\gamma\delta}{}^{\sigma\lambda}I_{\alpha\beta}{}^{\mu\nu} - I_{\alpha\beta}{}^{\mu\sigma}I_{\gamma\delta}{}^{\nu\lambda} - I_{\gamma\delta}{}^{\mu\sigma}I_{\alpha\beta}{}^{\nu\lambda}\bigg] \\ &+ \bigg[q_{\lambda}q^{\mu}\bigg(\eta_{\alpha\beta}I_{\gamma\delta}{}^{\nu\lambda} + \eta_{\gamma\delta}I_{\alpha\beta}{}^{\nu\lambda}\bigg) + q_{\lambda}q^{\nu}\bigg(\eta_{\alpha\beta}I_{\gamma\delta}{}^{\mu\lambda} + \eta_{\gamma\delta}I_{\alpha\beta}{}^{\mu\lambda}\bigg) \\ &- q^{2}\bigg(\eta_{\alpha\beta}I_{\gamma\delta}{}^{\mu\nu} + \eta_{\gamma\delta}I_{\alpha\beta}{}^{\mu\nu}\bigg) - \eta^{\mu\nu}q_{\sigma}q_{\lambda}\bigg(\eta_{\alpha\beta}I_{\gamma\delta}{}^{\sigma\lambda} + \eta_{\gamma\delta}I_{\alpha\beta}{}^{\sigma\lambda}\bigg)\bigg] \\ &+ \bigg[2q_{\lambda}\bigg(I_{\alpha\beta}{}^{\lambda\sigma}I_{\gamma\delta\sigma}{}^{\nu}\pi^{\mu} + I_{\alpha\beta}{}^{\lambda\sigma}I_{\gamma\delta\sigma}{}^{\mu}\pi^{\nu} + I_{\gamma\delta}{}^{\lambda\sigma}I_{\alpha\beta\sigma}{}^{\nu}k^{\mu} + I_{\gamma\delta}{}^{\lambda\sigma}I_{\alpha\beta\sigma}{}^{\mu}k^{\nu}\bigg) \\ &+ q^{2}\bigg(I_{\alpha\beta\sigma}{}^{\mu}I_{\gamma\delta}{}^{\nu\sigma} + I_{\alpha\beta}{}^{\nu\sigma}I_{\gamma\delta\sigma}{}^{\mu}\bigg) + \eta^{\mu\nu}q_{\sigma}q_{\lambda}\bigg(I_{\alpha\beta}{}^{\lambda\rho}I_{\gamma\delta\rho}{}^{\sigma} + I_{\gamma\delta}{}^{\lambda\rho}I_{\alpha\beta\rho}{}^{\sigma}\bigg)\bigg] \\ &+ \bigg\{(k^{2} + \pi^{2})\big[\mathcal{P}_{\alpha\beta}{}^{\mu\sigma}\mathcal{P}_{\gamma\delta,\sigma}{}^{\nu} + \mathcal{P}_{\gamma\delta}{}^{\mu\sigma}\mathcal{P}_{\alpha\beta,\sigma}{}^{\nu} - \frac{1}{2}\eta^{\mu\nu}(\mathcal{P}_{\alpha\beta,\gamma\delta} - \eta_{\alpha\beta}\eta_{\gamma\delta})\bigg] \\ &+ \bigg(\mathcal{P}_{\gamma\delta}{}^{\mu\nu}\eta_{\alpha\beta}\pi^{2} + \mathcal{P}_{\alpha\beta}{}^{\mu\nu}\eta_{\gamma\delta}k^{2}\bigg)\bigg\}\bigg), \end{split}$$

where $I_{\alpha\beta,\gamma\delta} := \mathcal{P}_{\alpha\beta,\gamma\delta} + \frac{1}{2} \eta_{\alpha\beta} \eta_{\gamma\delta}$. These vertices are equivalent to the ones computed with the vertices given by De Witt [3–5] and Sannan [50]. We remark that the expression for τ_3 is simpler than the three-graviton vertex in these references.

We notice that the three-graviton vertex satisfies the identity

$$\tau^{\mu\nu}_{(3) \pi\rho,\sigma\tau}(l,q) \mathcal{P}^{\pi\rho}_{\alpha\beta} \mathcal{P}^{\sigma\tau}_{\gamma\delta} = \tau^{\mu\nu}_{(3) \alpha\beta,\gamma\delta}(l,q) \tag{B.4}$$

that will be used to simplify the expression of the amplitude.

• The four-graviton vertex with $k_1 + k_2 + k_3 + k_4 = 0$ is given in [31, 50]

$$\begin{split} \tilde{\tau}_{(4)\,\mu\nu,\sigma\tau,\iota\kappa}^{\rho\lambda}(k_1,k_2,k_3,k_4) &= -\frac{1}{32}(k_1 \cdot k_2 \eta^{\mu\nu} \eta^{\sigma\tau} \eta^{\rho\lambda} \eta_{\iota\kappa}) - \frac{1}{16}(k_1^{\sigma} k_1^{\tau} \eta^{\mu\nu} \eta^{\rho\lambda} \eta_{\iota\kappa}) \\ &- \frac{1}{16}(k_1^{\sigma} k_2^{\mu} \eta^{\nu\tau} \eta^{\rho\lambda} \eta_{\iota\kappa}) + \frac{1}{32}(k_1 \cdot k_2 \eta^{\mu\sigma} \eta^{\nu\tau} \eta^{\rho\lambda} \eta_{\iota\kappa}) + \frac{1}{16}(k_1 \cdot k_2 \eta^{\mu\nu} \eta^{\sigma\tau} \eta_{\iota}^{\rho} \eta_{\kappa}^{\lambda}) \\ &+ \frac{1}{8}(k_1^{\sigma} k_1^{\tau} \eta^{\mu\nu} \eta_{\iota}^{\rho} \eta_{\kappa}^{\lambda}) + \frac{1}{8}(k_1^{\sigma} k_2^{\mu} \eta^{\nu\tau} \eta_{\iota}^{\rho} \eta_{\kappa}^{\lambda}) - \frac{1}{16}(k_1 \cdot k_2 \eta^{\mu\sigma} \eta^{\nu\tau} \eta_{\iota}^{\rho} \eta_{\kappa}^{\lambda}) \\ &+ \frac{1}{4}(k_1 \cdot k_2 \eta^{\mu\nu} \eta^{\sigma\rho} \eta_{\tau\lambda} \eta_{\iota\kappa}) + \frac{1}{4}(k_1^{\sigma} k_1^{\tau} \eta^{\mu\rho} \eta^{\nu\lambda} \eta_{\iota\kappa}) + \frac{1}{8}(k_1^{\rho} k_2^{\lambda} \eta^{\mu\sigma} \eta^{\nu\tau} \eta_{\iota\kappa}) \\ &+ \frac{1}{2}(k_1^{\sigma} k_2^{\rho} \eta^{\tau\mu} \eta^{\nu\lambda} \eta_{\iota\kappa}) - \frac{1}{4}(k_1 \cdot k_2 \eta^{\nu\sigma} \eta^{\tau\rho} \eta^{\lambda\mu} \eta_{\iota\kappa}) + \frac{1}{4}(k_1^{\sigma} k_2^{\mu} \eta^{\tau\rho} \eta^{\lambda\nu} \eta_{\iota\kappa}) \\ &+ \frac{1}{4}(k_1^{\sigma} k_1^{\rho} \eta^{\tau\lambda} \eta^{\mu\nu} \eta_{\iota\kappa}) - \frac{1}{2}(k_1 \cdot k_2 \eta^{\mu\nu} \eta^{\tau\rho} \eta_{\iota}^{\lambda} \eta_{\kappa}^{\sigma}) - \frac{1}{2}(k_1^{\sigma} k_1^{\tau} \eta^{\nu\rho} \eta_{\iota}^{\lambda} \eta_{\kappa}^{\mu}) \\ &- \frac{1}{2}(k_1^{\rho} k_2^{\lambda} \eta_{\iota}^{\nu} \eta_{\kappa}^{\sigma} \eta^{\tau\mu}) - (k_1^{\sigma} k_2^{\rho} \eta_{\iota}^{\tau} \eta_{\kappa}^{\mu} \eta^{\nu\lambda}) - \frac{1}{2}(k_1^{\rho} k_2 \iota_1 \eta^{\lambda\sigma} \eta^{\tau\mu} \eta_{\kappa}^{\nu}) \\ &+ \frac{1}{4}(k_1 \cdot k_2 \eta^{\nu\rho} \eta^{\lambda\sigma} \eta_{\iota}^{\tau} \eta_{\kappa}^{\mu}) - \frac{1}{2}(k_1^{\sigma} k_1^{\rho} \eta^{\mu\nu} \eta_{\iota}^{\tau} \eta_{\kappa}^{\lambda}) - \frac{1}{4}(k_1 \cdot k_2 \eta^{\mu\rho} \eta^{\nu\lambda} \eta_{\iota}^{\tau} \eta_{\kappa}^{\lambda}) \\ &- \frac{1}{2}(k_1^{\sigma} k_1^{\rho} \eta^{\tau\lambda} \eta_{\iota}^{\mu} \eta_{\kappa}^{\nu}) - \frac{1}{4}(k_1^{\rho} k_2 \iota_1 \eta_{\kappa}^{\lambda} \eta^{\mu\sigma} \eta^{\nu\tau}) - (k_1^{\sigma} k_2^{\rho} \eta^{\tau\mu} \eta_{\iota}^{\nu} \eta_{\kappa}^{\lambda}) \\ &- \frac{1}{2}(k_1^{\sigma} k_1^{\rho} \eta^{\tau\lambda} \eta_{\iota}^{\mu} \eta_{\kappa}^{\nu}) + \frac{1}{2}(k_1 \cdot k_2 \eta^{\nu\sigma} \eta^{\tau\rho} \eta_{\iota}^{\lambda} \eta_{\kappa}^{\mu}) + Sym(k_1, k_2, k_3, k_4) \\ &- \frac{1}{2}(k_1^{\sigma} k_2^{\mu} \eta^{\tau\rho} \eta_{\iota}^{\lambda} \eta_{\kappa}^{\nu}) + \frac{1}{2}(k_1 \cdot k_2 \eta^{\nu\sigma} \eta^{\tau\rho} \eta_{\iota}^{\lambda} \eta_{\kappa}^{\mu}) + Sym(k_1, k_2, k_3, k_4) \\ &- \frac{1}{2}(k_1^{\sigma} k_2^{\mu} \eta^{\tau\rho} \eta_{\iota}^{\lambda} \eta_{\kappa}^{\nu}) + \frac{1}{2}(k_1 \cdot k_2 \eta^{\nu\sigma} \eta^{\tau\rho} \eta_{\iota}^{\lambda} \eta_{\kappa}^{\mu}) + Sym(k_1, k_2, k_3, k_4) \\ &- \frac{1}{2}(k_1^{\sigma} k_2^{\mu} \eta^{\tau\rho} \eta_{\iota}^{\lambda} \eta_{\kappa}^{\nu}) + \frac{1}{2}(k_1 \cdot k_2 \eta^{\nu\sigma} \eta^{\tau\rho} \eta_{\iota}^{\lambda} \eta_{\kappa}^{\mu}) + Sym(k_1, k_2, k_3, k_4) \\ &- \frac{1}{2}(k_1^{\sigma} k_2^{\mu} \eta^{\tau\rho} \eta_{\iota}^{\lambda} \eta_{\kappa}^{\nu}) + \frac{1}{2}(k_1^{\sigma} k_2^{\mu} \eta^{\tau\rho}$$

we introduce the short hand notation

$$\tau^{\mu\nu}_{(4)\,\gamma\delta,\sigma\tau,\iota\theta}(k_1,k_2,k_3,k_4) := \tilde{\tau}^{\mu\nu}_{(4)\,\alpha\beta,\gamma\delta,\epsilon\eta}(k_1,k_2,k_3,k_4) \mathcal{P}^{\alpha\beta}_{\gamma\delta} \mathcal{P}^{\gamma\delta}_{\tau\tau} \mathcal{P}^{\epsilon\eta}_{\iota\theta} . \tag{B.6}$$

References

- [1] G. 't Hooft and M. J. G. Veltman, "One Loop Divergencies in the Theory of Gravitation," Ann. Inst. H. Poincare Phys. Theor. A **20** (1974), 69-94
- [2] M. J. G. Veltman, "Quantum Theory of Gravitation," Conf. Proc. C 7507281 (1975) 265.
- [3] B. S. DeWitt, "Quantum Theory of Gravity. 1. The Canonical Theory," Phys. Rev. 160 (1967), 1113-1148
- [4] B. S. DeWitt, "Quantum Theory of Gravity. 2. The Manifestly Covariant Theory," Phys. Rev. 162 (1967), 1195-1239
- [5] B. S. DeWitt, "Quantum Theory of Gravity. 3. Applications of the Covariant Theory," Phys. Rev. 162 (1967), 1239-1256
- [6] J. F. Donoghue, "General Relativity as an Effective Field Theory: the Leading Quantum Corrections," Phys. Rev. D **50** (1994), 3874-3888 [arXiv:gr-qc/9405057 [gr-qc]].
- [7] M. J. Duff, "Quantum Tree Graphs and the Schwarzschild Solution," Phys. Rev. D 7 (1973), 2317-2326
- [8] Y. Iwasaki, "Quantum Theory of Gravitation Vs. Classical Theory. Fourth-Order Potential," Prog. Theor. Phys. 46 (1971), 1587-1609
- [9] N. E. J. Bjerrum-Bohr, J. F. Donoghue and B. R. Holstein, "Quantum corrections to the Schwarzschild and Kerr metrics," Phys. Rev. D 68 (2003), 084005 [arXiv:hep-th/0211071 [hep-th]].

- [10] B. R. Holstein and J. F. Donoghue, "Classical Physics and Quantum Loops," Phys. Rev. Lett. 93 (2004) 201602 [hep-th/0405239].
- [11] J. F. Donoghue and T. Torma, "On the Power Counting of Loop Diagrams in General Relativity," Phys. Rev. D **54** (1996), 4963-4972 [arXiv:hep-th/9602121 [hep-th]].
- [12] N. E. J. Bjerrum-Bohr, P. H. Damgaard, G. Festuccia, L. Planté and P. Vanhove, "General Relativity from Scattering Amplitudes," Phys. Rev. Lett. 121 (2018) no.17, 171601 [arXiv:1806.04920 [hep-th]].
- [13] D. A. Kosower, B. Maybee and D. O'Connell, "Amplitudes, Observables, and Classical Scattering," JHEP **02** (2019), 137 [arXiv:1811.10950 [hep-th]].
- [14] C. Cheung, I. Z. Rothstein and M. P. Solon, "From Scattering Amplitudes to Classical Potentials in the Post-Minkowskian Expansion," Phys. Rev. Lett. 121 (2018) no.25, 251101 [arXiv:1808.02489 [hep-th]].
- [15] Z. Bern, C. Cheung, R. Roiban, C. H. Shen, M. P. Solon and M. Zeng, "Scattering Amplitudes and the Conservative Hamiltonian for Binary Systems at Third Post-Minkowskian Order," Phys. Rev. Lett. 122 (2019) no.20, 201603 [arXiv:1901.04424 [hep-th]].
- [16] Z. Bern, C. Cheung, R. Roiban, C. H. Shen, M. P. Solon and M. Zeng, "Black Hole Binary Dynamics from the Double Copy and Effective Theory," JHEP 10 (2019), 206 [arXiv:1908.01493 [hep-th]].
- [17] M. Z. Chung, Y. T. Huang and J. W. Kim, "Classical Potential for General Spinning Bodies," JHEP 09 (2020), 074 [arXiv:1908.08463 [hep-th]].
- [18] G. Kälin and R. A. Porto, "From Boundary Data to Bound States," JHEP 01 (2020), 072 [arXiv:1910.03008 [hep-th]].
- [19] C. Cheung and M. P. Solon, "Classical Gravitational Scattering at $\mathcal{O}(G^3)$ from Feynman Diagrams," JHEP **06** (2020), 144 [arXiv:2003.08351 [hep-th]].
- [20] G. Mogull, J. Plefka and J. Steinhoff, "Classical Black Hole Scattering from a Worldline Quantum Field Theory," [arXiv:2010.02865 [hep-th]].
- [21] A. K. Collado, P. Di Vecchia, R. Russo and S. Thomas, "The Subleading Eikonal in Supergravity Theories," JHEP 10 (2018), 038 [arXiv:1807.04588 [hep-th]].
- [22] A. Koemans Collado, P. Di Vecchia and R. Russo, "Revisiting the Second Post-Minkowskian Eikonal and the Dynamics of Binary Black Holes," Phys. Rev. D **100** (2019) no.6, 066028 [arXiv:1904.02667 [hep-th]].
- [23] A. Cristofoli, P. H. Damgaard, P. Di Vecchia and C. Heissenberg, "Second-Order Post-Minkowskian Scattering in Arbitrary Dimensions," JHEP **07** (2020), 122 [arXiv:2003.10274 [hep-th]].
- [24] G. U. Jakobsen, "Schwarzschild-Tangherlini Metric from Scattering Amplitudes," [arXiv:2006.01734 [hep-th]].
- [25] R. Emparan and H. S. Reall, "Black Holes in Higher Dimensions," Living Rev. Rel. 11 (2008), 6 [arXiv:0801.3471 [hep-th]].
- [26] R. Emparan, R. Suzuki and K. Tanabe, "The Large D Limit of General Relativity," JHEP **06** (2013), 009 [arXiv:1302.6382 [hep-th]].

- [27] W. D. Goldberger and I. Z. Rothstein, "An Effective Field Theory of Gravity for Extended Objects," Phys. Rev. D **73** (2006), 104029 [arXiv:hep-th/0409156 [hep-th]].
- [28] N. E. J. Bjerrum-Bohr, J. F. Donoghue and B. R. Holstein, "On the Parameterization Dependence of the Energy Momentum Tensor and the Metric," Phys. Rev. D 75 (2007), 108502 [arXiv:gr-qc/0610096 [gr-qc]].
- [29] P. Fromholz, E. Poisson and C. M. Will, "The Schwarzschild Metric: It's the Coordinates, Stupid!," Am. J. Phys. 82 (2014), 295 [arXiv:1308.0394 [gr-qc]].
- [30] Citation for the 2020 Nobel prize award to Roger Penrose https://www.nobelprize.org/prizes/physics/2020/press-release/
- [31] L. Planté, "Some aspects on effective field theory of gravity", Phd thesis, Université Paris-Saclay 2016
- [32] J. Galusha, "Investigations into the Classical Limit of Quantum Field Theory", Master thesis Niels Bohr International Academy 2018
- [33] M. Levy and J. Sucher, "Eikonal Approximation in Quantum Field Theory," Phys. Rev. 186 (1969), 1656-1670
- [34] R. N. Lee, "Presenting Litered: a Tool for the Loop Integrals Reduction," [arXiv:1212.2685 [hep-ph]].
- [35] R. N. Lee, "Litered 1.4: a Powerful Tool for Reduction of Multiloop Integrals," J. Phys. Conf. Ser. **523** (2014), 012059 [arXiv:1310.1145 [hep-ph]].
- [36] P. Vanhove, "The Physics and the Mixed Hodge Structure of Feynman Integrals," Proc. Symp. Pure Math. 88 (2014), 161-194 [arXiv:1401.6438 [hep-th]].
- [37] A. Cristofoli, N. E. J. Bjerrum-Bohr, P. H. Damgaard and P. Vanhove, "Post-Minkowskian Hamiltonians in General Relativity," Phys. Rev. D 100 (2019) no.8, 084040 [arXiv:1906.01579 [hep-th]].
- [38] J.C. Lagarias. "Euler's constant: Euler's work and modern developments." Bull. Amer. Math. Soc. (N.S.) **50** (2013), no. 4, 527–628.
- [39] D. Prinz, "Gravity-Matter Feynman Rules for Any Valence," [arXiv:2004.09543 [hep-th]].
- [40] F. R. Tangherlini, "Schwarzschild Field in N Dimensions and the Dimensionality of Space Problem," Nuovo Cim. 27 (1963), 636-651
- [41] J. F. Donoghue, B. R. Holstein, B. Garbrecht and T. Konstandin, "Quantum Corrections to the Reissner-Nordstrom and Kerr-Newman Metrics," Phys. Lett. B **529** (2002), 132-142 [erratum: Phys. Lett. B **612** (2005), 311-312] [arXiv:hep-th/0112237 [hep-th]].
- [42] A. Guevara, A. Ochirov and J. Vines, "Scattering of Spinning Black Holes from Exponentiated Soft Factors," JHEP **09** (2019), 056 [arXiv:1812.06895 [hep-th]].
- [43] M. Z. Chung, Y. T. Huang, J. W. Kim and S. Lee, "The Simplest Massive S-Matrix: from Minimal Coupling to Black Holes," JHEP 04 (2019), 156 [arXiv:1812.08752 [hep-th]].
- [44] N. Moynihan, "Kerr-Newman from Minimal Coupling," JHEP 01 (2020), 014 [arXiv:1909.05217 [hep-th]].
- [45] M. Z. Chung, Y. T. Huang and J. W. Kim, "Kerr-Newman Stress-Tensor from Minimal Coupling to All Orders in Spin," [arXiv:1911.12775 [hep-th]].

- [46] A. Guevara, A. Ochirov and J. Vines, "Black-Hole Scattering with General Spin Directions from Minimal-Coupling Amplitudes," Phys. Rev. D 100 (2019) no.10, 104024 [arXiv:1906.10071 [hep-th]].
- [47] A. Cristofoli, "Gravitational Shock Waves and Scattering Amplitudes," [arXiv:2006.08283 [hep-th]].
- [48] N. E. J. Bjerrum-Bohr, J. F. Donoghue and P. Vanhove, "On-Shell Techniques and Universal Results in Quantum Gravity," JHEP **02** (2014), 111 [arXiv:1309.0804 [hep-th]].
- [49] J. F. Donoghue, "Introduction to the Effective Field Theory Description of Gravity," in Advanced School on Effective Theories Almunecar, Spain, June 25-July 1, 1995, 1995. [arXiv:gr-qc/9512024 [gr-qc]].
- [50] S. Sannan, "Gravity as the Limit of the Type II Superstring Theory," Phys. Rev. D 34 (1986), 1749