

## STATISTICS WORKSHEET-5

**Q1 to Q10 are MCQs with only one correct answer. Choose the correct option.**

1. Using a goodness of fit, we can assess whether a set of obtained frequencies differ from a set of frequencies.

- a) Mean
- b) Actual
- c) Predicted
- d) Expected

D

2. Chisquare is used to analyse

- a) Score
- b) Rank
- c) Frequencies
- d) All of these

C

3. What is the mean of a Chi Square distribution with 6 degrees of freedom?

- a) 4
- b) 12
- c) 6
- d) 8

C

4. Which of these distributions is used for a goodness of fit testing?

- a) Normal distribution
- b) Chisquared distribution
- c) Gamma distribution
- d) Poission distribution

B

5. Which of the following distributions is Continuous

- a) Binomial Distribution
- b) Hypergeometric Distribution
- c) F Distribution
- d) Poisson Distribution

F

6. A statement made about a population for testing purpose is called?

- a) Statistic
- b) Hypothesis
- c) Level of Significance
- d) TestStatistic

B

7. If the assumed hypothesis is tested for rejection considering it to be true is called?

- a) Null Hypothesis
- b) Statistical Hypothesis

- c) Simple Hypothesis
- d) Composite Hypothesis

A

8. If the Critical region is evenly distributed then the test is referred as?

- a) Two tailed
- b) One tailed
- c) Three tailed
- d) Zero tailed

A

9. Alternative Hypothesis is also called as?

- a) Composite hypothesis
- b) Research Hypothesis
- c) Simple Hypothesis
- d) Null Hypothesis

→ B

10. In a Binomial Distribution, if 'n' is the number of trials and 'p' is the probability of success, then the mean value is given by  
a) np  
b) n

➔ A