



Bookmarks

▶ Machine Learning Course: Getting Started

▶ Week 1

▶ Week 2

▶ Week 3

▶ Week 4

▶ Week 5

▶ Week 6

▶ Week 7

▶ Week 8

▶ Week 9

▼ Week 10

Lecture 19
Principal Component Analysis

Lecture 20 Markov Models, Ranking and Semi-supervised Classification Examples

Week 10 Quiz

Quiz due Apr 11, 2017 07:30 MYT

Week 10 Discussion

Week 10 > Week 10 Quiz > Week 10 Quiz

Week 10 Quiz

🔖 Bookmark this page

Multiple Choice

1/1 point (graded)

Principle component analysis projects data into a lower dimension by minimizing _____

☐ the dimensionality of the projection

☒ the sum of squared errors of the projection ✓

☐ the absolut errors of the projection

☐ the total number of projections

Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)

Checkboxes

1/1 point (graded)

Check all that apply: PCA can be done using _____

☒ an eigendecomposition

☐ unconstrained least squares minimization

☐ random projections

☒ the singular value decomposition



Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)

Multiple Choice

1/1 point (graded)

True or False: The first principle component selects the direction of greatest variation in the data.

☒ TRUE ✓

☐ FALSE

Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)

Text Input

1/1 point (graded)

Probabilistic PCA introduces the _____ distribution to the PCA modeling framework.

Gaussian ✓

Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)

Multiple Choice

1/1 point (graded)

True or False: Kernel PCA must project data into a lower dimensional space than that of the original data.

☐ TRUE

☒ FALSE ✓

Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)

Text Input

1/1 point (graded)

A second-order Markov chain uses the previous ____ observations when making a prediction of the next observation.



Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)

Numerical Input

1/1 point (graded)

Enter the correct number: A 10x10 matrix can encode the transition probabilities of a ____-state Markov chain.



10

Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)

Numerical Input

1/1 point (graded)

The stationary distribution of a Markov chain is related to which eigenvector? Enter the index of this vector.



1

Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)

Multiple Choice

1/1 point (graded)

True or False: An iterative algorithm is required to find the maximum likelihood estimate of the transition matrix associated with an observed Markov chain.

☐ TRUE

☒ FALSE ✓

Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)

Multiple Choice

1/1 point (graded)

True or False: In the context of the lecture discussion, a random walk on a graph containing absorbing states is guaranteed to eventually terminate at one of these absorbing states.

☒ TRUE ✓

☐ FALSE

Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)

© All Rights Reserved



© 2012-2017 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

POWERED BY
OPEN EDX

