


RESEARCH ARTICLE

A full Bayesian model to handle structural ones and missingness in economic evaluations from individual-level data

Andrea Gabrio¹  | Alexina J. Mason² | Gianluca Baio¹

¹Department of Statistical Science,
University College London, London, UK

²Department of Health Services Research
and Policy, London School of Hygiene &
Tropical Medicine, London, UK

Correspondence

Andrea Gabrio, Department of Statistical
Science, University College London,
London WC1E 6BT, UK.
Email: ucakgab@ucl.ac.uk

Funding information

The Foundation BLANCEFLOR
Boncompagni Ludovisi, née Bildt; Mapi
Group

Economic evaluations from individual-level data are an important component of the process of technology appraisal, with a view to informing resource allocation decisions. A critical problem in these analyses is that both effectiveness and cost data typically present some complexity (eg, nonnormality, spikes, and missingness) that should be addressed using appropriate methods. However, in routine analyses, standardised approaches are typically used, possibly leading to biased inferences. We present a general Bayesian framework that can handle the complexity. We show the benefits of using our approach with a motivating example, the MenSS trial, for which there are spikes at one in the effectiveness and missingness in both outcomes. We contrast a set of increasingly complex models and perform sensitivity analysis to assess the robustness of the conclusions to a range of plausible missingness assumptions. We demonstrate the flexibility of our approach with a second example, the PBS trial, and extend the framework to accommodate the characteristics of the data in this study. This paper highlights the importance of adopting a comprehensive modelling approach to economic evaluations and the strategic advantages of building these complex models within a Bayesian framework.

KEYWORDS

Bayesian statistics, economic evaluations, hurdle models, missing data

1 | INTRODUCTION

Economic evaluation alongside randomised clinical trials (RCTs) is an important and increasingly popular component of the process of technology appraisal.¹ The typical analysis of individual-level data involves the comparison of two interventions for which suitable measures of clinical benefits and costs are observed on each patient enrolled in the trial, often at different time points throughout the follow-up. For simplicity, we generically term the clinical benefits as *effectiveness* and, thus, indicate the economic outcome variables as (e , c).

Typically, effectiveness is measured through multiattribute utility instruments (eg, the EQ-5D 3L: <http://www.euroqol.org>), the costs are obtained using clinic resource use records and both are summarised into cross-sectional quantities, eg, quality-adjusted life years (QALYs). The main objective of the economic analysis is to combine the population average effectiveness and costs and use the precepts of decision theory to determine the most “cost-effective” intervention, given

current evidence, as well as to assess the uncertainty in the decision-making process, induced by the uncertainty in the model inputs.²⁻¹⁰

In routine analyses, trial-based cost-effectiveness analyses (CEAs) are usually performed under a frequentist approach in which the two outcome variables (e , c) are modelled independently. Baseline adjustments are often included in the model using simple regression analyses.¹¹⁻¹³ This, often implicitly, assumes normality for the underlying cost and effectiveness data, or at least that the sample size is large enough for the means to be (approximately) normally distributed. In addition, almost invariably, the relationship between the outcomes and the baseline characteristics is assumed to be linear.

There are several potential issues with this setting: Firstly, the assumption of independence between costs and effectiveness is often questionable. While this is a recognised problem in the CEA literature, particularly under a Bayesian framework,^{7,14,15} and although it may introduce bias in the statistical modelling and, a fortiori, in the economic evaluation,^{14,16} appropriate methods to deal with correlation have historically found little application in routine analyses.¹⁷

Secondly, because both costs and effectiveness are usually characterised by a large degree of skewness, the assumption of normality is unlikely to hold and alternative approaches have been proposed in the literature. Examples include nonparametric bootstrapping¹⁸ and, particularly within a Bayesian approach, the use of more appropriate parametric modelling.^{15,16} Nonparametric bootstrapping mostly relies on using simple averages that often give similar results to those assuming normality.¹⁹ Conversely, modelling based on different parametric distributions (eg, Gamma for the costs and Beta for the QALYs) often allows improvement in the model fit to the observed data and appropriately captures skewness.

Thirdly, data may exhibit spikes at one or both of the boundaries of the range for the underlying distribution. For example, some patients in a trial may not accrue any cost at all (ie, $c_i = 0$), thus invalidating the assumptions for the Gamma distribution, which is defined on the range $(0, +\infty)$. Similarly, we may observe individuals who are associated with perfect health, ie, unit QALY,²⁰ which makes it difficult to use a Beta distribution, defined on the open interval $(0, 1)$. A simple solution is to add/subtract a small constant ϵ to the entire set of observed values for the cost/effectiveness variable, thus artificially rescaling it in the desired interval.²¹ Despite being very easy to implement, this strategy is potentially problematic as the results are likely to be strongly affected by the actual choice of the scaling parameter ϵ and no clear guideline exists about the value to use (eg, 0.1, 0.01, ...). In addition, when the proportion of these values is substantial, they may induce high skewness in the data and the application of simple methods may lead to biased inferences.²² A more efficient solution suggested to handle this issue is the application of *hurdle models*.²²⁻²⁴ These are mixture models defined by two components: the first one is a mass distribution at the spike, whereas the second is a parametric model applied to the natural range of the relevant variable. Usually, a logistic regression is used to estimate the probability of incurring a “structural” value (eg, 0 for the costs, or 1 for the QALYs); this is then used to weight the mean of the *nonstructural* values estimated in the second component. Hurdle models have been discussed and applied in CEA mainly for handling structural zero costs.²⁴⁻²⁶

Finally, individual-level data from RCTs are almost invariably affected by the problem of missing data. Numerous methods are available for handling missingness in the wider statistical literature, each relying on specific assumptions whose validity must be assessed on a case-by-case basis. While some guidelines exist for performing CEAs in the presence of missing outcome values,²⁷ they tend not to be consistently followed in published studies.²⁸⁻³¹ Analyses that are limited to the observed data (complete case analysis) are inefficient and may yield biased inferences.^{26,32-34} Multiple imputation (MI³⁵) is a more flexible method, which increasingly represents the *de facto* standard in clinical studies.^{36,37} In a nutshell, MI proceeds by replacing each missing data point with a value simulated from a suitable model. M complete (ie, without missing data) replicates of the original data set are thus created, each of which is then analysed separately using standard methods. The individual estimates are pooled using meta-analytic tools such as *Rubin's rules*,³⁵ to reflect the inherent uncertainty in imputing the missing values. For historical reasons, as well as on the basis of theoretical considerations, the number of replicated data sets M is usually in the range 5-10.^{35,38,39}

As a consequence of the separation between the imputation and the analysis steps, MI requires the property of *congeniality*, ie, the imputation model needs to be specified as equally or less restrictive than the analysis model.⁴⁰ In addition, in many applications, MI is based upon assuming a *missing-at-random* (MAR) mechanism, ie, the observed data can explain fully the reason for why some observations are missing. However, this may not be reasonable in practice (eg, for self-reported questionnaire data) and it is important to explore whether the resulting inferences are robust to a range of plausible *missing-not-at-random* (MNAR) mechanisms, which cannot be explained fully by the observed data. Neither MAR nor MNAR assumptions can be tested using the available data alone, and thus, it is crucial to perform sensitivity analysis to explore how variations in assumptions about the missing values impact the results.^{41,42}

Building on the existent literature, we show how models that simultaneously account for different potential sources of bias can be efficiently specified under a full Bayesian parametric framework, which has several advantages in comparison to a frequentist counterpart, specifically in health care technology assessments.^{7,10} Firstly, by virtue of its modular nature, Bayesian modelling is very flexible, which means that a basic structure can be relatively easily extended to account for the increasing complexity required to formally allow for the several features described above. We exploit this in Section 3. In addition, the Bayesian approach naturally allows for the principled incorporation of external evidence (eg, expert opinions) through the use of prior distributions. This is often crucial for conducting sensitivity analysis to a plausible range of missingness assumptions including MNAR,^{43,44} particularly when the evidence produced by the current study is limited, as is the case for small pilot trials, whose objective is to aid decision making about larger investments. Examples include the conduct of a full-scale trial, or the introduction in the market of a new cancer drug, based on the extrapolation of survival data produced over a short follow-up.

Moreover, we note that MI can be considered as an approximation to a full Bayesian analysis on different levels. First, MI separates the imputation and analysis steps in two estimation procedures while, within a full Bayesian approach, the parameters of interest are estimated simultaneously with the imputation of the missing values and no additional analysis or *ad hoc* pooling is necessary. Even though it has been shown that MI performs well in most standard situations, when the complexity of the analysis increases, a full Bayesian approach is likely to be a preferable option as it naturally allows the propagation of uncertainty to the wider economic model and to perform sensitivity analysis. Second, due to the small number of replicates that are kept in practice, MI can be thought of as a fully Bayesian analysis based on a few simulations. Interestingly, the often-quoted objection to Bayesian modelling, ie, that it is too computationally intensive in comparison to simpler frequentist counterparts, is likely to dissolve in the presence of extremely complex models, which would require tailor-made routines for the optimisation of nonstandard multivariate likelihood functions, thus effectively surrendering their computational advantage over intensive but efficient sampling methods such as Markov Chain Monte Carlo (MCMC).

The main contribution of this work is to provide a unified framework that allows jointly tackling the features in CEA discussed above. We use a real case study based on a small pilot trial as our motivating example. Starting from the original analysis, we progressively expand our basic model. We specifically focus on appropriately modelling spikes at the boundary and missingness, as they have substantial implications in terms of inferences and, crucially, cost-effectiveness results. We then use another case study to show the flexibility of our approach and accommodate the specific characteristics of the data in this second trial.

The paper is structured as follows. In Section 2, we present the case study used as motivating example and describe the data. Section 3 defines the general structure of the statistical model used in the analyses and how it can be tailored to deal with the different features affecting the data. Initially, for simplicity, we present each model under a complete case scenario that will then be extended to account for missingness. Section 4 compares the results from three alternative models under both a complete cases and all cases scenario assuming MAR. The robustness of the results to alternative MNAR assumptions is then explored. Section 5 summarises the inferences for each model from a decision-maker perspective and compares their implications in terms of cost effectiveness. Section 6 presents the second case study and the models used in the analysis, and summarises the results of the economic evaluation. Section 7 discusses the proposed framework and suggests some improvements for future work. Finally, the Appendix includes additional material related to model assessment and the computer code for our analysis.

2 | MOTIVATING EXAMPLE: THE MenSS TRIAL

We use as a motivating example the MenSS trial,⁴⁵ a pilot RCT conducted in the UK public care sector to evaluate the cost effectiveness of a new interactive digital intervention (the Men's Safer Sex [MenSS] website). This new intervention provides individually tailored advice on barriers to condom use to reduce the incidence of sexually transmitted infections (STIs) in young men. A total of 159 men aged 16 years and over with female sexual partners and recent unprotected sex or suspected acute STI were recruited from three English sexual health clinics. Participants were randomised to receive either usual clinical care only (comparator, $n_1 = 75$), or a combination of usual care and the MenSS website (active intervention, $n_2 = 84$). Sexual health-related resource use was collected via participant responses to questionnaires at 3, 6, and 12 months. For each individual i , utility scores u_{ij} were computed based on a generic health-related quality-of-life questionnaire, the EQ-5D 3L, collected at baseline ($j = 0$) and, then, at 3, 6, and 12 months ($j = 1, 2, J = 3$). Quality-adjusted life years and total costs (measured in pounds) were calculated by combining the utilities u_{ij} and costs c_{ij} collected at each




FIGURE 1 Histograms of the distributions of the complete case quality-adjusted life years (QALYs) and costs, expressed in pounds, in the control (A-B) and intervention (C-D) groups. For both variables and in both arms, skewness of the observed data is apparent. A, QALYs (control); B, Costs (control); C, QALYs (intervention); D, Costs (intervention)

time point as

$$e_i = \sum_{j=1}^J (u_{ij} + u_{i,j-1}) \frac{\delta_j}{2} \quad \text{and} \quad c_i = \sum_{j=1}^J c_{ij}, \quad (1)$$

where $\delta_j = \frac{\text{Time}_j - \text{Time}_{j-1}}{\text{Unit of time}}$ is the fraction of the time unit (12 months) between consecutive measurements, eg, $\delta_2 = (6 \text{ months} - 3 \text{ months}) / 12 \text{ months} = 0.25$. For the utilities, this approach is often referred to as the *area under the curve*.⁴⁶

The number of participants completing utility and cost questionnaires at every time point was 27 (36%) and 19 (23%) for the control and intervention groups, respectively. Figure 1 shows the histograms of the distributions of the complete case QALYs and costs in the control (panels A and B) and intervention (panels C and D) groups, respectively.

The data clearly present some of the features described in Section 1. A relatively high degree of skewness characterises the empirical distributions of QALYs and costs in both treatment groups. In particular, the substantial proportion of individuals incurring a perfect health status (which we term “structural ones”) observed in both control (33%) and intervention (42%) groups effectively induces spikes at 1 in the QALYs. Finally, a large proportion of missingness characterises both cost and utility data due to poor follow-up rates. With the exception of baseline data, which are only collected for the utilities, at each time point, the two outcomes are either both observed or both missing. A statistical summary of the observed cases and missingness levels for the utility and cost variables by follow-up period is shown in Table 1.

TABLE 1 Number and proportion of observed cases at each time point for the utility and cost data (self-recorded questionnaires), presented by the trial group (baseline data only related to the utilities). The number of individuals having valid data at each time point (complete cases) is also reported at the bottom of the table. Over the trial period, both dropout missingness and intermittent missingness occur; at each time point, only unit nonresponse is observed

Time	Type of Outcome	Control ($n_1 = 75$) observed (%)	Intervention ($n_2 = 84$) observed (%)
Baseline	Utilities	72 (96%)	72 (86%)
3 months	Utilities and costs	34 (45%)	23 (27%)
6 months	Utilities and costs	35 (47%)	23 (27%)
12 months	Utilities and costs	43 (57%)	36 (43%)
Complete cases	Utilities and costs	27 (36%)	19 (23%)

The original analysis was performed under a frequentist approach using standard ordinary least squares (OLS) regression.⁴⁵ Baseline utility regression adjustment was incorporated in the model, assuming MAR for all variables and restricting the analysis to the complete cases. However, most of the features described in Section 1 were not explicitly taken into account.

3 | MODELLING FRAMEWORK

In this section, we firstly present our general modelling framework for cost-effectiveness data. The model improves the typical approach used in routine analyses by accounting for correlation between the outcomes. Then, it is extended to handle structural values and missing data using three alternative specifications with increasing complexity. Throughout, we refer to our motivating example to demonstrate the flexibility of our full Bayesian approach in dealing with the idiosyncrasies highlighted above; we also note that these are likely to be encountered in many practical cases, thus making our arguments applicable in general.

Assume that some patient-level data are collected from a trial on $i = 1, \dots, n$ individuals who are randomly allocated to either a control ($t = 1$) or intervention ($t = 2$) group, with sample sizes n_1 and n_2 , respectively. We denote by e_{it} and c_{it} the effectiveness and cost outcome variables for the i th person in group t of the trial. To simplify the notation, unless necessary, we suppress the treatment subscript t .

To account for correlation between the outcomes, in general, we can specify the joint distribution $p(e, c)$ as

$$p(e, c) = p(c)p(e | c) = p(e)p(c | e), \quad (2)$$

where, for example, $p(e)$ is the *marginal* distribution of the effectiveness and $p(c | e)$ is the *conditional* distribution of the costs given the effectiveness.¹⁵ Note that, while it is possible to use interchangeably either factorisation in Equation 2, without loss of generality, we describe our analysis in the following through a marginal distribution for the effectiveness (QALYs) and a conditional distribution for the costs.

For each individual, we consider a marginal distribution $p(e_i | \theta_e)$ indexed by a set of parameters θ_e comprising a location ϕ_{ie} and a set of *ancillary* parameters ψ_e typically including some measure of *marginal* variance, σ_e^2 . We can model the location parameter using a generalised linear structure, eg,

$$g_e(\phi_{ie}) = \alpha_0 [+ \dots],$$

where α_0 is the intercept and the notation $[+ \dots]$ indicates that other terms (eg, quantifying the effect of relevant covariates) may or may not be included in the model. In the absence of covariates or assuming that a centred version $x_i^* = (x_i - \bar{x})$ is used, the parameter $\mu_e = g_e^{-1}(\alpha_0)$ represents the population average effectiveness.

For the costs, we consider a conditional model $p(c_i | e_i, \theta_c)$, which explicitly depends on the effectiveness variable, as well as on a set of quantities θ_c , again comprising a location and ancillary parameters. Note that, in this case, ψ_c includes a *conditional* variance τ_c^2 , which can be typically expressed as a function of the marginal variance σ_e^2 .^{7,15} The location can be modelled as a function of the effectiveness variable as

$$g_c(\phi_{ic}) = \beta_0 + \beta_1(e_i - \mu_e) [+ \dots].$$




FIGURE 2 Joint distribution $p(e, c)$, expressed in terms of a marginal distribution for the effectiveness and a conditional distribution for the costs, respectively indicated with a solid red line and a dashed blue line. The parameters indexing the corresponding distributions or modules are indicated with different Greek letters, whereas i denotes the individual index. The solid black and magenta arrows show the dependence relationships between the parameters within and between the two models, respectively. The dashed magenta arrow indicates that the ancillary parameters of the cost model may be expressed as a function of the corresponding effectiveness parameters. The dots enclosed in the square brackets indicate the potential inclusion of other covariates at the mean level for both modules [Colour figure can be viewed at wileyonlinelibrary.com]

Here, $(e_i - \mu_e)$ is the centred version of the effectiveness variable, while β_1 quantifies the correlation between costs and effectiveness. Assuming other covariates are also either centred or absent, $\mu_c = g_c^{-1}(\beta_0)$ is the population average cost.

Figure 2 shows a graphical representation of the general modelling framework described above. The effectiveness and cost distributions are represented in terms of combined “modules”—the blue and the red boxes—in which the random quantities are linked through logical relationships. This ensures the full characterisation of the uncertainty for each variable in the model. Notably, this is general enough to be extended to any suitable distributional assumption, as well as to handle covariates in either or both the modules.

In the rest of the section, we present three alternative specifications of the general structure depicted in Figure 2 to model effectiveness and cost data. These are (1) Normal marginal for the effectiveness and Normal conditional for the costs (which is identical to a Bivariate Normal distribution for the two outcomes), (2) Beta marginal for the effectiveness and Gamma conditional for the costs, and (3) hurdle model. First, we present each assuming a complete cases scenario and then extend the structure to all cases. The latter scenario includes the complete cases and additionally imputes the outcome values for all the remaining individuals in the trial, either under MAR (for all models) or alternative MNAR scenarios (for the hurdle model only).

3.1 | Complete cases

3.1.1 | Bivariate Normal

Arguably, the easiest way of jointly modelling two variables is to assume Bivariate normality, which in our context can be factorised into marginal and conditional Normal distributions for e_i and $c_i | e_i$. This is the closest modelling structure to those underpinning a typical frequentist analysis, while also accounting for potential correlation between the outcomes.

In line with current recommendations (and the original analysis of the MenSS trial), we adjust for the baseline utilities—using a centred version $(u_{i0} - \bar{u}_0)$. We model $e_i | \theta_e \sim \text{Normal}(\phi_{ie}, \sigma_e^2)$, using an identity link function for the location parameter

$$g_e(\phi_{ie}) = \phi_{ie} = \alpha_0 + \alpha_1(u_{i0} - \bar{u}_0).$$

Here, the parameter α_1 quantifies the impact of the centred baseline utilities on the QALYs, while $\mu_e = \alpha_0$ and σ_e^2 represent the marginal (population level) mean and variance, respectively.

As for the costs, we model $c_i | e_i, \theta_c \sim \text{Normal}(\phi_{ic}, \tau_c^2)$, where the conditional mean and variance are defined as

$$g_c(\phi_{ic}) = \phi_{ic} = \beta_0 + \beta_1(e_i - \mu_e) \quad \text{and} \quad \tau_c^2 = \sigma_c^2 - \sigma_e^2 \beta_1^2.$$

The model parameters are thus $\theta_e = (\alpha_0, \alpha_1, \sigma_e^2)$ and $\theta_c = (\beta_0, \beta_1, \mu_e, \sigma_c^2, \sigma_e^2)$ —note that the marginal mean and variance of the effectiveness link the two modules and therefore feature in both sets of parameters.

The model is completed by assigning suitable prior distributions to the elements of $\theta = (\theta_e, \theta_c)$; for example, independent Normal priors can be assumed for the regression parameters, while Uniform or Half-Cauchy priors can be assigned on the scale of the standard deviations.⁴⁷

3.1.2 | Beta-Gamma

The second model we consider assumes a Beta marginal for the QALYs and a Gamma conditional for the costs. In particular, we parameterise the Beta distribution in terms of the mean ϕ_{ie} and the scale parameter $\tau_{ie} = \left(\frac{\phi_{ie}(1-\phi_{ie})}{\sigma_e^2} - 1 \right)$ as $e_i | \theta_e \sim \text{Beta}(\phi_{ie}\tau_{ie}, (1-\phi_{ie})\tau_{ie})$ and model the location as

$$g_e(\phi_{ie}) = \text{logit}(\phi_{ie}) = \alpha_0 + \alpha_1(u_{i0} - \bar{u}_0).$$

The costs are modelled as $c_i | e_i, \theta_c \sim \text{Gamma}(\phi_{ic}\tau_{ic}, \tau_{ic})$, where the shape parameter is defined as the product of the location ϕ_{ic} and the rate τ_{ic} . The generalised linear model for the location is

$$g_c(\phi_{ic}) = \log(\phi_{ic}) = \beta_0 + \beta_1(e_i - \mu_e).$$

The marginal means for the QALYs and total costs can then be obtained using the respective inverse link functions

$$\mu_e = \frac{\exp(\alpha_0)}{1 + \exp(\alpha_0)} \quad \text{and} \quad \mu_c = \exp(\beta_0).$$

Notice that, in comparison to the Bivariate Normal of Section 3.1.1, the Beta-Gamma model reflects more closely the range and skewness of the observed data. Nevertheless, this modelling structure also fails to directly account for the structural values, eg, unit QALYs. In the presence of structural values, it is necessary to rescale the observed data, eg, by applying the Beta model to $e_i^* = e_i - \varepsilon$ for some $\varepsilon \rightarrow 0$.

The model is again completed by placing suitable priors on the parameters. For example, we can use independent Normal priors for the regression coefficients $(\alpha_0, \beta_0, \alpha_1, \beta_1)$ and a Uniform or Half-Cauchy prior for σ_e . Notice, however, that a little more care is needed in defining a prior distribution for σ_e . In fact, by the mathematical properties of the Beta distribution, the variance is bounded by a function of the mean

$$\sigma_e^2 \leq \mu_e(1 - \mu_e) = v.$$

Consequently, we can place an informative prior on the standard deviation $\sigma_e \sim \text{Uniform}(0, \sqrt{v})$, which coupled with a prior for μ_e , induces a suitable prior for τ_e as well. Note that, even by starting with vague distributions for (μ_e, σ_e) , the resulting prior for the Beta scale τ_e may not be vague at all.

3.1.3 | Hurdle model

To overcome the limitations of the model in Section 3.1.2 in terms of the structural ones, we expand it to a hurdle version. Specifically, for each subject in the trial $i = 1, \dots, n$, we define an indicator variable d_{ie} taking value 1 if the i th individual is associated with a unit QALYs ($e_i = 1$) and 0 otherwise ($e_i < 1$). This is then modelled as

$$d_{ie} := \mathbb{I}(e_i = 1) \sim \text{Bernoulli}(\pi_{ie})$$

$$\text{logit}(\pi_{ie}) = \gamma_0 + \gamma_1(u_{i0} - \bar{u}_0) [+ \dots], \quad (3)$$

where π_{ie} is the individual probability of unit QALYs, which is estimated on the logit scale as a function of a baseline parameter γ_0 and the centred baseline utilities, whose effect is captured by the parameter γ_1 . As for the effectiveness and cost models, other covariates can be additively included in the model of d_{ie} . We specifically distinguish the baseline utilities from any other covariate as they are likely to be particularly informative in predicting whether an individual is associated with a structural one in the QALYs. All the logistic regression parameters should be given suitable prior probability distributions (eg, Normal). Within this framework, the quantity

$$\bar{\pi}_e = \frac{\exp(\gamma_0)}{1 + \exp(\gamma_0)} \quad (4)$$

represents the estimated marginal probability of unit QALYs. The parameters $\bar{\pi}_e$ and $(1 - \bar{\pi}_e)$ in effect represent the weights used to mix the two components.

Depending on the value of d_{ie} , we can partition the observed data on the QALYs into two subsets. In the first subset, defined as the n^1 subjects for whom $d_{ie} = 1$, we define a variable $e_i^1 = 1$. Conversely, the second subset consists of the $n^{<1} = (n - n^1)$ subjects for whom $d_{ie} = 0$, and for these individuals, we define a variable $e_i^{<1}$. Because this is less than 1, we can model it directly using a Beta distribution characterised by an overall mean $\mu_e^{<1}$, in line with the specification we

have shown in Section 3.1.2. Using the estimated value for $\bar{\pi}_e$ from Equation 4, we can compute the overall population average effectiveness measure in both treatment groups μ_{et} as the linear combination

$$\mu_{et} = (1 - \bar{\pi}_{et})\mu_{et}^{<1} + \bar{\pi}_{et}.$$

In the absence of structural zeros, the conditional model for the costs is exactly as specified in Section 3.1.2.

3.2 | All cases

When missingness occurs in the QALYs and cost variables, no change to the model structure is required under MAR for both the Bivariate Normal and Beta-Gamma specifications. For the hurdle model, when e_i is missing, it is not possible to directly define the value for d_{ie} . However, unit QALYs can only be observed if $u_{ij} = 1$ for all time points $j = 0, \dots, J$. Consequently, if an individual i is such that u_{ij} is missing at some time point j and $u_{ij} \neq 1$ at any other time point, then by necessity $d_{ie} = 0$. However, for all individuals having $u_{ij} = 1$ at all observed time points but with at least one missing value at some other time point, then d_{ie} is unknown.

Incomplete covariates need to be explicitly modelled to impute their missing values. For simplicity, we consider the case where the only covariate included in the model is the baseline utility; however, the same approach can be extended to any other type of partially observed covariates. In the Bivariate Normal and the Beta-Gamma formulations, we can handle missingness in u_{i0} by assuming a suitable model. One simple choice is to assume the same distribution for u_{i0} as for the outcome e_i , ie, Normal or Beta, respectively, eg, Appendix A shows the implementation for the Beta-Gamma model.

Similarly to Section 3.1.3, we can formulate another hurdle model for u_{i0} . More specifically, first we specify a model for the individuals with a nonunit utility value. Again, a simple solution is to base this on the same distributions assumed for the QALYs. Secondly, we estimate the probability of observing a structural one in the utilities as

$$\begin{aligned} d_{iu} &:= \mathbb{I}(u_i = 1) \sim \text{Bernoulli}(\pi_{iu}) \\ \text{logit}(\pi_{iu}) &= \eta_0 [+ \dots], \end{aligned}$$

where d_{iu} is the indicator variable for the structural ones in the baseline utilities.

3.2.1 | Sensitivity analysis (MNAR)

Finally, hurdle models also offer a convenient framework for exploring the robustness of the results to some departures from MAR and, therefore, allow to perform a simple type of sensitivity analysis to the missingness assumptions. Two relevant cases are

- (a) the individuals for whom utility values are missing throughout the follow-up, ie, $u_{ij} = \text{NA}$ for all $j = 1, \dots, J$;
- (b) the individuals for whom all the observed utilities are equal to 1, but with at least one time point j at which $u_{ij} = \text{NA}$.

For both these cases, it is impossible to compute the value of the indicator d_{ie} according to the information from the observed data. However, we can arbitrarily set the value of d_{ie} to either 1 or 0 using different configurations, eg, by varying the number of structural values potentially observed in a given scenario. Because these configurations are based on assumptions about the missing values that cannot be verified from the data at hand (but are in fact arbitrarily set by the experimenter), they effectively represent a way to assess the robustness of the results to some departures from MAR.

In the MenSS trial, there are $n_1^* = 13$ (12%) individuals in the control and $n_2^* = 22$ (26%) in the intervention group who fall within case *a* or *b*. Thus, we perform sensitivity analysis by defining a set of alternative MNAR assumption scenarios for these individuals and assess the robustness of the results across them. The four different scenarios considered are summarised in Table 2:

We choose these scenarios in order to assess how different “extreme” combinations of the number of potential structural ones in the intervention and control group can impact the results.

4 | RESULTS

We fitted all models using JAGS,⁴⁸ a software specifically designed for the analysis of Bayesian models using Markov Chain Monte Carlo (MCMC) simulation, which can be interfaced with R through the package R2jags.⁴⁹ Samples from the posterior distribution of the parameters of interest generated by JAGS and saved to the R workspace are then used to

TABLE 2 Alternative (missing-not-at-random) MNAR scenarios considered in the Men's Safer Sex study for the hurdle model. In each scenario, individuals who are potentially associated with a unit QALYs in the control ($n_1^* = 13$) and intervention ($n_2^* = 22$) groups are assigned to either the structural or nonstructural components by setting the value of the indicator d_{ie} equal to 1 or 0, respectively

Scenario	Control ($n_1^* = 13$)	Intervention ($n_2^* = 22$)
MNAR1	$d_{ie} = 1$	$d_{ie} = 1$
MNAR2	$d_{ie} = 0$	$d_{ie} = 0$
MNAR3	$d_{ie} = 1$	$d_{ie} = 0$
MNAR4	$d_{ie} = 0$	$d_{ie} = 1$

Note. QALYs, quality-adjusted life years.

produce summary statistics and plots. We ran two chains with 20 000 iterations per chain, using a burn-in of 10 000, for a total sample of 20 000 iterations for posterior inference. For each unknown quantity in the model, we assessed convergence and autocorrelation of the MCMC simulations using diagnostic measures such as the *potential scale reduction factor* and the *effective sample size*.⁵⁰ The total running time required for the models to produce representative samples from the posterior distributions of interest ranged from 5 to 10 minutes.

Alternative prior distributions are considered to check that we are not incorporating any unintended information into the models through the priors. For example, we varied the upper boundary of the uniform distribution and specified alternative prior forms (Half-Cauchy and Half-Normal) for the standard deviations or chose different values for the variance of normally distributed regression parameters. Results were robust to these specifications. Although the hurdle model as described in Section 3.1.3 cannot be directly written in JAGS, it can be implemented using a simple “coding trick.” Appendix A presents all the technical details and the JAGS script.

4.1 | Complete and all cases (MAR)

Following the original analysis, we first consider only the complete cases and adjust for baseline utilities at the mean level for the QALYs in each model. We initially include in each model three more covariates (age, ethnicity and employment status). However, because posterior results for mean QALYs and costs were almost unaffected by these variables, we simplify the models and retain the covariates only in the linear predictor of Equation 3 to estimate the probability of structural ones in the hurdle model. We then extend the framework to all cases under MAR, where the baseline utilities are explicitly modelled as detailed in Section 3.2 and again as functions of age, ethnicity, and employment status. We report the results under MAR as this is the default (and often implicit) missingness assumption used by practitioners in routine analyses. We then conduct a sensitivity analysis to assess the robustness of the results to alternative departures from MAR.

Figure 3 shows the posterior distributions of the mean QALYs and costs for both treatment groups under a complete (red) and all (blue) cases scenario for each model, assuming MAR.

The posterior distributions of the mean QALYs (panels A and B) present some discrepancies between the complete and all cases scenarios, with magnitude varying according to the treatment group and model considered. In general, the results for all cases are lower in the control group and higher in the intervention group in comparison to those obtained using the complete cases. As for the mean costs (panels C and D), the results associated with a Gamma distribution are substantially more skewed compared with those obtaining using the Normal model, especially in the intervention group. In addition, the Gamma model typically leads to mean estimates that are systematically lower under the all cases scenario.

We compare the fit of the different models using the deviance information criterion (DIC).⁵¹ The DIC is a measure of comparative predictive ability based on the model deviance and a penalty for model complexity. When comparing a set of models based on the same data, the one associated with the lowest DIC is the best-performing, amongst those assessed. DIC is not uniquely defined in the presence of missing data and its use and interpretation are not straightforward.^{52,53} No measure of model comparison can provide a full picture of model fit when dealing with partially observed data. We can compare models based on the fit to the observed data alone, but we must remember that this provides no information about the fit to the unobserved data. In our analysis, we consider a DIC based on the observed data and calculated only




FIGURE 3 Posterior distributions for the marginal mean parameters of the quality-adjusted life years (QALYs) (A-B) and cost variables (C-D), expressed in pounds, in each group of the trial under either a complete (red) or all (blue) cases scenario. The posterior results are presented for all model specifications considered (Bivariate Normal, Beta-Gamma, and hurdle model), and for each of these, the posterior mean estimates and associated 90% highest posterior density (HPD) interval bounds are reported. A, QALYs (control); B, QALYs (intervention); C, Costs (control); D, Costs (intervention) [Colour figure can be viewed at wileyonlinelibrary.com]

for the modules that are in common between the models, ie, excluding the contribution from the structural indicators for the hurdle model. In our analysis, we consider a DIC based on the observed data and calculated only for the modules that are in common between the models, ie, excluding the contribution from the structural indicators for the hurdle model. The Bivariate Normal model is always associated with the highest DIC under both a complete and all cases scenarios (536 and 445). The Beta-Gamma (386 and 60) and, especially, the hurdle model (−50 and −2419) substantially improve the model fit to the observed data.




FIGURE 4 Imputed quality-adjusted life years (QALYs) in the control and intervention groups based on the Bivariate Normal, Beta-Gamma, and hurdle model. Imputations are summarised in terms of posterior means and 90% highest posterior density intervals (coloured dots and lines), whereas an x symbol is used to denote the observed cases. Imputed values are also distinguished according to whether the corresponding baseline utilities were either observed (blue) or missing (red). The solid black line represents the upper bound for the utilities, set at the value of 1. A, Bivariate Normal (control); B, Bivariate Normal (intervention); C, Beta-Gamma (control); D, Beta-Gamma (intervention); E, Hurdle model (control); F, Hurdle model (intervention) [Colour figure can be viewed at wileyonlinelibrary.com]

4.1.1 | Imputations under MAR

Figure 4 depicts the observed QALYs in both treatment groups (indicated with black crosses) as well as summaries of the posterior distributions for the imputed values, obtained from each model. Imputations are distinguished based on whether the corresponding baseline utility value is observed or missing (blue or red lines and dots, respectively) and are summarised in terms of posterior mean and 90% highest posterior density (HPD) intervals.

There are clear differences in the imputed values and corresponding credible intervals between the three models in both treatment groups. Neither the Bivariate Normal nor the Beta-Gamma models produce imputed values that capture the structural one component in the data. In addition, as to be expected, the Bivariate Normal fails to respect the natural support for the observed QALYs, with many of the imputations exceeding the unit threshold bound. These unrealistic imputed values highlight the inadequacy of the Normal distribution for the data and may lead to distorted inferences.




FIGURE 5 Density strip plots for the posterior distributions of the probability of structural ones (A) and the marginal mean quality-adjusted life years (QALYs) (B) under missing-at-random (MAR) and four alternative missing-not-at-random (MNAR) scenarios. For each scenario, results are presented for the control (red) and the intervention (blue) groups. Mean posterior values and associated 90% highest posterior density (HPD) interval bounds are indicated with tick marks and reported aside for each quantity. A, Probability of structural ones; B, Mean QALYs [Colour figure can be viewed at wileyonlinelibrary.com]

Conversely, imputations under the hurdle model are more realistic, as they can replicate values in the whole range of the observed data, including the structural ones. Imputed unit QALYs with no discernible interval are only observed in the intervention group due to the original data composition, ie, individuals associated with a unit baseline utility and missing QALYs are almost exclusively present in the intervention group.

4.2 | Sensitivity analysis (MNAR)

For each of the alternative MNAR scenarios described in Section 3.2.1, as well as for the analysis under MAR, Figure 5 shows posterior density strips for the structural one probability $\bar{\pi}_e$ and the marginal mean QALYs μ_e , in the control (red) and intervention (blue) groups.

Estimates under MAR indicate that the new intervention is associated with a probability of observing a structural one and a mean QALYs that are on average higher compared with the control. Although similar results are obtained under MNAR1, the estimated quantities are highly unstable across the other three MNAR scenarios. Specifically, under MNAR2, the probability of structural ones is substantially reduced in both groups and induces a zero mean difference in the QALYs. Under MNAR3 and MNAR4, the differences between the estimated probabilities and mean QALYs in the two groups are increased in magnitude and lead to opposite mean differentials.

5 | ECONOMIC EVALUATION

We complete the analysis by assessing the cost effectiveness of the new intervention with respect to the control, comparing the results of the different models under MAR (Section 4.1) and the alternative MNAR scenarios explored for the hurdle model (Section 4.2). We specifically rely on the examination of the cost-effectiveness plane (CEP⁵⁴) and the cost-effectiveness acceptability curve (CEAC⁵⁵) to summarise the economic analysis.

The CEP (Figure 6A) is a graphical representation of the joint distribution for the population average effectiveness and costs increments, indicated respectively as $\Delta_e = (\mu_{e2} - \mu_{e1})$ and $\Delta_c = (\mu_{c2} - \mu_{c1})$, under the three model specifications




FIGURE 6 Cost-effectiveness planes (CEPs) (A) and cost-effectiveness acceptability curves (CEACs) (B) associated with the hurdle (blue dots and line), Bivariate Normal (red dots and line), and Beta-Gamma (green dots and line) models. In the CEPs, the incremental cost-effectiveness ratios (ICERs) based on the results from the three model specifications under missing at random (MAR) are indicated with corresponding darker-coloured dots, whereas the portion of the plane on the right-hand side of the straight line passing through the plot (evaluated at $k = £20\,000$) denotes the sustainability area. For the CEACs, in addition to the results under MAR (solid lines), the probability values for the four missing-not-at-random (MNAR) models described in Section 4.2 are represented with different types of dashed lines. A, Cost-effectiveness plane; B, Cost-effectiveness acceptability curve [Colour figure can be viewed at wileyonlinelibrary.com]

(light red for the Bivariate Normal, light green for the Beta-Gamma, and light blue for the hurdle model). The slope of the straight line crossing the plane is the “willingness to pay” threshold (often indicated as k) and can be considered as the amount of budget the decision maker is willing to spend to increase the health outcome of one unit and effectively used to trade clinical benefits for money. Points lying below this straight line fall in the so-called *sustainability area*⁷ and suggest that the active intervention is more cost effective than the control. This is because, in this area, the new intervention is either more effective and less expensive (in the South-Eastern quadrant) or it produces an increase in benefits that more than offsets the increase in the costs (points in the North-Eastern quadrant below the line). In the graph, which for simplicity only displays the results associated with the all cases under MAR, we also show the incremental cost-effectiveness ratio (ICER) computed under each model, as darker-coloured dots. This is defined as

$$\text{ICER} = \frac{E[\Delta_c]}{E[\Delta_e]},$$

and it quantifies the cost per incremental unit of QALYs. For all three models, more than 70% of the samples fall in the sustainability area and are associated with negative ICERs, suggesting that the intervention can be considered as cost effective by producing a QALYs gain at virtually no extra costs, or even saving money.

The CEAC (Figure 6B) is obtained by computing the proportion of points lying in the sustainability area upon varying the willingness to pay threshold k . Based on general recommended guidelines,¹ we consider a range for k up to £30 000 per QALY gained. The CEAC estimates the probability of cost effectiveness, thus providing a simple summary of the uncertainty associated with *optimal* decision making suggested by the ICER. For each model, the results under MAR are reported using solid lines with different colours, ie, red for the Bivariate Normal, green for the Beta-Gamma, and blue for the hurdle model. In addition, the results associated with the four MNAR scenarios are reported using different types of dashed lines. Under MAR, for the Bivariate Normal and Beta-Gamma models, the CEACs indicate the cost effectiveness of the new intervention with a probability above 0.8 for most values of k . Conversely, under the hurdle model, the curve is shifted downward by 0.24 and 0.16 with respect to the Bivariate Normal and Beta-Gamma models, respectively, and suggests a more uncertain conclusion. Perhaps unsurprisingly, none of these results is robust to the alternative MNAR

scenarios explored. The CEAC plot clearly shows a large sensitivity of the cost-effectiveness probability with respect to the assumed number of structural ones in both treatment groups. Indeed, the curves span a huge probability range from 0.2 under MNAR4 to 1 under MNAR3. This implies a considerable change in the output of the decision process and severely undermines the validity of the conclusions obtained under MAR.

6 | SECOND EXAMPLE: THE PBS TRIAL

The positive behaviour support (PBS⁵⁶) trial is a multicentre RCT involving community intellectual disability services and service users with mild to severe intellectual disability and challenging behaviour. Positive behaviour support is a multi-component intervention that is designed to foster prosocial actions and enhance the person's quality of life and his/her integration within the local community. Participants ($n = 244$) were enrolled from a total of $S = 23$ sites and randomly allocated to staff teams trained to deliver PBS in addition to treatment as usual (reference intervention, $n_2 = 108$), or to staff teams trained to deliver treatment as usual alone (comparator, $n_1 = 136$). Measures for quality of life (EQ-5D-3L) and health-related cost (family and paid carer records) were collected from each site s at baseline 6 and 12 months. Utility values lower than zero, representing health states "worse than death," are recorded for some individuals throughout the study. quality-adjusted life years and cost variables are computed as in Equation 1. The number of complete cases is 108 (79%) and 96 (89%) for the control and intervention group, respectively.

The data in the PBS trial share the same complexities found in the MenSS study. Both QALYs and costs are partially observed and have skewed empirical distributions, and some individuals in both treatment groups are associated with a structural one in the QALYs (5%). The original analysis ignored these features and was performed on the complete cases using standard regression methods, adjusting for differences at baseline utilities and costs, and accounting for the multilevel structure through random effects.

6.1 | Models

We show the flexibility of our framework by adapting the models in Section 3 to accommodate the characteristics of the PBS data. Specifically, we scale the QALYs to avoid negative values when using a Beta distribution; replace the Gamma distribution for the costs with a LogNormal distribution, which improves the fit to the observed data; and account for the multilevel structure.

We assess and compare the performance of three models. These are (1) Bivariate Normal for the two outcomes, (2) Beta marginal for the effectiveness and LogNormal conditional for the costs, and (3) hurdle model. All models account for the multilevel structure in the data and include three categorical covariates (living condition, level of disability, and type of carer) to estimate the mean QALYs and costs.

We first apply the models to the complete cases and then extend the analysis to all cases under MAR. No sensitivity analysis as in Section 3.2.1 can be performed for the PBS study because, for each missing individual, we observe a utility value that is lower than one at at least one time point, ie, none of them can be a structural one. In the next section, we present the specification of the Beta-LogNormal model to show how the framework in Section 3 has been modified to address the characteristics of the PBS data.

6.1.1 | Beta-LogNormal

The second model consists of a Beta marginal for the QALYs and a LogNormal conditional for the costs. Because the Beta distribution does not allow negative values, we scale the QALYs on $(0, 1)$ through the transformation $e_i^* = \frac{e_i - \min(e_i)}{\max(e_i) - \min(e_i)}$ and fit the model to these transformed variables. A similar scaling is applied to the baseline utilities u_{i0}^* when these are modelled in the all cases scenario.

The Beta distribution is parameterised as in Section 3.1.2, while the multilevel structure is accounted for through site-specific regression coefficients for the baseline utility and cost terms (varying-slope only model). We choose this specification as inference from a varying-intercept model was similar to that obtained from ignoring the multilevel structure, while the introduction of site-specific coefficients for the other covariate terms lead to almost identical results compared with the proposed approach.

The QALYs are modelled as $e_i^* | \theta_e \sim \text{Beta}(\phi_{ie}\tau_{ie}, (1 - \phi_{ie})\tau_{ie})$ with location

$$\text{logit}(\phi_{ie}) = \alpha_0 + \alpha_{1s}(u_{i0} - \bar{u}_0) [+ \dots],$$

where α_{1s} are the baseline utility structured coefficients associated with the $s = 1, \dots, S$ different sites.

The costs are modelled as $c_i | e_i^*, \theta_c \sim \text{LogNormal}(\phi_{ic}, \tau_c)$, where the mean and standard deviation parameters (ϕ_{ic}, τ_c) are defined on the log scale. Baseline costs (c_0) are included in the cost model using the same multilevel specification used for u_0 in the QALYs model. It is not possible to include the centred QALYs in the regression as modelling e_i^* on a transformed scale does not allow to identify the marginal mean μ_e as in Section 3.1.2. Consequently, the cost location is

$$\phi_{ic} = \beta_0 + \beta_1(e_i^*) + \beta_{2s}(c_{i0} - \bar{c}_0) [+ \dots],$$

where β_{2s} are the site-specific baseline cost regression coefficients.

We retrieve the marginal means on the natural scale for both outcomes through Monte Carlo integration. At each iteration of the posterior distribution for the model parameters in the MCMC output, we generate a large number of samples for e_i and c_i and take the expectation over these values to obtain Monte Carlo estimates of the marginal means μ_e and μ_c .

To complete the model, we specify Normal priors for the structured coefficients $\alpha_{1s} \sim \text{Normal}(0, \sigma_\alpha)$ and $\beta_{2s} \sim \text{Normal}(0, \sigma_\beta)$ with shared standard deviations σ_α and σ_β . Finally, we choose independent Normal priors for the other regression coefficients α and β and a Uniform or Half-Cauchy prior for the standard deviations.

6.2 | Results

All models are fitted and assessed using the same software configuration and diagnostic measures as in Section 4. Due to space constraints, we only present the economic results for each model in the PBS study for the all cases scenario under MAR. A comparison of the posterior inference between the models for both the complete and all cases scenarios, the imputed QALYs for each model under MAR, and a summary and visual description of the PBS data are available on the Web Appendix.




FIGURE 7 Cost-effectiveness planes (CEPs) (A) and cost-effectiveness acceptability curves (CEACs) (B) associated with the hurdle (blue dots and line), Bivariate Normal (red dots and line), and Beta-LogNormal (green dots and line) models. In the CEPs, the incremental cost-effectiveness ratios (ICERs) based on the results from the three model specifications under MAR are indicated with corresponding darker-coloured dots, whereas the portion of the plane on the right-hand side of the straight line passing through the plot (evaluated at $k = £20\,000$) denotes the sustainability area. A, Cost-effectiveness plane; B, Cost-effectiveness acceptability curve. MNAR, missing not at random [Colour figure can be viewed at wileyonlinelibrary.com]

Figure 7 shows the CEP and CEAC for the Bivariate Normal (red dots and line), Beta-LogNormal (green dots and line), and hurdle model (blue dots and line) for the all cases scenario under MAR.

For all three models, almost all samples in the CEPs (Figure 7A) are located in the North-Eastern quadrant and most of them fall in the sustainability area at a willingness to pay $k = £20\,000$. Results from the Bivariate Normal model suggests a higher cost effectiveness of the new intervention and are associated with a lower ICER compared with those of the other two models. This is reflected in the CEACs (Figure 7B), which show a probability of cost effectiveness that is on average 20% higher for the Bivariate Normal with respect to the Beta-LogNormal and hurdle model.

7 | DISCUSSION

In CEAs alongside RCTs, analysts typically rely on standard models that ignore or at best fail to properly account for potentially important features in the data, such as the correlation between costs and effectiveness, skewness in the distribution of the observed data, the presence of structural values, and almost invariably, missing data. In this paper, we have presented a general Bayesian framework that is able to overcome these problems. Our approach represents a considerable improvement with respect to the current practice and can be implemented in a relatively easy way using freely available software. This is a key advantage that can encourage practitioners to move away from likely biased methods and promote the use of our framework in routine analyses.

The analysis of both case studies shows notable variations in the results, compared with those of the original analyses. In the MenSS trial, accounting for the structural ones and missingness uncertainty has a considerable impact on the cost effectiveness of the new intervention and future research prioritisation. In particular, the sensitivity of the final conclusions to the missingness scenarios suggests that the results obtained under MAR are not robust to the departures explored. This is expected in light of the large proportion of missing values and the limited auxiliary information available. The original analysis ignored missingness uncertainty, and therefore, its conclusions are unlikely to provide a reliable picture about the economic assessment. Our analysis of the MenSS trial illustrates how a model that accounts for the complexities of the data can be specified using our framework and shows the difficulty of drawing any conclusion from this study given the limited evidence available and the lack of robustness of the results to the missing data assumptions. In the PBS trial, the results of the economic evaluation change substantially when skewness is accounted for. This second example demonstrates the flexibility of our framework and compares alternative model specifications in a setting where the data have a multilevel structure and the proportion of missing values is lower with respect to the MenSS trial. In both cases, the hurdle model represents the best model amongst those assessed as it captures both skewness and structural values, while the other specifications fail to deal with at least one of these features.

Our results are obtained with specific reference to the two case studies. However, the MenSS and PBS trials are very much representative of the “typical” data set used in CEAs alongside RCTs. Thus, it is highly likely that the same features (and potentially the same contradictions in the results, upon varying the complexity of the modelling assumptions) apply to many real cases. This is a very important, if somewhat overlooked problem, as it can thwart the validity of simplistic models that, while established amongst practitioners, may lead to misleading cost-effectiveness conclusions and bias the decision-making process.

Missing data pose a serious threat to the economic evaluation as, when confronted with a partially observed data set, each analysis makes assumptions about the missing values that cannot be verified from the data at hand. Any measure of fit or predictive accuracy, such as the DIC or posterior predictive checks,⁵⁰ can only provide information about the observed data and therefore tell just part of the story.^{52,53} Thus, the use of sensitivity analysis to explore the impact on the results of different plausible missingness assumptions, including MNAR, becomes essential. The Bayesian approach naturally allows to perform these assessments through the incorporation of external evidence (eg, expert opinions) in the model using prior distributions while ensuring consistency and the correct propagation of uncertainty throughout the model.

We have demonstrated one possible way of assessing the robustness of the results to some “extreme” scenarios that can be incorporated in the framework at no extra cost in terms of model complexity. If the results are not robust to the departures explored, these scenarios offer a starting point for further analyses where more advanced methods can be used to explicitly allow for the variability in the MNAR values, eg, selection models or pattern mixture models.⁴²⁻⁴⁴ Practitioners could incorporate strategies at the design stage of the trial to collect data from external sources, which can be used to guide the missingness assumptions in the statistical analysis. For example, additional information can be obtained through the elicitation of experts' knowledge or the inclusion of auxiliary variables that are thought to be predictive of missingness.

Finally, a potentially relevant question concerning missing variables derived from repeated questionnaires, eg, EQ-5D, is whether imputation should be carried out at the time scores (utilities) or total scores (QALYs) level. The performance of these alternative imputation strategies has only recently been compared in the health economic literature.⁵⁷⁻⁵⁹ The cross-sectional modelling framework used in trial-based economic evaluations has some limitations for handling missingness because it does not allow for the incorporation of the information from the partially observed utilities and costs in the study or the time dependence between the responses. These limitations can be overcome using a modelling framework that explicitly accounts for the longitudinal nature of the data. In future work, we plan to develop a longitudinal model to efficiently deal with missingness while also jointly accounting for the complexities of the data in economic evaluations. This, however, requires a radical change of the model paradigm currently used by practitioners and a level of statistical knowledge that would undermine the ease of implementation of our framework in routine analyses and therefore goes beyond the objective of this paper.

In conclusion, in this work, we have presented a flexible Bayesian analytic framework that can (a) jointly model costs and effectiveness, (b) account for skewness and structural values, and (c) assess the robustness of the results under a set of differing missingness assumptions. Routine analyses almost exclusively rely on methods that ignore at least some of the complexities of the data. The original contribution of this work consists in the joint implementation of methods that account for these complexities within a unique and flexible framework that is relatively easy to apply. It is important that the features of CEA individual-level data are simultaneously addressed to avoid biased results, which may in turn lead to misleading cost-effectiveness conclusions. The availability of methodological and practical tools such as the ones presented in this paper has the potential to improve the work of modellers and regulators alike, thus advancing the fields of economic evaluation of health care interventions.

ACKNOWLEDGEMENTS

Mr Andrea Gabrio is partially funded in his PhD programme at University College London by a research grant sponsored by The Foundation BLANCEFLOR Boncompagni Ludovisi, née Bildt. Dr Gianluca Baio is partially supported as the recipient of an unrestricted research grant sponsored by Mapi Group at University College London. Finally, we wish to thank Ms Julia V. Bailey, Ms Angela Hassiotis, and Ms Rachael Hunter at University College London for providing the MenSS and PBS trial data and advice on the original economic model.

ORCID

Andrea Gabrio  <https://orcid.org/0000-0002-7650-4534>

REFERENCES

1. NICE. Guide to the methods of technological appraisal. London, UK: National Institute for Health and Care Excellence; 2013.
2. Briggs AH. Handling uncertainty in cost-effectiveness models. *PharmacoEconomics*. 2000;17(5):479-500.
3. O'Hagan A, McCabe C, Akehurst R, et al. Incorporation of uncertainty in health economic modelling studies. *PharmacoEconomics*. 2004;23(6):529-536.
4. Sculpher MJ, Claxton K, Drummond M, McCabe C. Whither trial-based economic evaluation for health decision making? *Health Econ*. 2005;15(7):677-687.
5. Spiegelhalter DJ, Best NG. Bayesian approaches to multiple sources of evidence and uncertainty in complex cost-effectiveness modelling. *Statist Med*. 2003;22(23):3687-3709.
6. Claxton K. The irrelevance of inference: a decision making approach to stochastic evaluation of health care technologies. *J Health Econ*. 1999;18(3):341-364.
7. Baio G. *Bayesian Methods in Health Economics*. Boca Raton, FL: CRC Press; 2012.
8. Jackson CH, Thompson SG, Sharples LD. Accounting for uncertainty in health economic decision models by using model averaging. *J R Stat Soc A Stat Soc*. 2009;172(2):383-404.
9. Briggs A, Sculpher M, Claxton K. *Decision Modelling for Health Economic Evaluation*. Oxford, UK: Oxford University Press; 2006.
10. Spiegelhalter DJ, Abrams KR, Myles JP. *Bayesian Approaches to Clinical Trials and Health-Care Evaluation*. Chichester, UK: John Wiley & Sons; 2004.
11. Manca A, Hawkins N, Sculpher MJ. Estimating mean QALYs in trial-based cost-effectiveness analysis: the importance of controlling for baseline utility. *Health Econ*. 2005;14(5):487-496.
12. Hunter RM, Baio G, Butt T, Morris S, Round J, Freemantle N. An educational review of the statistical issues in analysing utility data for cost-utility analysis. *PharmacoEconomics*. 2015;33(4):355-366.

13. Committee for Medicinal Products for Human Use (CHMP). Guideline on adjustment for baseline covariates. London, UK: European Medicines Agency; 2013. http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2013/06/WC500144946.pdf
14. O'Hagan A, Stevens JW. A framework for cost-effectiveness analysis from clinical trial data. *Health Econ.* 2001;10(4):303-315.
15. Nixon RM, Thompson SG. Methods for incorporating covariate adjustment, subgroup analysis and between-centre differences into cost-effectiveness evaluations. *Health Econ.* 2005;14(12):1217-1229.
16. Thompson SG, Nixon RM. How sensitive are cost-effectiveness analyses to choice of parametric distributions? *Med Decis Making.* 2005;25(4):416-423.
17. Briggs A, Gray A. The distribution of health care costs and their statistical analysis for economic evaluation. *J Health Serv Res Policy.* 1998;3(4):233-245.
18. Rascati KL, Smith LJ, Neilands T. Dealing with skewed data: an example using asthma-related costs of medicaid clients. *Clin Ther.* 2001;23(3):481-498.
19. O'Hagan A, Stevens JW. Assessing and comparing costs: how robust are the bootstrap and methods based on asymptotic normality? *Health Econ.* 2003;12(1):33-49.
20. Basu A, Manca A. Regression estimators for generic health-related quality of life and quality-adjusted life years. *Med Decis Making.* 2012;32(1):56-69.
21. Cooper NJ, Sutton AJ, Mugford M, Abrams KR. Use of Bayesian Markov chain Monte Carlo methods to model cost-of-illness data. *Med Decis Making.* 2003;23(1):38-53.
22. Mihaylova B, Briggs A, O'Hagan A, Thompson SG. Review of statistical methods for analysing healthcare resources and costs. *Health Econ.* 2011;20(8):897-916.
23. Ntzoufras I. *Bayesian Modelling Using WinBUGS*. Hoboken, NJ: John Wiley & Sons; 2009.
24. Baio G. Bayesian models for cost-effectiveness analysis in the presence of structural zero costs. *Statist Med.* 2014;33(11):1900-1913.
25. Tooze JA, Grunwald GK, Jones RH. Analysis of repeated measures data with clumping at zero. *Statist Methods Med Res.* 2002;11(4):341-355.
26. Härkänen T, Maljanen T, Lindfors O, Virtala E, Knekt P. Confounding and missing data in cost-effectiveness analysis: comparing different methods. *Heal Econ Rev.* 2013;3(1):3-8.
27. Ramsey SD, Willke RJ, Glick H, et al. Cost-effectiveness analysis alongside clinical trials II-an ISPOR good research practices task force report. *Value Health.* 2015;18(2):161-172.
28. Groenwold RHH, Donders ART, Roes KCB, Harrell FE, Moons KGM. Dealing with missing outcome data in randomized trials and observational studies. *Am J Epidemiol.* 2012;175(3):210-217.
29. Wood AM, White IR, Thompson SG. Are missing outcome data adequately handled? A review of published randomized controlled trials in major medical journals. *Clin Trials.* 2004;1(4):368-376.
30. Noble SM, Hollingworth W, Tilling K. Missing data in trial-based cost-effectiveness analysis: the current state of play. *Health Econ.* 2012;21(2):187-200.
31. Gabrio A, Mason AJ, Baio G. Handling missing data in within-trial cost-effectiveness analysis: a review with future recommendations. *PharmacoEconomics-Open.* 2017;1(2):79-97.
32. Briggs A, Clark T, Wolstenholme J, Clarke P. Missing...presumed at random: cost-analysis of incomplete data. *Health Econ.* 2003;12(5):377-392.
33. Manca A, Palmer S. Handling missing data in patient-level cost-effectiveness analysis alongside randomised clinical trials. *Appl Health Econ Health Policy.* 2005;4(2):65-75.
34. Faria R, Gomes M, Epstein D, White IR. A guide to handling missing data in cost-effectiveness analysis conducted within randomised controlled trials. *PharmacoEconomics.* 2014;32(12):1157-1170.
35. Rubin DB. *Multiple Imputation for Nonresponse in Surveys*. Hoboken, NJ: John Wiley & Sons; 1987.
36. Díaz-Ordaz K, Kenward MG, Grieve R. Handling missing values in cost effectiveness analyses that use data from cluster randomized trials. *J R Stat Soc Ser A.* 2014;177(2):457-474.
37. Burton A, Billingham LJ, Bryan S. Cost-effectiveness in clinical trials: using multiple imputation to deal with incomplete cost data. *Clin Trials.* 2007;4(2):154-161.
38. Schafer JL. *Analysis of Incomplete Multivariate Data*. Boca Raton, FL: Chapman & Hall; 1997.
39. Schafer JL, Olsen MK. Multiple imputation for multivariate missing-data problems: a data analyst's perspective. *Multivar Behav Res.* 1998;33(4):545-571.
40. Van Buuren S, Groothuis-Oudshoorn K. MICE: multivariate imputation by chained equations in R. *J Stat Softw.* 2011;45(3):1-67.
41. Carpenter JR, Kenward MG, White IR. Sensitivity analysis after multiple imputation under missing at random: a weighting approach. *Stat Methods Med Res.* 2007;16(3):259-275.
42. Molenberghs G, Fitzmaurice G, Kenward MG, Tsiatis A, Verbeke G. *Handbook of Missing Data Methodology*. Boca Raton, FL: CRC Press; 2015.
43. Daniels MJ, Hogan JW. *Missing Data in Longitudinal Studies: Strategies for Bayesian Modeling and Sensitivity Analysis*. Boca Raton, FL: CRC Press; 2008.
44. Mason A, Richardson S, Plewis I, Best N. Strategy for modelling nonrandom missing data mechanisms in observational studies using Bayesian methods. *J Off Stat.* 2012;28(2):279-302.
45. Bailey JV, Webster R, Hunter R, et al. The men's safer sex project: intervention development and feasibility randomised controlled trial of an interactive digital intervention to increase condom use in men. *Health Technol Assess.* 2016;20(91):1-124.

46. Drummond MF, Schulpher MJ, Claxton K, Stoddart GL, Torrance GW. *Methods for the Economic Evaluation of Health Care Programmes*. 3rd ed. Oxford, UK: Oxford University Press; 2005.
47. Gelman A. Prior distributions for variance parameters in hierarchical models. *Bayesian Anal.* 2006;1(3):515-534.
48. Plummer M. JAGS: Just Another Gibbs Sampler. 2010. <http://mcmc-jags.sourceforge.net/>
49. Su Y-S, Yajima M. Package R2jags. 2015. <http://mcmc-jags.sourceforge.net/>
50. Gelman A, Carlin JB, Stern HS, Rubin DB. *Bayesian Data Analysis*. 2nd ed. Boca Raton, FL: Chapman and Hall/CRC; 2004.
51. Spiegelhalter DJ, Best NG, Carlin BP, van der Linde A. Bayesian measures of model complexity and fit. *J R Stat Soc.* 2002;64(4):583-639.
52. Celeux G, Forbes S, Robert CP, Titterton DM. Deviance information criteria for missing data models. *Bayesian Anal.* 2006;1(4):651-674.
53. Mason A, Richardson S, Best N. Two-pronged strategy for using DIC to compare selection models with non-ignorable missing responses. *Bayesian Anal.* 2012;7(1):109-146.
54. Black WC. The CE plane: a graphic representation of cost-effectiveness. *Med Decis Making.* 1990;10(3):212-214.
55. Van Hout BA, Al MJ, Gordon GS, Rutten FFH. Costs, effects and C/E-ratios alongside a clinical trial. *Health Econ.* 1994;3:309-319.
56. Hassiotis A, Poppe M, Strydom A, et al. Positive behaviour support training for staff for treating challenging behaviour in people with intellectual disabilities: a cluster RCT. *Health Technol Assess.* 2018;22(15):1-110.
57. Lambert PC, Billingham LJ, Cooper NJ, Sutton AJ, Abrams KR. Estimating the cost-effectiveness of an intervention in a clinical trial when partial cost information is available: a Bayesian approach. *Health Econ.* 2008;17(1):67-81.
58. Eekhout I. *Don't Miss Out!: Incomplete Data Can Contain Valuable Information* [PhD thesis]. Amsterdam, The Netherlands: EMGO+ Institute for Health and Care Research, Department of Epidemiology and Biostatistics, VU University Medical Center; 2014.
59. Simons CL, Rivero-Arias O, Yu L-M, Simon J. Multiple imputation to deal with missing EQ-5D-3L data: should we impute individual domains or the actual index? *Qual Life Res.* 2015;24(4):805-815.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Gabrio A, Mason AJ, Baio G. A full Bayesian model to handle structural ones and missingness in economic evaluations from individual-level data. *Statistics in Medicine*. 2018;1-22. <https://doi.org/10.1002/sim.8045>

APPENDIX A

MODEL CODE AND IMPLEMENTATION

A.1 | Implementation “trick”

The model described in Section 3.1.3 uses a different sampling distribution for the QALYs, depending on the observed value of the indicator d_{ie}

$$e_i | d_{ie} \sim \begin{cases} p(e_i | d_{ie} = 0) = p(e_i | \theta^{<1}), & \text{if } e_i < 1 \\ p(e_i | d_{ie} = 1) = p(e_i | \theta^1), & \text{if } e_i = 1, \end{cases}$$

where the model for $e_i = 1$ is degenerate at a point mass at 1, while that for $e_i < 1$ is defined in terms of a Beta distribution. We can conveniently rewrite this more succinctly and with specific reference to our case as

$$e_i \sim \text{Beta} \left(\phi_{ie}^{d_{ie}} \tau_{ie}^{d_{ie}}, \left(1 - \phi_{ie}^{d_{ie}} \right) \tau_{ie}^{d_{ie}} \right).$$

If we set $\phi_{ie}^1 = 1$ and select τ_{ie}^1 in order to induce a variance as close to 0 as possible, the two specifications are identical. Unfortunately, it is not possible to do so in the BUGS/JAGS language because the Beta distribution is specified in the open interval (0, 1), and thus, setting $\phi_{ie}^1 = 1$ implies that $\tau_{ie}^1 = 0$, which is not allowed.

However, the required behaviour is very closely mimicked if we define our model with

$$\text{logit}(\phi_{ie}^1) = \alpha_0^1 [+ \dots]$$




FIGURE A1 Sensitivity analysis for the choice of the standard deviation for the distribution of the structural ones in the quality-adjusted life years (QALYs). For each value of σ_e^1 tested, posterior means and 90% highest posterior density intervals for the mean QALYs parameters are respectively represented with dots and lines (red for the control and blue for the intervention group) [Colour figure can be viewed at wileyonlinelibrary.com]

and set $\alpha_0^1 = \text{logit}(0.999999)$ and $\sigma_e \approx 0$, which implies $\mu_e \approx 1$ with virtually no uncertainty. In other words, we can specify extremely informative priors on the parameters θ^1 so that the implied distribution for the structural ones components of the mixture is concentrated around 1 with essentially no uncertainty. More importantly, with such a prior, no amount of data can modify the posterior. The critical aspect of this strategy, however, is that inferences may be potentially sensitive to the way such priors are specified, that is, whether a small variation in the hyperprior values can affect the posterior estimates.

In fact, the estimation of the other parameters is not really affected by this choice, provided that the encoded prior really induces the variance towards zero. It is also plausible that different values for σ_e^1 have an impact on measures of model fit, such as the DIC. This is essentially due to the fact that the population is really comprised of two groups, one of which shows QALYs that are identically one. Thus, the closer the approximation to zero for the variance, the better the fit to the observed data and, therefore, the smaller the resulting DIC.

With this in mind, we have used different values for σ_e^1 to assess the impact on the mean QALYs estimates. Fixing the value of the mean for the ones group to $\mu_e^1 = 0.999999$ corresponds to an upper bound for the standard deviation of 0.0001 (see Section 3.1.2). We have explored a range of possibilities by progressively decreasing this value and assessed their impact on posterior results.

Figure A1 shows the sensitivity of the inferences across the alternative specifications for σ_e^1 . Results in terms of mean posterior estimates and 90% HPD intervals were almost unchanged in all the cases. Thus, we can assert that model performance was unaffected by the choice of the value for σ_e^1 . We also observe that the DIC becomes smaller when the standard deviation parameter decreases and the best-fitting model is the one associated with the smallest values, although the results are hardly different from both an estimation and convergence perspective for all the parameters.

A.2 | Code

The complete JAGS code for the hurdle model used in the analysis is given below.

```
model {
  # data variables
  # e, c and u denote the QALYs, costs and baseline utilities
```



```

# d.e and d.u denote the structural one indicators for e and u
# age, ethnicity and employment are covariates in the model of d.e and d.u

# control group (t = 1)

for(i in 1 : N1) {

  # 1. Module for the structural ones in the QALYs
  d.e1[i] ~ dbern(pi.e[i, 1])
  logit(pi.e[i, 1]) <- gamma0[1] + gamma1[1] * (u1[i] - mean(u1[])) +
    gamma2[1] * (age1[i] - mean(age1[])) + gamma3[ethnicity1[i], 1] + gamma4[employment1[i], 1]

  #2. Module for the structural ones in the baseline utilities
  d.u1[i] ~ dbern(pi.u[i, 1])
  logit(pi.u[i, 1]) <- eta0[1] + eta1[1] * (age1[i] - mean(age1[])) + eta2[ethnicity1[i], 1] + eta3[employment1[i], 1]

  #3. Marginal module for the QALYs
  e1[i] ~ dbeta(phi.e[i, 1] * tau.e[i, 1], (1 - phi.e[i, 1]) * tau.e[i, 1])
  tau.e[i, 1] <- phi.e[i, 1] * (1 - phi.e[i, 1]) / pow(sigma.e[d.e1[i] + 1], 2) - 1
  logit(phi.e[i, 1]) <- alpha0[d.e1[i]+1, 1] + alpha1[d.e1[i]+1, 1] * (u1[i] - mean(u1[]))

  #4. Marginal module for the baseline utilities
  u1[i] ~ dbeta(mu.u[d.u1[i] + 1, 1] * tau.u[d.u1[i] + 1, 1], (1 - mu.u[d.u1[i] + 1, 1]) * tau.u[d.u1[i] + 1, 1])

  #5. Conditional module for the costs
  c1[i] ~ dgamma(phi.c[i, 1] * tau.c[i, 1], tau.c[i, 1])
  tau.c[i, 1] <- phi.c[i, 1] / pow(sigma.c[1], 2)
  log( phi.c[i, 1]) <- beta0[1] + beta1[1] * (e1[i] - mu.e[1])
}

#intervention group (t = 2)

for(i in 1 : N2) {

  #1. Module for the structural ones in the QALYs
  d.e2[i] ~ dbern(pi.e[i, 2])
  logit(pi.e[i, 2]) <- gamma0[2] + gamma1[2] * (u2[i] - mean(u2[])) +
    gamma2[2] * (age2[i] - mean(age2[])) + gamma3[ethnicity2[i], 2] + gamma4[employment2[i], 2]

  #2. Module for the structural ones in the baseline utilities
  d.u2[i] ~ dbern(pi.u[i, 2])
  logit(pi.u[i, 2]) <- eta0[2] + eta1[2] * (age2[i] - mean(age2[])) + eta2[ethnicity2[i], 2] + eta3[employment2[i], 2]

  #3. Marginal module for the QALYs
  e2[i] ~ dbeta(phi.e[i, 2] * tau.e[i, 2], (1 - phi.e[i, 2]) * tau.e[i, 2])
  tau.e[i, 2] <- phi.e[i, 2] * (1 - phi.e[i, 2]) / pow(sigma.e[d.e2[i] + 1], 2) - 1
  logit(phi.e[i, 2]) <- alpha0[d.e2[i] + 1, 2] + alpha1[d.e2[i] + 1, 2] * (u2[i] - mean(u2[]))

  #4. Marginal module for the baseline utilities
  u2[i] ~ dbeta(mu.u[d.u2[i] + 1, 2] * tau.u[d.u2[i] + 1, 2], (1 - mu.u[d.u2[i] + 1, 2]) * tau.u[d.u2[i] + 1, 2])

  #5. Conditional module for the costs
  c2[i] ~ dgamma(phi.c[i, 2] * tau.c[i, 2], tau.c[i, 2])
  tau.c[i, 2] <- phi.c[i, 2] / pow(sigma.c[2], 2)
  log( phi.c[i, 2]) <- beta0[2] + beta1[2] * (e2[i] - mu.e[2])
}

#Priors
#priors for module 1 and 2

for(t in 1 : 2) {
  gamma0[t] ~ dlogis(0, 1)
  gamma1[t] ~ dnorm(0, 0.00001)
  gamma2[t] ~ dnorm(0, 0.00001)

  eta0[t] ~ dlogis(0, 1)
  eta2[t] ~ dnorm(0, 0.00001)

  #priors on coefficients for categorical covariates
  #(setting reference category as 0)
  gamma3[1, t] <- 0
  gamma4[1, t] <- 0

  eta2[1, t] <- 0

```

```

    eta3[1, t] <- 0
  }

# set priors for all other categories
# use blocking to improve model convergence
# mu and tau values provided as data variables with zero means and small precisions (0.00001)
# ethnicity has different numbers of categories between arms

gamma3[2:14, 1] ~ dnmnorm(mu1.gamma3[], tau1.gamma3[, ])
gamma3[2:12, 2] ~ dnmnorm(mu2.gamma3[], tau2.gamma3[, ])
gamma4[2:6, 1] ~ dnmnorm(mu1.gamma4[], tau1.gamma4[, ])
gamma4[2:6, 2] ~ dnmnorm(mu2.gamma4[], tau2.gamma4[, ])

eta2[2:14, 1] ~ dnmnorm(mu1.eta2[], tau1.eta2[, ])
eta2[2:12, 2] ~ dnmnorm(mu2.eta2[], tau2.eta2[, ])
eta3[2:6, 1] ~ dnmnorm(mu1.eta3[], tau1.eta3[, ])
eta3[2:6, 2] ~ dnmnorm(mu2.eta3[], tau2.eta3[, ])

for(t in 1 : 2) {
  # priors for model 3
  # priors for the ones group in the QALYs
  alpha0[2, t] <- logit(0.999999)
  alpha1[2, t] <- 0
  sigma.e[2, t] <- 0.00001
  # priors for the non-ones group in the QALYs
  alpha0[1, t] ~ dnorm(0, 0.000001)
  alpha1[1, t] ~ dnorm(0, 0.000001)
  sigma.e[1, t] ~ dunif(0, sd.limit.e[t])
  sd.limit.e[t] <- pow(mu.e[1, t] * (1 - mu.e[1, t]), 0.5)

  # priors for model 4
  # priors for the ones group in the baseline utilities
  tau.u[2, t] <- mu.u[2, t] * (1 - mu.u[2, t]) / pow(sigma.u[2, t], 2) - 1
  logit(mu.u[2, t]) <- delta0[2, t]
  delta0[2, t] <- logit(0.999999)
  sigma.u[2, t] <- 0.00001
  # priors for the non-ones group in the baseline utilities
  tau.u[1, t] <- mu.u[1, t] * (1 - mu.u[1, t]) / pow(sigma.u[1, t], 2) - 1
  logit(mu.u[1, t]) <- delta0[1, t]
  delta0[1, t] ~ dnorm(0, 0.00001)
  sigma.u[1, t] ~ dunif(0, sd.limit.u[t])
  sd.limit.u[t] <- pow(mu.u[1, t] * (1 - mu.u[1, t]), 0.5)

  # priors for module 5
  beta0[t] ~ dnorm(0, 0.00001)
  sigma.c[t] ~ dunif(0, 1000)
  beta1[t] ~ dnorm(0, 0.00001)

  # obtain marginal probabilities for weighting
  p[t] <- ilogit(gamma0[t])

  # obtain the weighted marginal mean QALYs
  mu.e[t] <- p[t] + (1-p[t]) * ilogit(alpha0[t])
}

# compute incremental QALYs and costs
Delta_e <- mu.e[2] - mu.e[1]
Delta_c <- mu.c[2] - mu.c[1]
}

```