

WHITMORE, Harold James

DOB: March 8, 1945 | Age: 69 yrs | Sex: Male | MRN: SMC-04471932

Encounter Date: September 25, 2014 | Provider: Sandra Liu, MD — Medical Oncology | Doc ID: NCI-2014-0925-PSA

Oncology Surveillance Visit

Post-Radiotherapy PSA Surveillance — Approaching Nadir

Subjective

69-year-old, 8 months post completion of prostate IMRT, continuing leuprolide ADT. Reports hot flashes, mild fatigue, decreased libido (expected ADT effects). No bone pain, hematuria, or obstructive symptoms. Mood stable.

Vital Signs

BP (mmHg)	HR (bpm)	RR	Temp	SpO2	Weight	BMI
126/74	64	15	36.6 C	98% RA	92.0 kg	30.0

Surveillance Labs

Test	Result	Units	Reference Range	Flag
PSA, total	0.2	ng/mL	Post-treatment trend	
Testosterone, total	11	ng/dL	< 20 on ADT (castrate)	
Hemoglobin A1c	7.6	%	< 7.5 (individualized)	H
Creatinine	1.1	mg/dL	0.7 - 1.3	

Assessment

- Prostate cancer s/p definitive IMRT + ADT — excellent biochemical response; PSA approaching nadir at 0.2 ng/mL with castrate testosterone.
- ADT-related vasomotor symptoms — tolerable.
- ADT-associated worsening of glycemia (A1c drift to 7.6%) — refer endocrinology to intensify.
- Bone health on ADT — DEXA up to date.

Plan

- Continue leuprolide per 18-month plan (final dose anticipated spring 2015).
- PSA + testosterone every 3 months.
- Endocrinology to adjust diabetes regimen (see endocrine note).
- Continue calcium/vitamin D; encourage weight-bearing exercise.
- Manage hot flashes conservatively.