

WHITMORE, Harold James

DOB: March 8, 1945 | Age: 75 yrs | Sex: Male | MRN: SMC-04471932
Encounter Date: September 15, 2020 | Provider: David Mercer, MD — Orthopaedic Surgery | Doc ID: SMC-2020-0915-OP

Operative Report
Right Total Knee Arthroplasty

Operative Details

Surgeon	David Mercer, MD — Orthopaedic Surgery
Pre-op Diagnosis	Severe right knee osteoarthritis (KL grade 4), failed conservative management
Procedure	Right total knee arthroplasty (cemented, cruciate-retaining)
Anesthesia	Spinal with adductor canal block
Estimated blood loss	120 mL · Tourniquet time 58 min

Indication & Pre-op Optimization

75-year-old with end-stage right knee OA, progressive pain limiting mobility despite NSAIDs (limited by CKD), physiotherapy, and intra-articular corticosteroid injections (2018, 2019). Cardiology cleared for surgery; HF compensated (recent EF ~38%). Apixaban held 48 h pre-op; bridging not required. Tobacco-free.

Procedure Summary

Standard medial parapatellar approach. Femoral and tibial cuts made with measured resection. Trial reduction with good tracking and stability. Components cemented (femoral, tibial baseplate with polyethylene insert, patellar resurfacing). Tourniquet released, hemostasis confirmed. Wound closed in layers. Tolerated procedure well.

Post-op Plan

- VTE prophylaxis: resume apixaban 24 h post-op (held intra-op) — therapeutic dose covers both AF and prophylaxis; monitor for bleeding.
- Multimodal analgesia avoiding nephrotoxic NSAIDs: acetaminophen 1 g q8h scheduled, low-dose oxycodone PRN, periarticular block.
- Early mobilization with physiotherapy day 0-1; weight-bearing as tolerated.
- Continue home cardiac/HF/diabetes medications; monitor fluid status (HF) and renal function.
- Anticipated discharge day 2-3 with home PT.

Peri-operative Medications

Medication	Dose	Route	Frequency	Indication
Acetaminophen	1000 mg	PO	Every 8 h scheduled	Analgesia
Oxycodone	5 mg	PO	Every 4-6 h PRN	Breakthrough pain
Apixaban	5 mg	PO	Twice daily (resume 24h post-op)	AF / VTE prophylaxis
Cefazolin	2 g	IV	Pre-op + 2 doses	Surgical prophylaxis