

WHITMORE, Harold James

DOB: March 8, 1945 | Age: 68 yrs | Sex: Male | MRN: SMC-04471932
Encounter Date: September 2, 2013 | Provider: Marcus Webb, MD — Urology | Doc ID: URO-2013-0902-BX

Urology — Prostate Biopsy Procedure & Clinic Note

Transrectal Ultrasound-Guided 12-Core Prostate Biopsy

History

68-year-old referred for PSA 7.8 ng/mL (07/08/2013). Confirmatory PSA on 08/12/2013 was 8.1 ng/mL, free PSA 9% (low, raising malignancy suspicion). DRE: smooth enlargement without nodule. Counseled on biopsy risks (bleeding, infection, urinary retention); consent obtained.

Procedure

Transrectal ultrasound: prostate volume 38 cc; no discrete hypoechoic lesion. Periprostatic nerve block with lidocaine 1% (10 mL). Twelve-core systematic biopsy obtained (6 per lobe). Patient tolerated well; minimal bleeding. Prophylaxis: ciprofloxacin 500 mg PO (pre- and post-procedure x3 days).

Immediate Course

No retention; voided prior to discharge. Advised re hematuria/hemospermia expected for days to weeks. Pathology pending — see Surgical Pathology report (09/09/2013).

Plan

- Await pathology; office follow-up to discuss results in 1-2 weeks.
- Continue tamsulosin 0.4 mg nightly.
- Return precautions for fever, inability to void, or heavy bleeding.

Medications This Encounter

Medication	Dose	Route	Frequency	Indication
Ciprofloxacin	500 mg	PO	BID x3 days	Biopsy prophylaxis
Lidocaine 1%	10 mL	Local	Once	Periprostatic block
Tamsulosin	0.4 mg	PO	Nightly	LUTS / BPH