

# Tutorial: Reminder (?) on mixture model and the EM algorithm - correction

MSc in Statistics for Smart Data – Introduction to graph analysis and modeling

Julien Chiquet, November the 13th, 2018

## 1 Gaussian Mixture Models

We consider a collection of random variables  $(X_1, \dots, X_n)$  associated with  $n$  individuals drawn from  $Q$  populations. The label of each individual describes the population (or class) to which it belongs and is unobserved. The  $Q$  classes have *a priori* distribution  $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_Q)$  with  $\alpha_q = \mathbb{P}(i \in q)$ . The hidden random indicator variables  $(Z_{iq})_{i \in \mathcal{P}, q \in \mathcal{Q}}$  describe the label of each individuals, that is,

$$\alpha_q = \mathbb{P}(Z_{iq} = 1) = \mathbb{P}(i \in q), \quad \text{such that } \sum_{q=1}^Q \alpha_q = 1.$$

Remark that we have  $\mathbf{Z}_i = (Z_{i1}, \dots, Z_{iQ}) \sim \mathcal{M}(1, \boldsymbol{\alpha})$ . The distribution of  $X_i$  conditional on the label of  $i$  is assumed to be a univariate gaussian distribution with unknown parameters, that is,  $X_i | Z_{iq} = 1 \sim \mathcal{N}(\mu_q, \sigma_q^2)$ .

## 2 Questions

2.1 *Likelihood.* Write the model complete-data loglikelihood.

We denote the vector of parameters to be estimated by  $\boldsymbol{\mu} = (\mu_1, \dots, \mu_Q)$ ,  $\boldsymbol{\sigma}^2 = (\sigma_1^2, \dots, \sigma_Q^2)$ ,  $\boldsymbol{\tau} = (\tau_{iq, i=1, \dots, n; q=1, \dots, Q})$ . The negative complete-data loglikelihood is derived as follows

$$\begin{aligned} \log L(\boldsymbol{\mu}, \boldsymbol{\sigma}^2, \boldsymbol{\tau}; \mathbf{x}) &= \sum_{i=1}^n \log \left( \prod_{q=1}^Q \alpha_q f(x_i; \mu_q, \sigma_q^2)^{Z_{iq}} \right) \\ &= \sum_{i=1}^n \sum_{q=1}^Q Z_{iq} \left( \log \frac{\alpha_q}{\sigma \sqrt{2\mathbb{P}i}} \exp \left\{ \frac{1}{2\sigma^2} (x_i - \mu_q)^2 \right\} \right) \\ &= \sum_{i=1}^n \sum_{q=1}^Q Z_{iq} \left( \log \alpha_q - \log \sigma - \log(\sqrt{2\mathbb{P}i}) - \frac{1}{2\sigma_q^2} (x_i - \mu_q)^2 \right) \end{aligned} \tag{1}$$

## 2.2 E-step.

For fixed values of  $\hat{\mu}_q, \hat{\sigma}_q^2$  and  $\hat{\alpha}_q$ , give the expression of the estimates of the posterior probabilities  $\tau_{iq} = \mathbb{P}(Z_{iq} = 1|X_i)$ .

$$\tau_{iq} = \frac{\hat{\alpha}_q f(x_i; \hat{\mu}_q, \hat{\sigma}_q^2)}{\sum_{q=1}^Q \hat{\alpha}_q f(x_i; \hat{\mu}_q, \hat{\sigma}_q^2)}, \quad (2)$$

where  $f$  is the density of the normal distribution.

## 2.3 M-step.

The maximization step consists in solving the following optimization problem

$$\arg \max_{\sigma_q, \mu_q, \alpha_q} \underbrace{\sum_{i=1}^n \sum_{q=1}^Q \hat{\tau}_{iq} \left( \log \alpha_q - \log \sigma - \log(\sqrt{2\mathbb{P}i}) - \frac{1}{2\sigma_q^2} (x_i - \mu_q)^2 \right)}_{Q(\sigma, \mu, \alpha; \hat{\tau})} \quad (3)$$

Consider first the mixture coefficients. We solve the above maximization problem under the constraint that the mixture coefficients sum to 1. This can be dealt with the Lagrange multiplier technique. By deriving the objective function w.r.t  $\alpha_q$ , we get

$$\frac{\sum_i \tau_{iq}}{\alpha_q} + \lambda = 0 \Leftrightarrow \alpha_q = \frac{\sum_i \tau_{iq}}{-\lambda} \quad (4)$$

where  $\lambda$  corresponds to the Lagrange multiplier associated with the constraint  $\sum_q \alpha_q = 1$ . If we sum the latter result over all  $q$ , we get that  $1 = \sum_q \tau_{iq}/(-\lambda)$ . In other words,  $\lambda = -\sum_q \tau_{iq}$  so that finally

$$\hat{\alpha}_q = \frac{\sum_{i=1}^n \tau_{iq}}{\sum_{i=1}^n \sum_{q=1}^Q \tau_{iq}} \quad (5)$$

Concerning,  $\hat{\mu}_q$ , null gradient condition leads to

$$\sum_i \frac{\tau_{iq}}{2\sigma_q^2} (x_i - \mu_q) = 0 \Leftrightarrow \mu_q = \frac{\sum_i \tau_{iq} x_i}{\sum_i \tau_{iq}} \quad (6)$$

Similarly, for  $\hat{\sigma}_q$ , we get

$$\sum_{i=1}^n \tau_{iq} \left( -\frac{1}{2\sigma_q^2} + \frac{1}{2\sigma_q^4} (x_i - \mu_q)^2 \right) = 0 \Leftrightarrow \sigma_q^2 = \frac{\sum_{i=1}^n \tau_{iq} (x_i - \mu_q)^2}{\sum_{i=1}^n \tau_{iq}} \quad (7)$$

- *Implementation.*

```

get.cloglik <- function(X, Z, alpha, mu, sigma) {
  n <- length(X); Q <- length(alpha)
  xs <- sweep(sweep(matrix(X,n,Q),2,mu,"-"),2,sigma,"/")
  return(sum(t(Z)*(log(alpha)-log (sigma)-.5*(log(2*pi)+t(xs)^2))))
}

EM.mixture <- function(X, Q,
                       init.cl=kmeans(X,Q)$cl, max.iter=100, eps=1e-5) {
  ## INITIALIZATIONS
  n <- length(X)
  tau <- matrix(0,n,Q)
  tau[cbind(1:n,init.cl)] <- 1
  loglik <- vector("numeric", max.iter)
  cloglik <- vector("numeric", max.iter)
  iter <- 0; cond <- FALSE

  while (!cond) {
    iter <- iter + 1
    ## M step
    alpha <- colMeans(tau)
    mu <- colMeans(tau * matrix(X,n,Q)) / alpha
    sigma <- sqrt(colMeans(tau*sweep(matrix(X,n,Q),2,mu,"-")^2)/alpha)
    ## E step
    prob <- sapply(1:Q, function(q) alpha[q]*dnorm(x,mu[q],sigma[q]))
    likelihoods <- rowSums(prob)
    tau <- prob / likelihoods
    Z <- 1*(tau >= .5)
    ## check consistency
    loglik[iter] <- sum(log(likelihoods))
    cloglik[iter] <- get.cloglik(X, Z, alpha, mu, sigma)
    if (iter > 1)
      cond <- (iter>=max.iter) | abs(cloglik[iter]-cloglik[iter-1]) < eps
  }

  return(list(alpha = alpha ,
             mu = mu ,
             sigma = sigma,
             tau = tau ,
             cl = apply(Z,1,which.max),
             cloglik = cloglik[1:iter],
             loglik = loglik[1:iter]))
}

```

## 2.4 Examples

We test ICL and BIC on a simple example with 4 groups


Let us start with the data generation.

```
mu1 <- 5 ; sigma1 <- 1; n1 <- 100
mu2 <- 10 ; sigma2 <- 1; n2 <- 200
mu3 <- 15 ; sigma3 <- 2; n3 <- 50
mu4 <- 20 ; sigma4 <- 3; n4 <- 100
c1 <- rep(1:4,c(n1,n2,n3,n4))
x <- c(rnorm(n1,mu1,sigma1),rnorm(n2,mu2,sigma2),
      rnorm(n3,mu3,sigma3),rnorm(n4,mu4,sigma4))
n <- length(x)

## we randomize the class ordering
rnd <- sample(1:n)
c1 <- c1[rnd]
x <- x[rnd]


alpha <- c(n1,n2,n3,n4)/n

curve(alpha[1]*dnorm(x,mu1,sigma1) +
      alpha[2]*dnorm(x,mu2,sigma2) +
      alpha[3]*dnorm(x,mu3,sigma3) +
      alpha[4]*dnorm(x,mu4,sigma4),
      col="blue", lty=1, from=0,to=30, n=1000,
      main="Theoretical Gaussian mixture and its components",
      xlab="x", ylab="density")
curve(alpha[1]*dnorm(x,mu1,sigma1), col="red", add=TRUE, lty=2)
curve(alpha[2]*dnorm(x,mu2,sigma2), col="red", add=TRUE, lty=2)
curve(alpha[3]*dnorm(x,mu3,sigma3), col="red", add=TRUE, lty=2)
curve(alpha[4]*dnorm(x,mu4,sigma4), col="red", add=TRUE, lty=2)
rug(x)
```




Suppose that we know the number of components, i.e. 4.

```
out <- EM.mixture(x, Q=4)
plot(out$loglik, main="data log-likelihood", type="l")
```



```
curve(alpha[1]*dnorm(x,mu1,sigma1) +
      alpha[2]*dnorm(x,mu2,sigma2) +
      alpha[3]*dnorm(x,mu3,sigma3) +
      alpha[4]*dnorm(x,mu4,sigma3), col="blue",
      lty=1, from=0,to=30, n=1000,
      main="Theoretical Gaussian mixture and estimated components",
      xlab="x", ylab="density")
curve(out$alpha[1]*dnorm(x,out$mu[1],out$sigma[1]), col="red", add=TRUE, lty=2)
curve(out$alpha[2]*dnorm(x,out$mu[2],out$sigma[2]), col="red", add=TRUE, lty=2)
curve(out$alpha[3]*dnorm(x,out$mu[3],out$sigma[3]), col="red", add=TRUE, lty=2)
curve(out$alpha[4]*dnorm(x,out$mu[4],out$sigma[4]), col="red", add=TRUE, lty=2)
rug(x)
```




```
## the confusion table gives rather good results
table(out$c1,c1)
```

```
##      c1
##      1  2  3  4
##  1  0 172  1  1
##  2  91  0  0  0
##  3   9  28 45 25
##  4   0   0  4 74
```

The number of component mixture is hard to recover because of the last two mixed components

```
seq.Q <- 2:10
crit.EM <- sapply(seq.Q, function(Q) {
  out <- EM.mixture(x, Q)
  df <- Q-1 + 2 * Q
  return(c(BIC = -2*tail(out$loglik,1) + log(n)*df,
           ICL = -2*tail(out$cloglik,1) + log(n)*df ))
})
matplot(seq.Q, t(crit.EM), type="l", col=c("red", "blue"))
legend("topleft", c("BIC", "ICL"), col=c("red", "blue"),lty=1)
```



```
Q.hat <- seq.Q[which.min(crit.EM[1, ])]
out <- EM.mixture(x, Q=Q.hat)
par(mfrow=c(1,1))
curve(alpha[1]*dnorm(x,mu1,sigma1) +
      alpha[2]*dnorm(x,mu2,sigma2) +
      alpha[3]*dnorm(x,mu3,sigma3) +
      alpha[4]*dnorm(x,mu4,sigma3), col="blue",
      lty=1, from=0,to=30, n=1000,
      main="Theoretical Gaussian mixture and estimated components",
      xlab="x", ylab="density")
for (q in 1:Q.hat) {
  curve(out$alpha[q]*dnorm(x,out$mu[q],out$sigma[q]), col="red", add=TRUE, lty=2)
```

```
}  
rug(x)
```

**Theoretical Gaussian mixture and estimated components**

