

Termination in Convex Sets of Distributions

Ana Sokolova and Harald Woracek



CALCO 2017, Ljubljana, 14.6.2017

Termination in Convex Sets of Distributions

Ana Sokolova and Harald Woracek



CALCO 2017, Ljubljana, 14.6.2017

Convex Algebras

infinitely many
finitary operations

convex
combinations

binary ones
“suffice”

- algebras

$$(A, \sum_{i=1}^n p_i (-)_i)$$

$$p_i \in [0, 1], \sum_{i=1}^n p_i = 1$$

- convex (affine) maps

$$h \left(\sum_{i=1}^n p_i a_i \right) = \sum_{i=1}^n p_i h(a_i)$$

satisfying

- Projection

$$\sum_{i=1}^n p_i a_i = a_k, \quad p_k = 1$$

- Barycenter

$$\sum_{i=1}^n p_i \left(\sum_{j=1}^m p_{i,j} a_j \right) = \sum_{j=1}^m \left(\sum_{i=1}^n p_i p_{i,j} \right) a_j$$

Eilenberg-Moore Algebras

convex algebras
abstractly

$\mathcal{EM}(\mathcal{D})$

- objects

$$\begin{array}{c} \mathcal{D}A \\ \downarrow a \\ A \end{array}$$

satisfying

$$\begin{array}{ccc} A & \xrightarrow{\eta} & \mathcal{D}A \\ & \searrow a & \downarrow a \\ & & A \end{array}$$

$$\begin{array}{ccc} \mathcal{D}\mathcal{D}A & \xrightarrow{\mu} & \mathcal{D}A \\ \mathcal{D}a \downarrow & & \downarrow a \\ \mathcal{D}A & \xrightarrow{a} & A \end{array}$$

- morphisms

$$\begin{array}{c} \mathcal{D}A \\ \downarrow a \\ A \end{array} \xrightarrow{h} \begin{array}{c} \mathcal{D}B \\ \downarrow b \\ B \end{array}$$


$$\begin{array}{ccc} \mathcal{D}A & \xrightarrow{\mathcal{D}h} & \mathcal{D}B \\ a \downarrow & & \downarrow b \\ A & \xrightarrow{h} & B \end{array}$$

Probabilistic Automata

without termination

belief-state
transformers

coalgebras on
 $\mathcal{EM}(\mathcal{D})$



possible behaviour

$$\begin{array}{c} \frac{1}{3}x_1 + \frac{2}{3}x_2 \\ \downarrow a \\ \frac{2}{3}x_2 + \frac{1}{6}x_3 + \frac{1}{6}x_4 \end{array}$$

$$\begin{array}{c} \frac{1}{3}x_1 + \frac{2}{3}x_2 \\ \downarrow a \\ \frac{8}{9}x_2 + \frac{1}{9}x_3 \end{array}$$

termination?

$$\begin{array}{c} \frac{1}{3}x_1 + \frac{2}{3}x_3 \\ \downarrow a \\ ? \end{array}$$

Probabilistic Automata

without termination

belief-state
transformers

coalgebras on
 $\mathcal{EM}(\mathcal{D})$
with carriers
free algebras

$\mathcal{D}\mathcal{D}S$
 $\downarrow \mu$
 $\mathcal{D}S$

free
algebra

\mathcal{P}_c

nonempty convex
powerset

$\mathcal{D}\mathcal{D}S$
 $\downarrow \mu$
 $\mathcal{D}S$


L

constant exponent

[Bonchi Silva S. '17]

Probabilistic Automata

termination?




Markov Chains

no termination

belief-state
transformers

coalgebras on
 $\mathcal{EM}(\mathcal{D})$



possible behaviour

$$\begin{array}{c} \frac{1}{2}x_1 + \frac{1}{2}x_2 \\ \downarrow \\ \frac{2}{3}x_2 + \frac{1}{12}x_3 + \frac{1}{4}x_4 \end{array}$$

Markov Chains

no termination

belief-state
transformers

coalgebras on
 $\mathcal{EM}(\mathcal{D})$
with carriers
free algebras

$\mathcal{D}\mathcal{D}S$
 $\downarrow \mu$
 $\mathcal{D}S$

free
algebra


$\mathcal{D}\mathcal{D}S$
 $\downarrow \mu$
 $\mathcal{D}S$

for the
identity functor

termination?

Markov Chains

termination?



The Problem

- Given a convex algebra, is it possible to extend it by a single element?

YES!

- If yes, what are all the possible ways?

there are many possible ways

we can give full description for...

single naturally functorial way

The Cases of Interest[★]

we can give full description for:

- Free convex algebras

$$\mathbb{D}_S = \begin{array}{c} \mathcal{D}\mathcal{D}S \\ \downarrow \mu \\ \mathcal{D}S \end{array}$$

- Convex subsets of convex subsets[★] of a vector space

in particular

$$\mathcal{P}_c \mathbb{D}_S = \mathcal{P}_c \begin{array}{c} \mathcal{D}\mathcal{D}S \\ \downarrow \mu \\ \mathcal{D}S \end{array}$$

nonempty

$\mathcal{P}_c \mathbb{D}$

convex subset
of a vector
space

$$X = \mathbb{D}_S$$

Free convex algebras

carrier

$$X_* = X + 1 = X \cup \{*\}$$

Possible extensions X_* are:

- the black-hole extension

$$px + (1 - p)* = *$$

- $*$ imitates a point $w \in X$

$$px + (1 - p)* = px + (1 - p)w$$

- $*$ imitates one of the extremal points $s \in S$ on all other points, and **adheres** this point

$$px + (1 - p)* = px + (1 - p)s, \quad x \neq s$$

$$ps + (1 - p)* = *$$

these are all extensions!

Functoriality

Given a functor $F: \mathcal{EM}(\mathcal{D}) \rightarrow \mathcal{EM}(\mathcal{D})$ with

- $\mathbb{X} \leq F\mathbb{X}$
- $F\mathbb{X}$ has carrier $X + 1 = X \cup \{*\}$
- $$\begin{array}{ccc} F\mathbb{X} & \xrightarrow{Ff} & F\mathbb{Y} \\ \iota_X \uparrow & & \uparrow \iota_Y \\ \mathbb{X} & \xrightarrow{f} & \mathbb{Y} \end{array}$$

unique / single
functorial
extension

Then $F\mathbb{X}$ must be
the black-hole extension !

$$\mathbb{X} = \mathcal{P}_c \mathbb{D}$$

Convex subsets of...

Possible extensions \mathbb{X}_* are:

- the black-hole extension
- * imitates a “point” $C \in \mathcal{P}_c(\text{Vis}(D))$
- * imitates one of the extremal “points” of \mathbb{D} on all other points, and **adheres** this point
- * imitates $C \in \mathcal{P}_c(\text{Vis}(D))$ on P and **adheres** $X \setminus P$

visibility hull

these are all extensions if $D - D$ is linearly bounded!

$$|C| \geq 2$$

$$\text{conv}\{A \in X \mid A \not\bullet C\} \subseteq P \neq X$$

prime ideal

$$\mathbb{X} = \mathcal{P}_c \mathbb{D}$$

$$D \subseteq V$$





vector
space

Visibility hull

$$\text{Vis}(D) = \{v \in V \mid \forall d \in D. \forall p \in (0, 1). pv + (1 - p)d \in D\}$$

all points
from
which all of D
is visible

$V \setminus D$ is not
transparent



$$\mathbb{X} = \mathcal{P}_c \mathbb{D}$$

Convex subsets of...

Possible extensions \mathbb{X}_* are:

- the black-hole extension
- * imitates a “point” $C \in \mathcal{P}_c(\text{Vis}(D))$
- * imitates one of the extremal “points” of \mathbb{D} on all other points, and **adheres** this point
- * imitates $C \in \mathcal{P}_c(\text{Vis}(D))$ on P and **adheres** $X \setminus P$

visibility hull

these are all extensions if $D - D$ is linearly bounded!

$|C| \geq 2$

$\text{conv}\{A \in X \mid A \not\prec C\} \subseteq P \neq X$

prime ideal

somewhat implicit description

Summing-up

Thank You!

- Convex algebras are important for the semantics of probabilistic systems
- We looked at one-point extensions, for termination.

Every convex algebra can be extended by a single point

- What are all the possible ways?

there are many possible ways

we can give full description for...

single naturally functorial way