

# Convex Algebras for Probabilistic Systems

Ana Sokolova



TRENDS '17

# Convex Algebras

- algebras
- convex (affine) maps

# Convex Algebras

- algebras

$$(A, \sum_{i=1}^n p_i(-)_i)$$

- convex (affine) maps

# Convex Algebras

- algebras

$$(A, \sum_{i=1}^n p_i (-)_i)$$

$$p_i \in [0, 1], \sum_{i=1}^n p_i = 1$$

- convex (affine) maps

# Convex Algebras

infinitely many  
finitary operations

- algebras

$$(A, \sum_{i=1}^n p_i (-)_i)$$

$$p_i \in [0, 1], \sum_{i=1}^n p_i = 1$$

- convex (affine) maps

# Convex Algebras

infinitely many  
finitary operations

convex  
combinations

- algebras

$$(A, \sum_{i=1}^n p_i (-)_i)$$

$$p_i \in [0, 1], \sum_{i=1}^n p_i = 1$$

- convex (affine) maps

# Convex Algebras

infinitely many  
finitary operations

convex  
combinations

binary ones  
“suffice”

- algebras

$$(A, \sum_{i=1}^n p_i (-)_i)$$

$$p_i \in [0, 1], \sum_{i=1}^n p_i = 1$$

- convex (affine) maps

# Convex Algebras

infinitely many  
finitary operations

convex  
combinations

binary ones  
“suffice”

- algebras

$$(A, \sum_{i=1}^n p_i (-)_i)$$

$$p_i \in [0, 1], \sum_{i=1}^n p_i = 1$$

- convex (affine) maps

$$h \left( \sum_{i=1}^n p_i a_i \right) = \sum_{i=1}^n p_i h(a_i)$$



# Convex Algebras

infinitely many  
finitary operations

convex  
combinations

binary ones  
“suffice”

- algebras

$$(A, \sum_{i=1}^n p_i (-)_i)$$

$$p_i \in [0, 1], \sum_{i=1}^n p_i = 1$$

- convex (affine) maps

$$h \left( \sum_{i=1}^n p_i a_i \right) = \sum_{i=1}^n p_i h(a_i)$$

satisfying

- Projection

$$\sum_{i=1}^n p_i a_i = a_k, \quad p_k = 1$$

- Barycenter

$$\sum_{i=1}^n p_i \left( \sum_{j=1}^m p_{i,j} a_j \right) = \sum_{j=1}^m \left( \sum_{i=1}^n p_i p_{i,j} \right) a_j$$



# Eilenberg-Moore Algebras

convex algebras  
abstractly

$\mathcal{EM}(\mathcal{D})$

- objects

$$\begin{array}{c} \mathcal{D}A \\ \downarrow a \\ A \end{array}$$

satisfying

$$\begin{array}{ccc} A & \xrightarrow{\eta} & \mathcal{D}A \\ & \searrow a & \downarrow a \\ & & A \end{array}$$

$$\begin{array}{ccc} \mathcal{D}\mathcal{D}A & \xrightarrow{\mu} & \mathcal{D}A \\ \mathcal{D}a \downarrow & & \downarrow a \\ \mathcal{D}A & \xrightarrow{a} & A \end{array}$$

- morphisms

$$\begin{array}{c} \mathcal{D}A \\ \downarrow a \\ A \end{array} \xrightarrow{h} \begin{array}{c} \mathcal{D}B \\ \downarrow b \\ B \end{array}$$

$$\begin{array}{ccc} \mathcal{D}A & \xrightarrow{\mathcal{D}h} & \mathcal{D}B \\ a \downarrow & & \downarrow b \\ A & \xrightarrow{h} & B \end{array}$$



# What is $\mathcal{D}$ ?

Probability distribution functor on **Sets**

$$\mathcal{D}X = \{\mu: X \rightarrow [0, 1] \mid \sum_{x \in X} \mu(x) = 1\}$$

# Free Convex Algebras

# Free Convex Algebras

$$\mathbb{D}_X = (\mathcal{D}X, \sum_{i=1}^n p_i(-)_i)$$

# Free Convex Algebras

$$\mathbb{D}_X = (\mathcal{D}X, \sum_{i=1}^n p_i(-)_i) \quad p_i \in [0, 1], \sum_{i=1}^n p_i = 1$$

# Free Convex Algebras

carried by distributions

$$\mathbb{D}_X = (\mathcal{D}X, \sum_{i=1}^n p_i(-)_i)$$

$$p_i \in [0, 1], \sum_{i=1}^n p_i = 1$$

# Free Convex Algebras

carried by distributions

convex combinations as expected

$$\mathbb{D}_X = (\mathcal{D}X, \sum_{i=1}^n p_i(-)_i) \quad p_i \in [0, 1], \sum_{i=1}^n p_i = 1$$



# Free Convex Algebras

carried by distributions

convex combinations as expected

$$\mathbb{D}_X = (\mathcal{D}X, \sum_{i=1}^n p_i(-)_i) \quad p_i \in [0, 1], \sum_{i=1}^n p_i = 1$$

$$\sum p_i \xi_i = \xi \quad \Leftrightarrow \quad \forall x \in X. \xi(x) = \sum p_i \xi_i(x)$$

# Free Convex Algebras

carried by distributions

convex combinations as expected

wherever there are distributions, there is convexity

$$\mathbb{D}_X = (\mathcal{D}X, \sum_{i=1}^n p_i(-)_i) \quad p_i \in [0, 1], \sum_{i=1}^n p_i = 1$$

$$\sum p_i \xi_i = \xi \quad \Leftrightarrow \quad \forall x \in X. \xi(x) = \sum p_i \xi_i(x)$$

# Free Convex Algebras

carried by distributions


convex combinations as expected

wherever there are distributions, there is convexity

$$\mathbb{D}_X = (\mathcal{D}X, \sum_{i=1}^n p_i(-)_i) \quad p_i \in [0, 1], \sum_{i=1}^n p_i = 1$$

$$\sum p_i \xi_i = \xi \quad \Leftrightarrow \quad \forall x \in X. \xi(x) = \sum p_i \xi_i(x)$$

finitely generated free convex algebras are simplexes




# Convexity in Probabilistic Systems Semantics

# Traces

## Generative PTS

$\mathcal{D} (1 + A \times (-))$




$$\text{tr}(x_1)(ab) = \frac{1}{6} \quad \text{tr}(x_1)(ac) = \frac{1}{8}$$

$$\text{tr}: X \rightarrow \mathcal{D}A^*$$

# Traces via determinisation

## Generative PTS

$$\mathcal{D}(1 + A \times (-))$$



$$\text{tr}(x_1)(ab) = \frac{1}{6} \quad \text{tr}(x_1)(ac) = \frac{1}{8}$$

$$\text{tr}: X \rightarrow \mathcal{D}A^*$$

trace = bisimilarity after  
determinisation

Happens in  
convex algebra

# Trace axioms for generative PTS

Axioms for bisimilarity



$$p \cdot a \cdot (p_1 E_1 \oplus p_2 E_2) \equiv p_1 \cdot a \cdot p E_1 \oplus p_2 \cdot a \cdot p E_2 \quad (D)$$

# Trace axioms for generative PTS

Axioms for bisimilarity



$$p \cdot a \cdot (p_1 E_1 \oplus p_2 E_2) \equiv p_1 \cdot a \cdot p E_1 \oplus p_2 \cdot a \cdot p E_2 \quad (D)$$

[Silva, S. MFPS'11]



# Trace axioms for generative PTS

Axioms for bisimilarity



$$p \cdot a \cdot (p_1 E_1 \oplus p_2 E_2) \equiv p_1 \cdot a \cdot p E_1 \oplus p_2 \cdot a \cdot p E_2 \quad (D)$$

Happens in  
convex algebra

[Silva, S. MFPS'11]

# Trace axioms for generative PTS

Axioms for bisimilarity



$$p \cdot a \cdot (p_1 E_1 \oplus p_2 E_2) \equiv p_1 \cdot a \cdot p E_1 \oplus p_2 \cdot a \cdot p E_2 \quad (D)$$

soundness and  
completeness



Happens in  
convex algebra

[Silva, S. MFPS'11]

# Trace axioms for generative PTS

## Generative PTS

$\mathcal{D}(1 + A \times (-))$



$$\frac{1}{4} \cdot a \cdot \frac{1}{2} \cdot c \cdot 1 \cdot * \stackrel{(D)}{=} \frac{1}{2} \cdot a \cdot \frac{1}{4} \cdot c \cdot 1 \cdot *$$

$$\begin{aligned} \left( \frac{1}{2} \cdot a \cdot \frac{1}{3} \cdot b \cdot 1 \cdot * \right) \oplus \left( \frac{1}{4} \cdot a \cdot \frac{1}{2} \cdot c \cdot 1 \cdot * \right) &\stackrel{(Cong)}{=} \left( \frac{1}{2} \cdot a \cdot \frac{1}{3} \cdot b \cdot 1 \cdot * \right) \oplus \left( \frac{1}{2} \cdot a \cdot \frac{1}{4} \cdot c \cdot 1 \cdot * \right) \\ &\stackrel{(D)}{=} \frac{1}{2} \cdot a \cdot \left( \frac{1}{3} \cdot b \cdot 1 \cdot * \oplus \frac{1}{4} \cdot c \cdot 1 \cdot * \right) \end{aligned}$$

# The quest for completeness

Inspired lots of new research:


- A. S., H. Woracek [Congruences of convex algebras JPAA'15](#)
- S. Milius [Proper functors CALCO'17](#)

# The quest for completeness

Inspired lots of new research:

- A. S., H. Woracek *Congruences of convex algebras* JPAA'15

- S. Milius *Proper functors* CALCO'17




if  $f.p. = f.g.$  and  
then completeness

# The quest for completeness

Inspired lots of new research:

- A. S., H. Woracek *Congruences of convex algebras* JPAA'15
- S. Milius *Proper functors* CALCO'17



if f.p. = f.g. and  
then completeness



does not hold

# The quest for completeness

Inspired lots of new research:

- A. S., H. Woracek *Congruences of convex algebras* JPAA'15

- S. Milius *Proper functors* CALCO'17

our axiomatisation would  
be proven complete if  
only one particular  
convex functor were  
proper

if  $f.p. = f.g.$  and  
then completeness

does not hold

# The quest for completeness

Inspired lots of new research:

- A. S., H. Woracek *Congruences of convex algebras* JPAA'15

- S. Milius *Proper functors* CALCO'17

our axiomatisation would  
be proven complete if  
only one particular  
convex functor were  
proper

it works !

if  $f.p. = f.g.$  and  
then completeness

does not hold



# The quest for completeness

Inspired lots of new research:

- A. S., H. Woracek *Congruences of convex algebras* JPAA'15

- S. Milius *Proper functors* CALCO'17

our axiomatisation would  
be proven complete if  
only one particular  
convex functor were  
proper

it works !

if  $f.p. = f.g.$  and  
then completeness

does not hold

[S., Woracek :-)'17]

# Finely generated, finely presentable

# Finitely generated, finitely presentable

finitely generated (f.g.) = quotients of free finitely generated ones

# Finitely generated, finitely presentable

finitely generated (f.g.) = quotients of free finitely generated ones

finitely presentable (f.p.) = quotients of free finitely generated ones  
under finitely generated congruences

# Finitely generated, finitely presentable

finitely generated (f.g.) = quotients of free finitely generated ones

finitely presentable (f.p.) = quotients of free finitely generated ones  
under finitely generated congruences

smallest congruence  
containing a finite set of pairs

# Finitely generated, finitely presentable

finitely generated (f.g.) = quotients of free finitely generated ones

finitely presentable (f.p.) = quotients of free finitely generated ones  
under finitely generated congruences

smallest congruence  
containing a finite set of pairs

## Theorem

Every congruence of convex algebras is f.g.  
Hence f.p. = f.g.



# Finitely generated, finitely presentable

finitely generated (f.g.) = quotients of free finitely generated ones

finitely presentable (f.p.) = quotients of free finitely generated ones  
under finitely generated congruences

smallest congruence  
containing a finite set of pairs

## Theorem

Every congruence of convex algebras is f.g.  
Hence f.p. = f.g.

[S., Woracek JPAA'15]


# Determinisations



# Determinisations

## Generative PTS


$$X \rightarrow \mathcal{D} (1 + A \times X)$$



# Determinisations

## Generative PTS



$$X \rightarrow \mathcal{D} (1 + A \times X)$$



# Determinisations

## Generative PTS



$$X \rightarrow \mathcal{D} (1 + A \times X)$$



# Determinisations

## Generative PTS

$$X \rightarrow \mathcal{D} (1 + A \times X)$$




[Silva, S. MFPS'11]

[Jacobs, Silva, S. JCSS'15]


# Determinisations

Generative PTS

$$X \rightarrow \mathcal{D} (1 + A \times X)$$



belief-state  
transformer



[Silva, S. MFPS'11]


[Jacobs, Silva, S. JCSS'15]

# Belief-state transformers

# Belief-state transformers

MC


$$X \rightarrow \mathcal{D}(X)$$



# Belief-state transformers

MC

$$X \rightarrow \mathcal{D}(X)$$






# Belief-state transformers

MC

$$X \rightarrow \mathcal{D}(X)$$




$$\begin{array}{c} \frac{1}{3}x_1 + \frac{2}{3}x_2 \quad \dots \\ \Downarrow \\ \frac{7}{9}x_2 + \frac{1}{18}x_3 + \frac{1}{6}x_4 \end{array}$$

# Belief-state transformers

MC

$$X \rightarrow \mathcal{D}(X)$$




belief-state  
transformer

$$\begin{array}{c} \frac{1}{3}x_1 + \frac{2}{3}x_2 \quad \dots \\ \Downarrow \\ \frac{7}{9}x_2 + \frac{1}{18}x_3 + \frac{1}{6}x_4 \end{array}$$

# Belief-state transformers

MC

$$X \rightarrow \mathcal{D}(X)$$



belief-state  
transformer


belief state

$$\begin{array}{c} \frac{1}{3}x_1 + \frac{2}{3}x_2 \quad \dots \\ \Downarrow \\ \frac{7}{9}x_2 + \frac{1}{18}x_3 + \frac{1}{6}x_4 \end{array}$$

# Belief-state transformers

MC

$$X \rightarrow \mathcal{D}(X)$$



belief-state  
transformer

belief state

$$\frac{1}{3} \left( \frac{1}{3}x_2 + \frac{1}{6}x_3 + \frac{1}{2}x_4 \right) + \frac{2}{3}(1x_2)$$


$$\begin{array}{c} \frac{1}{3}x_1 + \frac{2}{3}x_2 \quad \dots \\ \Downarrow \\ \frac{7}{9}x_2 + \frac{1}{18}x_3 + \frac{1}{6}x_4 \end{array}$$

# Belief-state transformers

# Belief-state transformers

PA


$$X \rightarrow (\mathcal{PD}(X))^A$$



# Belief-state transformers

PA


$$X \rightarrow (\mathcal{PD}(X))^A$$



# Belief-state transformers

PA

$$X \rightarrow (\mathcal{PD}(X))^A$$




$$\begin{array}{ccc}
 & \frac{1}{3}x_1 + \frac{2}{3}x_2 & \dots \\
 & \swarrow a & \searrow a \\
 \frac{8}{9}x_2 + \frac{1}{9}x_3 & \dots & \frac{2}{3}x_2 + \frac{1}{6}x_3 + \frac{1}{6}x_4
 \end{array}$$



# Belief-state transformers

PA

$$X \rightarrow (\mathcal{PD}(X))^A$$




belief-state  
transformer

$$\begin{array}{ccc}
 & \frac{1}{3}x_1 + \frac{2}{3}x_2 & \dots \\
 & \swarrow a & \searrow a \\
 \frac{8}{9}x_2 + \frac{1}{9}x_3 & \dots & \frac{2}{3}x_2 + \frac{1}{6}x_3 + \frac{1}{6}x_4
 \end{array}$$

# Belief-state transformers

PA

$$X \rightarrow (\mathcal{PD}(X))^A$$



belief-state  
transformer


belief state

$$\begin{array}{ccccc}
 & & \frac{1}{3}x_1 + \frac{2}{3}x_2 & & \dots \\
 & \swarrow a & & \searrow a & \\
 \frac{8}{9}x_2 + \frac{1}{9}x_3 & & \dots & & \frac{2}{3}x_2 + \frac{1}{6}x_3 + \frac{1}{6}x_4
 \end{array}$$

# Belief-state transformers

PA

$$X \rightarrow (\mathcal{PD}(X))^A$$



belief-state  
transformer


belief state

$$\begin{array}{ccc}
 & \frac{1}{3}x_1 + \frac{2}{3}x_2 & \dots \\
 \swarrow a & & \searrow a \\
 \frac{8}{9}x_2 + \frac{1}{9}x_3 & \dots & \frac{2}{3}x_2 + \frac{1}{6}x_3 + \frac{1}{6}x_4
 \end{array}$$

# Belief-state transformers

PA

$$X \rightarrow (\mathcal{PD}(X))^A$$



belief-state  
transformer

belief state

$$\frac{1}{3} \left( \frac{2}{3}x_2 + \frac{1}{3}x_3 \right) + \frac{2}{3}(1x_2)$$


$$\frac{8}{9}x_2 + \frac{1}{9}x_3 \quad \dots \quad \frac{2}{3}x_2 + \frac{1}{6}x_3 + \frac{1}{6}x_4$$

# Belief-state transformer

PA

$$X \rightarrow \mathcal{P}(\mathcal{D}(X))^A$$


foundation ?



via a generalised  
determinisation



LTS on free  
convex algebra




# Belief-state transformer

PA

$$X \rightarrow \mathcal{P}(\mathcal{D}(X))^A$$

foundation ?




via a generalised  
determinisation



Minkowski  
sum

LTS on free  
convex algebra




# Belief-state transformer

PA

$$X \rightarrow \mathcal{P}(\mathcal{D}(X))^A$$

foundation ?




via a generalised  
determinisation



Minkowski  
sum

LTS on free  
convex algebra

[Bonchi, Silva, S. CONCUR'17]



# Probabilistic Automata



# Probabilistic Automata

Can be given different semantics:

1. Bisimilarity
2. Convex bisimilarity
3. Distribution bisimilarity

# Probabilistic Automata

Can be given different semantics:

1. Bisimilarity

strong  
bisimilarity

2. Convex bisimilarity

3. Distribution bisimilarity

# Probabilistic Automata

Can be given different semantics:

1. Bisimilarity

strong  
bisimilarity

2. Convex bisimilarity

probabilistic /  
combined  
bisimilarity

3. Distribution bisimilarity

# Probabilistic Automata

Can be given different semantics:

1. Bisimilarity

strong  
bisimilarity

2. Convex bisimilarity


probabilistic /  
combined  
bisimilarity

3. Distribution bisimilarity


belief-state  
bisimilarity

# Distribution bisimilarity


# Distribution bisimilarity



# Distribution bisimilarity




# Distribution bisimilarity







# Distribution bisimilarity




# Distribution bisimilarity



# Distribution bisimilarity




# Distribution bisimilarity



# Distribution bisimilarity

$\sim_d$   
largest distribution  
bisimulation

distribution  
bisimulation



transition in  
the  
belief-state  
transformer

$\sim_d$   
is LTS bisimilarity on  
the belief-state  
transformer

# Coinductive proof method for distribution bisimilarity

[Bonchi, Silva, S. CONCUR'17]

# Coinductive proof method for distribution bisimilarity


[Bonchi, Silva, S. CONCUR'17]

bisimulation  
up-to  
convex hull

$R$

# Coinductive proof method for distribution bisimilarity

[Bonchi, Silva, S. CONCUR'17]



$R$




bisimulation  
up-to  
convex hull




# Coinductive proof method for distribution bisimilarity

[Bonchi, Silva, S. CONCUR'17]




# Coinductive proof method for distribution bisimilarity

[Bonchi, Silva, S. CONCUR'17]




# Coinductive proof method for distribution bisimilarity

[Bonchi, Silva, S. CONCUR'17]



# Coinductive proof method for distribution bisimilarity


[Bonchi, Silva, S. CONCUR'17]



# Coinductive proof method for distribution bisimilarity

[Bonchi, Silva, S. CONCUR'17]

bisimulation  
up-to  
convex hull




convex and equivalence  
closure of  $R$

to prove  $\mu \sim_d \nu$   
it suffices to find a  
bisimulation up-to  
convex hull  $R$   
with  $\mu R \nu$

# Coinductive proof method for distribution bisimilarity

[Bonchi, Silva, S. CONCUR'17]






# Coinductive proof method for distribution bisimilarity

[Bonchi, Silva, S. CONCUR'17]

bisimulation  
up-to  
convex hull



convex and equivalence  
closure of  $R$

to prove  $\mu \sim_d \nu$   
it suffices to find a  
bisimulation up-to  
convex hull  $R$   
with  $\mu R \nu$

there always  
exists a finite  
one!

by S., Woracek JPAA'15]

# Termination?

- We looked at one-point extensions of convex algebras, for termination.
- What are all the possible ways?



# Termination?

- We looked at one-point extensions of convex algebras, for termination.

Every convex algebra can be extended by a single point

- What are all the possible ways?

# Termination?

- We looked at one-point extensions of convex algebras, for termination.

Every convex algebra can be extended by a single point

- What are all the possible ways?

there are many possible ways

we can give full description for...

single naturally functorial way

# Termination?

- We looked at one-point extensions of convex algebras, for termination.

Every convex algebra can be extended by a single point

- What are all the possible ways?

there are many possible ways

we can give full description for...

single naturally functorial way

MC and PA  
belief-state transformers

# Termination?

- We looked at one-point extensions of convex algebras, for termination.

Every convex algebra can be extended by a single point

- What are all the possible ways?

there are many possible ways

we can give full description for...

single naturally functorial way

MC and PA  
belief-state transformers

[S., Woracek CALCO'17]

$$\mathbb{X} = \mathbb{D}_X$$

# Free convex algebras

carrier

$$X_* = X + 1 = X \cup \{*\}$$

Possible extensions  $\mathbb{X}_*$  are:

- the black-hole extension

$$px + (1 - p)* = *$$

- $*$  imitates a point  $w \in X$

$$px + (1 - p)* = px + (1 - p)w$$

- $*$  imitates one of the extremal points  $s \in S$  on all other points, and **adheres** this point

$$px + (1 - p)* = px + (1 - p)s, \quad x \neq s$$

$$ps + (1 - p)* = *$$

these are all extensions!

# It's time to terminate this talk..

convexity is  
everywhere  
in probabilistic systems  
semantics

next: algorithms  
for computing  
distribution  
bisimilarity

Thank You!