

A NEW PERSPECTIVE ON THE STEIN–TOMAS THEOREM

Ana Emilia de Orellana.
aedo1@st-andrews.ac.uk

University of St Andrews
Joint work with Marc Carnovale and Jonathan Fraser

One World Fractals

The restriction/extension problem

Let S be a compact *Lebesgue null set* supporting a measure μ .
Let $f \in L^q(\mathbb{R}^d)$. When does $\hat{f}|_S$ *make sense*?

The restriction/extension problem

Let S be a compact *Lebesgue null set* supporting a measure μ .
Let $f \in L^{q'}(\mathbb{R}^d)$. When does $\widehat{f}|_S$ *make sense*?

Find the optimal range of $p, q \in [1, \infty]$ such that

$$\|\widehat{f}\|_{L^{p'}(\mu)} \lesssim \|f\|_{L^{q'}(\mathbb{R}^d)} \quad (\text{restriction}).$$

The restriction/extension problem

Let S be a compact *Lebesgue null set* supporting a measure μ .
Let $f \in L^q(\mathbb{R}^d)$. When does $\widehat{f}|_S$ *make sense*?

Find the optimal range of $p, q \in [1, \infty]$ such that

$$\|\widehat{f}\|_{L^{p'}(\mu)} \lesssim \|f\|_{L^q(\mathbb{R}^d)} \quad (\text{restriction}).$$

- If $f \in L^1(\mathbb{R}^d)$, \widehat{f} is continuous and bounded, so $\widehat{f} \in L^{p'}(\mu)$ for all $p' \geq 1$.

The restriction/extension problem

Let S be a compact *Lebesgue null set* supporting a measure μ .
Let $f \in L^{q'}(\mathbb{R}^d)$. When does $\widehat{f}|_S$ *make sense*?

Find the optimal range of $p, q \in [1, \infty]$ such that

$$\|\widehat{f}\|_{L^{p'}(\mu)} \lesssim \|f\|_{L^{q'}(\mathbb{R}^d)} \quad (\text{restriction}).$$

- If $f \in L^1(\mathbb{R}^d)$, \widehat{f} is continuous and bounded, so $\widehat{f} \in L^{p'}(\mu)$ for all $p' \geq 1$.
- If $f \in L^2(\mathbb{R}^d)$, $\widehat{f} \in L^2(\mathbb{R}^d)$, so \widehat{f} is only defined almost everywhere.

The relevant range is $1 < q' < 2$.

The restriction/extension problem

Let S be a compact *Lebesgue null set* supporting a measure μ .
Let $f \in L^{q'}(\mathbb{R}^d)$. When does $\widehat{f}|_S$ *make sense*?

Find the optimal range of $p, q \in [1, \infty]$ such that

$$\|\widehat{f}\|_{L^{p'}(\mu)} \lesssim \|f\|_{L^{q'}(\mathbb{R}^d)} \quad (\text{restriction}).$$

- If $f \in L^1(\mathbb{R}^d)$, \widehat{f} is continuous and bounded, so $\widehat{f} \in L^{p'}(\mu)$ for all $p' \geq 1$.
- If $f \in L^2(\mathbb{R}^d)$, $\widehat{f} \in L^2(\mathbb{R}^d)$, so \widehat{f} is only defined almost everywhere.

The relevant range is $1 < q' < 2$.

Equivalently, for all $f \in L^p(\mu)$ and $q > 2$,

$$\|\widehat{f\mu}\|_{L^q(\mathbb{R}^d)} \lesssim \|f\|_{L^p(\mu)} \quad (\text{extension}).$$

Some history

Let σ^{d-1} be the surface measure on S^{d-1} .

- Stein (1970's): Curvature causes Fourier transforms to decay.

$$\|\widehat{f\sigma^{d-1}}\|_{L^q(\mathbb{R}^d)} \lesssim \|f\|_{L^2(\sigma^{d-1})}, \quad (1)$$

holds for $q > \frac{4d}{d-1}$.

Some history

Let σ^{d-1} be the surface measure on S^{d-1} .

- Stein (1970's): Curvature causes Fourier transforms to decay.

$$\|\widehat{f\sigma^{d-1}}\|_{L^q(\mathbb{R}^d)} \lesssim \|f\|_{L^2(\sigma^{d-1})}, \quad (1)$$

holds for $q > \frac{4d}{d-1}$.

- Tomas (1975): For arbitrary $d \geq 2$, (1) holds for $q \geq \frac{2(d+1)}{d-1}$.

Some history

Let σ^{d-1} be the surface measure on S^{d-1} .

- Stein (1970's): Curvature causes Fourier transforms to decay.

$$\|\widehat{f\sigma^{d-1}}\|_{L^q(\mathbb{R}^d)} \lesssim \|f\|_{L^2(\sigma^{d-1})}, \quad (1)$$

holds for $q > \frac{4d}{d-1}$.

- Tomas (1975): For arbitrary $d \geq 2$, (1) holds for $q \geq \frac{2(d+1)}{d-1}$.

Why σ^{d-1} ?

Some history

Let σ^{d-1} be the surface measure on S^{d-1} .

- Stein (1970's): Curvature causes Fourier transforms to decay.

$$\|\widehat{f\sigma^{d-1}}\|_{L^q(\mathbb{R}^d)} \lesssim \|f\|_{L^2(\sigma^{d-1})}, \quad (1)$$

holds for $q > \frac{4d}{d-1}$.

- Tomas (1975): For arbitrary $d \geq 2$, (1) holds for $q \geq \frac{2(d+1)}{d-1}$.

Why σ^{d-1} ? Two reasons:

- $\sigma^{d-1}(B(e, r)) \lesssim r^{d-1}$, for all $e \in S^{d-1}$, $r > 0$.
- $\sup_{\xi \in \mathbb{R}^d} |\widehat{\sigma^{d-1}}(\xi)|^2 |\xi|^{d-1} < \infty$.

General measures

Tomas' proof was generalised by Mockenhaupt (2000), Mitsis (2002) and Bak-Seeger (2011).

Tomas' proof was generalised by Mockenhaupt (2000), Mitsis (2002) and Bak-Seeger (2011).

Theorem (Stein–Tomas Theorem)

Let μ be a non-zero, finite, compactly supported, Borel measure on \mathbb{R}^d with

$$\begin{aligned}\mu(B(x, r)) &\lesssim r^\alpha, \quad \forall x \in \mathbb{R}^d, r > 0; \\ \sup_{\xi \in \mathbb{R}^d} |\widehat{\mu}(\xi)|^2 |\xi|^\beta &< \infty.\end{aligned}$$

Then, if $q \geq 2 + \frac{4(d-\alpha)}{\beta}$, for all $f \in L^2(\mu)$,

$$\|\widehat{f\mu}\|_{L^q(\mathbb{R}^d)} \lesssim \|f\|_{L^2(\mu)}.$$

Fourier and Frostman dimensions

Given μ on \mathbb{R}^d , define the Frostman and Fourier dimensions of μ respectively as

$$\dim_{\text{Fr}} \mu = \sup\{\alpha \in \mathbb{R} : \mu(B(x, r)) \lesssim r^\alpha, \forall x \in \mathbb{R}^d, \forall r > 0\};$$

and

$$\dim_{\text{F}} \mu = \sup\{\beta \in \mathbb{R} : \sup_{\xi \in \mathbb{R}^d} |\widehat{\mu}(\xi)|^2 |\xi|^\beta < \infty\}.$$

Fourier and Frostman dimensions

Given μ on \mathbb{R}^d , define the Frostman and Fourier dimensions of μ respectively as

$$\dim_{\text{Fr}} \mu = \sup\{\alpha \in \mathbb{R} : \mu(B(x, r)) \lesssim r^\alpha, \forall x \in \mathbb{R}^d, \forall r > 0\};$$

and

$$\dim_{\text{F}} \mu = \sup\{\beta \in \mathbb{R} : \sup_{\xi \in \mathbb{R}^d} |\widehat{\mu}(\xi)|^2 |\xi|^\beta < \infty\}.$$

They satisfy that $\min\{\frac{\dim_{\text{F}} \mu}{2}, d\} \leq \dim_{\text{Fr}} \mu$.

Fourier and Frostman dimensions

Given μ on \mathbb{R}^d , define the Frostman and Fourier dimensions of μ respectively as

$$\dim_{\text{Fr}} \mu = \sup\{\alpha \in \mathbb{R} : \mu(B(x, r)) \lesssim r^\alpha, \forall x \in \mathbb{R}^d, \forall r > 0\};$$

and

$$\dim_{\text{F}} \mu = \sup\{\beta \in \mathbb{R} : \sup_{\xi \in \mathbb{R}^d} |\widehat{\mu}(\xi)|^2 |\xi|^\beta < \infty\}.$$

They satisfy that $\min\{\frac{\dim_{\text{F}} \mu}{2}, d\} \leq \dim_{\text{Fr}} \mu$.

In particular, the Stein–Tomas theorem can be stated (without the endpoint) as

Theorem (Stein–Tomas Theorem)

Let μ be a non-zero, finite, compactly supported, Borel measure on \mathbb{R}^d . Then if $q > 2 + \frac{4(d - \dim_{\text{Fr}} \mu)}{\dim_{\text{F}} \mu}$, for all $f \in L^2(\mu)$,

$$\|\widehat{f\mu}\|_{L^q(\mathbb{R}^d)} \lesssim \|f\|_{L^2(\mu)}.$$

Fourier and Frostman dimensions

Given μ on \mathbb{R}^d , define the Frostman and Fourier dimensions of μ respectively as

$$\dim_{\text{Fr}} \mu = \sup\{\alpha \in \mathbb{R} : \mu(B(x, r)) \lesssim r^\alpha, \forall x \in \mathbb{R}^d, \forall r > 0\};$$

and

$$\dim_{\text{F}} \mu = \sup\{\beta \in \mathbb{R} : \sup_{\xi \in \mathbb{R}^d} |\widehat{\mu}(\xi)|^2 |\xi|^\beta < \infty\}.$$

They satisfy that $\min\{\frac{\dim_{\text{F}} \mu}{2}, d\} \leq \dim_{\text{Fr}} \mu$.

In particular, the Stein–Tomas theorem can be stated (without the endpoint) as

Theorem (Stein–Tomas Theorem)

Let μ be a non-zero, finite, compactly supported, Borel measure on \mathbb{R}^d with $\dim_{\text{Fr}} \mu = \alpha$. Then if $q > 2 + \frac{4(d-\alpha)}{\dim_{\text{F}} \mu}$, for all $f \in L^2(\mu)$,

$$\|\widehat{f\mu}\|_{L^q(\mathbb{R}^d)} \lesssim \|f\|_{L^2(\mu)}.$$

Fourier and Frostman dimensions

In particular, the Stein–Tomas theorem can be stated (without the endpoint) as

Theorem (Stein–Tomas Theorem)

Let μ be a non-zero, finite, compactly supported, Borel measure on \mathbb{R}^d with $\dim_{\text{Fr}} \mu = \alpha$. Then if $q > 2 + \frac{4(d-\alpha)}{\dim_{\text{F}} \mu}$, for all $f \in L^2(\mu)$,

$$\|\widehat{f\mu}\|_{L^q(\mathbb{R}^d)} \lesssim \|f\|_{L^2(\mu)}.$$

Question

Can more information about the *dimensionality of μ* give us a better range for the restriction estimate to hold?

The Fourier spectrum

Given μ a finite, compactly supported, Borel measure on \mathbb{R}^d and $\theta \in (0, 1]$

$$\dim_{\mathbb{F}}^{\theta} \mu = \sup \left\{ s \in \mathbb{R} : \int_{\mathbb{R}^d} |\widehat{\mu}(\xi)|^{\frac{2}{\theta}} |\xi|^{\frac{s}{\theta} - d} d\xi < \infty \right\}.$$

$$\theta = 0 \quad \longrightarrow \quad \sup_{\xi \in \mathbb{R}^d} |\widehat{\mu}(\xi)|^2 |\xi|^s < \infty$$

The Fourier spectrum

Given μ a finite, compactly supported, Borel measure on \mathbb{R}^d and $\theta \in (0, 1]$

$$\dim_{\mathbb{F}}^{\theta} \mu = \sup \left\{ s \in \mathbb{R} : \int_{\mathbb{R}^d} |\widehat{\mu}(\xi)|^{\frac{2}{\theta}} |\xi|^{\frac{s}{\theta} - d} d\xi < \infty \right\}.$$
$$\theta = 0 \quad \longrightarrow \quad \sup_{\xi \in \mathbb{R}^d} |\widehat{\mu}(\xi)|^2 |\xi|^s < \infty$$

Some facts about the Fourier spectrum:

The Fourier spectrum

Given μ a finite, compactly supported, Borel measure on \mathbb{R}^d and $\theta \in (0, 1]$

$$\dim_{\mathbb{F}}^{\theta} \mu = \sup \left\{ s \in \mathbb{R} : \int_{\mathbb{R}^d} |\widehat{\mu}(\xi)|^{\frac{2}{\theta}} |\xi|^{\frac{s}{\theta} - d} d\xi < \infty \right\}.$$
$$\theta = 0 \quad \longrightarrow \quad \sup_{\xi \in \mathbb{R}^d} |\widehat{\mu}(\xi)|^2 |\xi|^s < \infty$$

Some facts about the Fourier spectrum:

- $\dim_{\mathbb{F}}^0 \mu = \dim_{\mathbb{F}} \mu$ and $\dim_{\mathbb{F}}^1 \mu = \dim_{\mathbb{S}} \mu$.

The Fourier spectrum

Given μ a finite, compactly supported, Borel measure on \mathbb{R}^d and $\theta \in (0, 1]$

$$\dim_{\mathbb{F}}^{\theta} \mu = \sup \left\{ s \in \mathbb{R} : \int_{\mathbb{R}^d} |\widehat{\mu}(\xi)|^{\frac{2}{\theta}} |\xi|^{\frac{s}{\theta} - d} d\xi < \infty \right\}.$$
$$\theta = 0 \quad \longrightarrow \quad \sup_{\xi \in \mathbb{R}^d} |\widehat{\mu}(\xi)|^2 |\xi|^s < \infty$$

Some facts about the Fourier spectrum:

- $\dim_{\mathbb{F}}^0 \mu = \dim_{\mathbb{F}} \mu$ and $\dim_{\mathbb{F}}^1 \mu = \dim_{\mathbb{S}} \mu$.
- For each $\theta \in [0, 1]$, $\dim_{\mathbb{F}} \mu \leq \dim_{\mathbb{F}}^{\theta} \mu \leq \dim_{\mathbb{S}} \mu$.

The Fourier spectrum

Given μ a finite, compactly supported, Borel measure on \mathbb{R}^d and $\theta \in (0, 1]$

$$\dim_{\mathbb{F}}^{\theta} \mu = \sup \left\{ s \in \mathbb{R} : \int_{\mathbb{R}^d} |\widehat{\mu}(\xi)|^{\frac{2}{\theta}} |\xi|^{\frac{s}{\theta} - d} d\xi < \infty \right\}.$$
$$\theta = 0 \quad \longrightarrow \quad \sup_{\xi \in \mathbb{R}^d} |\widehat{\mu}(\xi)|^2 |\xi|^s < \infty$$

Some facts about the Fourier spectrum:

- $\dim_{\mathbb{F}}^0 \mu = \dim_{\mathbb{F}} \mu$ and $\dim_{\mathbb{F}}^1 \mu = \dim_{\mathbb{S}} \mu$.
- For each $\theta \in [0, 1]$, $\dim_{\mathbb{F}} \mu \leq \dim_{\mathbb{F}}^{\theta} \mu \leq \dim_{\mathbb{S}} \mu$.
- $\theta \mapsto \dim_{\mathbb{F}}^{\theta} \mu$ is continuous, concave, and non-decreasing.


The Fourier spectrum

Given μ a finite, compactly supported, Borel measure on \mathbb{R}^d and $\theta \in (0, 1]$

$$\dim_{\mathbb{F}}^{\theta} \mu = \sup \left\{ s \in \mathbb{R} : \int_{\mathbb{R}^d} |\widehat{\mu}(\xi)|^{\frac{2}{\theta}} |\xi|^{\frac{s}{\theta} - d} d\xi < \infty \right\}.$$
$$\theta = 0 \quad \longrightarrow \quad \sup_{\xi \in \mathbb{R}^d} |\widehat{\mu}(\xi)|^2 |\xi|^s < \infty$$

Some facts about the Fourier spectrum:

- $\dim_{\mathbb{F}}^0 \mu = \dim_{\mathbb{F}} \mu$ and $\dim_{\mathbb{F}}^1 \mu = \dim_{\mathbb{S}} \mu$.
- For each $\theta \in [0, 1]$, $\dim_{\mathbb{F}} \mu \leq \dim_{\mathbb{F}}^{\theta} \mu \leq \dim_{\mathbb{S}} \mu$.
- $\theta \mapsto \dim_{\mathbb{F}}^{\theta} \mu$ is continuous, concave, and non-decreasing.
- $\dim_{\mathbb{F}}^{\theta} \mu \geq \dim_{\mathbb{F}} \mu + \theta(\dim_{\mathbb{S}} \mu - \dim_{\mathbb{F}} \mu)$.
- $\dim_{\mathbb{F}}^{\theta} \mu \leq \dim_{\mathbb{F}} \mu + d\theta$.



A Stein–Tomas type result for the Fourier spectrum


Theorem (Carnovale–Fraser–dO, 24++)

Let μ be a non-zero, finite, compactly supported, Borel measure on \mathbb{R}^d , with $\dim_{\text{Fr}} \mu = \alpha$ for some $0 < \alpha \leq d$. If

$$q > 2 + 2 \inf_{\substack{\theta \in [0,1] \\ \dim_{\text{Fr}}^{\theta} \mu > d\theta}} \frac{(d - \alpha)(2 - \theta)}{\dim_{\text{Fr}}^{\theta} \mu - \alpha\theta},$$

then for every $f \in L^2(\mu)$,

$$\|\widehat{f\mu}\|_{L^q(\mathbb{R}^d)} \lesssim \|f\|_{L^2(\mu)}.$$



A Stein–Tomas type result using the Fourier spectrum

Theorem (Carnovale–Fraser–dO, 24++)

Let μ be a non-zero, finite, compactly supported, Borel measure on \mathbb{R}^d , with $\dim_{\text{Fr}} \mu = \alpha$ for some $0 < \alpha \leq d$. If

$$q > 2 + 2 \inf_{\substack{\theta \in [0,1] \\ \dim_{\text{F}}^{\theta} \mu > d\theta}} \frac{(d - \alpha)(2 - \theta)}{\dim_{\text{F}}^{\theta} \mu - \alpha\theta},$$

then for every $f \in L^2(\mu)$,

$$\|\widehat{f\mu}\|_{L^q(\mathbb{R}^d)} \lesssim \|f\|_{L^2(\mu)}.$$

What is more, restriction will not hold for

$$q < \sup \left\{ \frac{2}{\theta} : \dim_{\text{F}}^{\theta} \mu < d\theta \right\}.$$

Restriction for the cone

Let

$$C^{d-1} = \{(t, |t|) : t \in \mathbb{R}^{d-1}, |t| \leq 1\} \subseteq \mathbb{R}^d,$$

be the cone on \mathbb{R}^d and ν^{d-1} the surface measure on C^{d-1} .

Conjecture (cone restriction conjecture)

The estimate $\|\widehat{f\nu^{d-1}}\|_{L^q(\mathbb{R}^d)} \lesssim_{p,q} \|f\|_{L^p(\nu^{d-1})}$ holds for all $f \in L^p(\nu^{d-1})$ if and only if

$$\frac{d-2}{p'} \geq \frac{d}{q} \text{ and } q > \frac{2(d-1)}{d-2},$$


and for $p = 2$ (solved) if and only if

$$q \geq \frac{2d}{d-2}.$$

- Cone restriction theorem: $q \geq \frac{2d}{d-2}$.
- Stein–Tomas: restriction holds for $q \geq \max \left\{ 4, \frac{2d}{d-2} \right\}$.
- Hambrook–Łaba: restriction **fails** for $q < \frac{2d}{d-1}$.

- Cone restriction theorem: $q \geq \frac{2d}{d-2}$.
- Stein–Tomas: restriction holds for $q \geq \max \left\{ 4, \frac{2d}{d-2} \right\}$.
- Hambrook–Łaba: restriction **fails** for $q < \frac{2d}{d-1}$.


$$\dim_{\mathbb{F}}^{\theta} \nu^{d-1} = \min \left\{ 2 + (d-1)\theta, (d-2) + \theta \right\}.$$



- Cone restriction theorem: $q \geq \frac{2d}{d-2}$.
- Stein–Tomas: restriction holds for $q \geq \max \left\{ 4, \frac{2d}{d-2} \right\}$.
- Hambrook–Łaba: restriction **fails** for $q < \frac{2d}{d-1}$.

$$\dim_{\mathbb{F}} \nu^{d-1} = \min \left\{ 2 + (d-1)\theta, (d-2) + \theta \right\}.$$


- Fourier spectrum: restriction **holds** for $q > \max \left\{ \frac{3d-4}{d-2}, \frac{2d}{d-2} \right\}$.
- Fourier spectrum: restriction **fails** for $q < \frac{2(d-1)}{d-2}$.



- Cone restriction theorem: $q \geq \frac{2d}{d-2}$.
- Stein–Tomas: restriction holds for $q \geq \max \left\{ 4, \frac{2d}{d-2} \right\}$.
- Hambrook–Łaba: restriction **fails** for $q < \frac{2d}{d-1}$.

$$\dim_{\mathbb{F}}^{\theta} \nu^{d-1} = \min \left\{ 2 + (d-1)\theta, (d-2) + \theta \right\}.$$

- Fourier spectrum: restriction **holds** for $q > \max \left\{ \frac{3d-4}{d-2}, \frac{2d}{d-2} \right\}$.
- Fourier spectrum: restriction **fails** for $q < \frac{2(d-1)}{d-2}$.



Thank you!