Use of SCN features: Yes
Max learning objects: Maximum objects/class
Strategy N° 3

PA Selected Samples prediction using PA training set,
Learning with selected samples classes with no low regional training instances, no extra training categories,
No Calanoida (civ-vi), Cyclopoida, Zooplankton classes in learning set

Confusion Matrix - In percent of Actual Value



Classification Report Matrix
max available learning objects per class




Predictions of discarded taxa from training
Predicted Taxa
Acartia spp. 45.9% (n=277) 0.0% 0.0% 33.3% (n=1) 0.0% 0.0% 1.2% (n=4)
Bivalvia (larvae) 0.0% 0.0% 0.0% 0.0% 0.0% 20.0% (n=5) 32.5% (n=109)
Bryozoa (larvae) 0.0% 0.0% 0.0% 0.0% 0.0% 0.0% 0.6% (n=2)
Calanoida (ci-ciii) 1.5% (n=9) 0.0% 0.0% 0.0% 0.0% 0.0% 0.0%
Calycophorae (nectophore) 0.7% (n=4) 16.7% (n=1) 3.2% (n=1) 0.0% 100.0% (n=3) 0.0% 0.0%
Centropages spp. 0.5% (n=3) 16.7% (n=1) 0.0% 0.0% 0.0% 0.0% 0.0%
Cirripedia (larvae) 3.2% (n=19) 0.0% 3.2% (n=1) 0.0% 0.0% 0.0% 3.3% (n=11)
Copepoda (nauplii) 0.0% 0.0% 0.0% 0.0% 0.0% 0.0% 1.5% (n=5)
Corycaeidae 10.8% (n=65) 0.0% 0.0% 33.3% (n=1) 0.0% 4.0% (n=1) 2.1% (n=7)
Decapoda-brachyura (zoeae) 1.3% (n=8) 0.0% 0.0% 0.0% 0.0% 0.0% 0.3% (n=1)
Decapoda-non brachyura (larvae) 0.2% (n=1) 50.0% (n=3) 0.0% 0.0% 0.0% 0.0% 0.3% (n=1)
Echinodermata (larvae) 0.2% (n=1) 0.0% 0.0% 0.0% 0.0% 0.0% 0.3% (n=1)
Evadne spp. 7.0% (n=42) 0.0% 19.4% (n=6) 33.3% (n=1) 0.0% 48.0% (n=12) 6.0% (n=20)
Fritillaria spp. 0.5% (n=3) 0.0% 0.0% 0.0% 0.0% 0.0% 0.3% (n=1)
Gastropoda (larvae/Limacina) 0.0% 0.0% 0.0% 0.0% 0.0% 0.0% 2.1% (n=7)
Hydrozoa (medusa) 0.2% (n=1) 0.0% 51.6% (n=16) 0.0% 0.0% 0.0% 0.9% (n=3)
Oikopleura spp. 1.3% (n=8) 0.0% 0.0% 0.0% 0.0% 0.0% 0.6% (n=2)
Paracalanus spp. 24.0% (n=145) 0.0% 0.0% 0.0% 0.0% 0.0% 2.4% (n=8)
Podon/Pleopsis spp. 1.0% (n=6) 0.0% 19.4% (n=6) 0.0% 0.0% 28.0% (n=7) 41.8% (n=140)
Polychaeta (larvae) 0.5% (n=3) 0.0% 3.2% (n=1) 0.0% 0.0% 0.0% 3.9% (n=13)
Pseudocalanus spp. 0.8% (n=5) 0.0% 0.0% 0.0% 0.0% 0.0% 0.0%
Tortanus spp. 0.5% (n=3) 16.7% (n=1) 0.0% 0.0% 0.0% 0.0% 0.0%
Calanoida (civ-vi) n=603
Calanus spp. n=6
Ctenophora (larvae) n=31
Cyclopoida (unid) n=3
Euphysa spp. (medusa) n=3
Ostracoda n=25
Zooplankton (unid) n=335
Actual discarded Taxa
100
80
60
40
20
0

Relative Abundance of Top Taxonomic Instances per Sample

Val Pred Val Pred Val Pred Val Pred Val Pred Val Pred Val Pred Val Pred Val Pred Val Pred

1.0
0.8
0.6
0.4
0.2
0.0

Relative Abundance

S11 S12 S13 S14 S15 S16 S17 S18 S19 S20

Sample Short ID

Most abundant classes (In Val or Pred)
- Acartia spp.
- Calanoida (ci–ciii)
- Calanoida (civ–vi)
- Cirripedia (larvae)
- Corycaeidae
- Evadne spp.
- Fritillaria spp.
- Oikopleura spp.
- Paracalanus spp.
- Podon/Pleopsis spp.
- Zooplankton (unid)
- Others