ESCHERICHIA COLI ATCC 8739 ADAPTS SPECIFICALLY TO SODIUM CHLORIDE, MONOSODIUM GLUTAMATE, AND BENZOIC ACID AFTER PROLONGED STRESS

Bryan Zong Lin Loo¹, Samuel Xin Zher Low¹, Zhen Qin Aw¹, Kun Cheng Lee¹, Jack Si Hao Oon¹, Chin How Lee¹ and Maurice Han Tong Ling^{1,2,*}

¹School of Chemical and Life Science, Singapore Polytechnic, Singapore ²Department of Zoology, The University of Melbourne, Australia

ABSTRACT

Escherichia coli lives in the human intestine and any form of adaptation may affect the human body. The effects of food additives on E. coli have been less studied compared to antibiotics. A recent study has demonstrated that E. coli is able to adapt to food additives by demonstrating global stress response. This study continues to study the evolution of E. coli in different food additives (sodium chloride, benzoic acid, monosodium glutamate) in different concentrations, singly or in combination, for over 83 passages. Adaptability of the cells was estimated with generation time and cell density at the stationary phase. Polymerase Chain Reaction (PCR)/ Restriction Fragments Length Polymorphism (RFLP) were used to analyze the adaptation at genomic level. Our results show that adaptation started to slow down and the gradients of generation time against passage are less steep compared with previous study, suggesting that most adaptive mutations occurred within the first 500 generations. In the genomic level, ecological specialization is observed as we find that the cells adapted through a different mechanism and diverge from each other although the resulting effect of the medium is the same. It suggests that different concentrations of food additives cause different types of chemical stress, instead of different levels of chemical stress.

Keywords: Escherichia coli, ATCC 8739, food additive, chemical stress

^{*} Corresponding author: mauriceling@acm.org

Citation: Loo, BZL, Low, SXZ, Aw, ZQ, Lee, KC, Oon, JSH, Lee, CH, Ling, MHT. 2014. Escherichia coli ATCC 8739 Adapts Specifically to Sodium Chloride, Monosodium Glutamate, and Benzoic Acid after Prolonged Stress. Asia Pacific Journal of Life Sciences 7(3): 243-258.

ACKNOWLEDGMENT

This project was funded by Singapore Polytechnic final year project fund (Account Number: 11-27801-45-2550).

REFERENCES

- [1] Koonin, E. V. 2009. Darwinian evolution in the light of genomics. *Nucleic Acids Research*, 37, 1011-34.
- [2] Saccheri, I. J., Rousset, F., Watts, P. C., Brakefield, P. M. & Cook, L. M. 2008. Selection and gene flow on a diminishing cline of melanic peppered moths. *Proceedings of the National Academy of Science USA*, 105, 16212-7.
- [3] Blount, Z. D., Borland, C. Z. & Lenski, R. E. 2008. Historical contingency and the evolution of a key innovation in an experimental population of *Escherichia coli*. *Proceedings of the National Academy of Science USA*, 105, 7899-906.
- [4] Okada, H., Negoro, S., Kimura, H. & Nakamura, S. 1983. Evolutionary adaptation of plasmid-encoded enzymes for degrading nylon oligomers. *Nature*, 306, 203-6.
- [5] Safran, R. J. & Nosil, P. 2012. Speciation: The origin of new species. *Nature Education* Knowledge, 3, 17.
- [6] Lenormard, T. 2012. From local adaptation to speciation: Specialization and reinforcement. *International Journal of Ecology* 2012, Article ID 508458.
- [7] Hurst, L. D. 2009. Fundamental concepts in genetics: genetics and the understanding of selection. *Nature Review Genetics*, 10, 83-93.
- [8] Mank, J. E., Nam, K. & Ellegren, H. 2009. Faster-Z evolution is predominantly due to genetic drift. *Molecular Biology and Evolution*, 27, 661-670.
- [9] Lenski, R. E., Rose, M. R., Simpson, S. C. & Tadler, S. C. 1991. Long-term experimental evolution in Escherichia coli. I. Adaptation and divergence during 2,000 generations. *American Naturalist*, 138, 1315-1341.
- [10] Cooper, V. S., Bennett, A. F. & Lenski, R. E. 2001a. Evolution of thermal dependence of growth rate of *Escherichia coli* populations during 20,000 generations in a constant environment. *Evolution*, 55, 889-96.
- [11] Cooper, V. S., Schneider, D., Blot, M. & Lenski R., E. 2001b. Mechanisms causing rapid and parallel losses of ribose catabolism in evolving populations of *Escherichia coli* B. *Journal of Bacteriology*, 183, 2834-41.
- [12] Goh, D. J. W, How, J. A., Lim, J. Z. R., Ng, W. C., Oon, J. S. H., Lee, K. C., Lee, C. H., & Ling, M. H. T. 2012. Gradual and Step-wise Halophilization enables *Escherichia coli* ATCC 8739 to adapt to 11% NaCl. *Electronic Physician*, 4, 527-535.

- [13] How, J. A., Lim, J. Z. R., Goh, D. J. W., Ng, W. C., Oon, J. S. H., Lee, K. C., Lee, C. H., & Ling, M. H. T. 2013. Adaptation of *Escherichia coli* ATCC 8739 to 11% NaCl. *Dataset Papers in Biology*, 2013, Article ID 219095.
- [14] Low, S. X. Z., Aw, Z. Q., Loo, B. Z. L., Lee, K. C., Oon, J. S. H., Lee, C. H., & Ling, M. H. T. 2013. Viability of *Escherichia coli* ATCC 8739 in Nutrient Broth, Luria-Bertani Broth and Brain Heart Infusion over 11 weeks. *Electronic Physician*, 5, 576-581.
- [15] Travisano, M. & Lenski, R. E. 1996. Long-term experimental evolution in *Escherichia coli*. IV. Targets of selection and the specificity of adaptation. *Genetics*, 143, 15-26.
- [16] Hays, H. C., Millner, P. A., Jones, J. K. & Rayner-Brandes, M. H. 2005. A novel and convenient self-drying system for bacterial preservation. *Journal of Microbiological Methods*, 63, 29-35.
- [17] Chaudhuri, R. R., Sebaihia, M., Hobman, J. L., Webber, M. A., Leyton, D. L., Goldberg, M. D., Cunningham, A. F., Scott-Tucker, A., Ferguson, P. R., Thomas, C. M., Frankel, G., Tang, C. M., Dudley, E. G., Roberts, I. S., Rasko, D. A., Pallen, M. J., Parkhill, J., Nataro, J. P., Thomson, N. R. & Henderson, I. R. 2010. Complete genome sequence and comparative metabolic profiling of the prototypical enteroaggregative *Escherichia coli* strain 042. *PLoS ONE*, 5, e8801.
- [18] Lestari, E. S., Severin, J. A. & Verbrugh, H. A. 2012. Antimicrobial resistance among pathogenic bacteria in Southeast Asia. Southeast Asian Journal of Tropical Medicine and Public Health, 43, 385-422.
- [19] Van Der Beek, D., Brouwer, M. C., Thwaites, G. E. & Tunkel, A. R. 2012. Advances in treatment of bacterial meningitis. *Lancet*, 380, 1693-702.
- [20] Russell, A. D. 1991. Mechanisms of bacterial resistance to non-antibiotics: food additives and food and pharmaceutical preservatives. *Journal of Applied Bacteriology*, 71, 191-201.
- [21] Garner, M. R., James, K. E., Callahan, M. C., Wiedmann, M. & Boor, K. J. 2006. Exposure to salt and organic acids increases the ability of *Listeria monocytogenes* to invade Caco-2 cells but decreases its ability to survive gastric stress. *Applied and Environmental Microbiology*, 72, 5384-95.
- [22] Kesseru, P., Kiss, I., Bihari, Z. & Polyak, B. 2002. The effects of NaCl and some heavy metals on the denitrification activity of *Ochrobactrum anthropi*. *Journal of Basic Microbiology*, 42, 268-76.
- [23] Glaasker, E., Tjan, F. S., Ter Steeg, P. F., Konings, W. N. & Poolman, B. 1998. Physiological response of *Lactobacillus plantarum* to salt and nonelectrolyte stress. *Journal of Bacteriology*, 180, 4718-23.
- [24] Verluyten, J., Messens, W. & De Vuyst, L. 2004. Sodium chloride reduces production of curvacin A, a bacteriocin produced by *Lactobacillus curvatus* strain LTH 1174, originating from fermented sausage. *Applied Environmental Microbiology*, 70, 2271-8.
- [25] Lee, C. H., Oon, J. S., Lee, K. C. & Ling, M. H. T. 2012. Escherichia coli ATCC 8739 adapts to the presence of sodium chloride, monosodium glutamate, and benzoic acid after extended culture. ISRN Microbiology, 2012, Article ID 965356.
- [26] Zhong, S., Miller, S. P., Dykhuizen, D. E. & Dean, A. M. 2009. Transcription, translation, and the evolution of specialists and generalists. *Molecular Biology and Evolution*, 26, 2661–2678.

- [27] Cheng, H. R. & Jiang, N. 2006. Extremely rapid extraction of DNA from bacteria and yeasts. *Biotechnology Letters*, 28, 55-9.
- [28] Sezonov, G., Joseleau-Petit, D. & D'ari, R. (2007) *Escherichia coli* physiology in Luria-Bertani Broth. *Journal of Bacteriology*, 189, 8746-9.
- [29] Nei, M. & Li, W. H. 1979. Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proceedings of the National Academy of Science USA*, 76, 5269-73.
- [30] Chay, Z. E., Lee, C. H., Lee, K. C., Oon, J. S. H., & Ling, M. H. T. 2010. Russel and Rao coefficient is a suitable substitute for Dice coefficient in studying restriction mapped genetic distances of *Escherichia coli*. *Computational and Mathematical Biology*, 1, 1.
- [31] Lee, C. H, Lee, K. C, Oon, J. S. H, & Ling, M. H. T. 2010. Bactome, I: Python in DNA fingerprinting. *The Python Papers*, 5, 6.
- [32] Ling, M. H. T. 2010. COPADS, I: Distances measures between two lists or sets. *The Python Papers* Source Codes, 2, 2.
- [33] De Visser, J. & Lenski, R. 2002. Long-term experimental evolution in *Escherichia coli*. XI. Rejection of non-transitive interations as cause of declining rate of adaptation. *BioMed Central Evolutionary Biology*, 2,1-8.
- [34] Korona, R. 1996. Genetic divergence and fitness convergence under uniform selection in experimental populations of bacteria. *Genetics*, 143, 637-44.
- [35] Cooper, V.S & Lenski, R.E. 2000. The population genetics of ecological specialization in evolving *Esherichia coli* populations. *Nature*, 407, 736-39.
- [36] Ray, S. & Peters, C. A. 2010. Adaptations in microbiological populations exposed to dinitrophenol and other chemical stressors. *Environmental Toxicology and Chemistry*, 29, 2161–68.
- [37] Miller, L. D., Russell, M. H. & Alexandre, G. 2009. Diversity in bacterial chemotactic responses and niche adaptation. *Advances in Applied Microbiology*, 66, 53-75.
- [38] Wu, X., Monchy, S., Taghavi, S., Zhu, W., Ramos, J. & Van Der Lelie, D. 2010. Comparative genomics and functional analysis of niche-specific adaptation in Pseudomonas putida. FEMS Microbiology Reviews, 35, 299-323.
- [39] Siezen, R. J., Tzeneva, V. A., Castioni, A., Wels, M., Phan, H. T., Rademaker, J. L., Starrenburg, M. J., Kleerebezem, M., Molenaar, D. & Van Hylckama Vlieg, J. E. 2010. Phenotypic and genomic diversity of *Lactobacillus plantarum* strains isolated from various environmental niches. *Environmental Microbiology*, 12, 758-73.
- [40] Loui, C., Chang, A. C., Lu, S. 2009. Role of the ArcAB two-component system in the resistance of Escherichia coli to reactive oxygen stress. *BMC Microbiology*, 9, 183.
- [41] Wang, Q., Yu, H., Xia, Y., Kang, Z. & Qi, Q. 2009. Complete PHB mobilization in Escherichia coli enhances the stress tolerance: a potential biotechnological application. *Microbial Cell Factory*, 8, 47.
- [42] Badaoui Najjar, M., Chikidas, M. & Montville, T. J. 2007. Changes in *Listeria monocytogenes* membrane fluidity in response to temperature stress. *Applied and Environmental Microbiology*, 73, 6429-35.
- [43] Friman V. P., Ghoul M., Molin S., Johansen H. K., & Buckling A. 2013. *Pseudomonas aeruginosa* adaptation to lungs of cystic fibrosis patients leads to lowered resistance to phage and protist enemies. *PLoS One*, 8, e75380.