
All transfer families — raw difference
Median
75th percentile
95th percentile
Observed maximum (no CI)
Same-model
pooled; n=5,160; 172 blocks; core30
Same init, different data order
pooled; n=11,760; 210 blocks; core56
Different init, same data order
pooled; n=4,500; 150 blocks; core30
Different init, different data order
pooled; n=26,768; 478 blocks; core56
Reproduction (prv1 ↔ orig)
pooled; n=870; 29 blocks; core30
160M → 410M original-seed lineage
dog; n=330; 11 blocks; core30
410M → 160M original-seed lineage
dog; n=300; 10 blocks; core30
160M → 410M cross-seed lineage
dog; n=300; 10 blocks; core30
410M → 160M cross-seed lineage
dog; n=300; 10 blocks; core30
Pythia 160M → OLMo 1B
dog; n=560; 16 blocks; core35
OLMo 1B → Pythia 160M
dog; n=330; 11 blocks; core30
Same init, different corpus
elephant; n=1,400; 40 blocks; core35
Different init, different corpus
elephant; n=700; 20 blocks; core35
Cross seed (different init, same corpus)
elephant; n=1,050; 30 blocks; core35
0.0
0.1
0.2
0.3
0.4
0.6
0.8
1.0
p(student) − p(control)