

Generalization near a bifurcation point

In this work, we attempt to theoretically arrive at the ability of an abstract generative deep neural network to generalize near critical points like divergences, bifurcations, etc when trained on data away from criticality.

1 Linear Toy Model

1.1 Data-generating process

Let $r \in \mathbb{R}$ be a one-dimensional control parameter. We model a bifurcation at $r = r_c$ by introducing a conditional Gaussian distribution whose covariance becomes singular in one direction as $r \rightarrow r_c$.

Conditional distribution. For each r , samples $\mathbf{x} \in \mathbb{R}^d$ are drawn from

$$\mathbf{x} \mid r \sim \mathcal{N}(\mathbf{0}, \Sigma(r)). \quad (1)$$

Critical covariance. Let $\Sigma(r)$ be the covariance of the distribution given in (??). Since the covariance has a symmetric positive semi-definite structure it has an eigen decomposition as follows,

$$\Sigma(r) = Q(r)\Lambda(r)Q(r)^T \quad (2)$$

where

$$\Lambda(r) = \text{diag}(\lambda_1(r), \dots, \lambda_d(r))$$

Now suppose that as $r \rightarrow r_c$, fluctuations in one direction become increasingly dominant. Mathematically,

$$\lambda_1(r) \sim |r - r_c|^{-\nu},$$

while the other eigenvalues remain bounded:

$$\lambda_k(r) = O(1), \quad k = 2, \dots, d.$$

Then

$$\frac{\lambda_k(r)}{\lambda_1(r)} \rightarrow \infty$$

One direction exhibits parametrically faster growth of variance than the others. [H] =====
 »»»> Stashed changes Without loss of generality, choose coordinates so that the critical direction is the first canonical basis vector $\mathbf{e}_1 = (1, 0, \dots, 0)^\top$. Define

$$\Sigma(r) = I_d + \frac{1}{|r - r_c|^\nu} \mathbf{e}_1 \mathbf{e}_1^\top, \quad (3)$$

where $\nu > 0$ is the critical exponent. The eigenvalues of $\Sigma(r)$ are

$$\sigma_1(r) = 1 + |r - r_c|^{-\nu}, \quad \sigma_k(r) = 1 \quad (k = 2, \dots, d).$$

As $r \rightarrow r_c$, the variance along $\mathbf{e}_1$ diverges as $|r - r_c|^{-\nu}$, mimicking the diverging fluctuations near a critical point. ~~The exponent ν is universal: for the period-doubling cascade in the logistic map at the Feigenbaum point, $\nu = \log 2 / \log \delta_F \approx 0.4498$, with $\delta_F \approx 4.6692$ the Feigenbaum constant.~~




Figure 1: **Data-generating process dynamics and covariance structure across the transition to criticality.** (Top) Representative sample trajectories x_t generated off-criticality ($r = 0.5, \tau = 2.0$, left) and near-criticality ($r = 0.01, \tau = 100.0$, right). As the control parameter $r \rightarrow r_c$, trajectories exhibit marked critical slowing down, characterized by enhanced fluctuation amplitudes and persistent memory. (Bottom) Corresponding empirical covariance matrices $\Sigma_{t,t'}$. Off-criticality (left), correlations remain localized along the main diagonal. Near criticality (right), long-range temporal correlations emerge as the leading eigenvalue diverges according to $\sigma_1(r) = 1 + |r - r_c|^{-\nu}$, establishing a dominant critical direction $\mathbf{e}_1$.

1.2 True score function

The score of a Gaussian $\mathcal{N}(\mathbf{0}, \Sigma)$ is computed by direct differentiation of its log-density. The density is

$$p(\mathbf{x}) = \frac{1}{(2\pi)^{d/2} \det(\Sigma)^{1/2}} \exp\left(-\frac{1}{2}\mathbf{x}^\top \Sigma^{-1} \mathbf{x}\right),$$

so

$$\log p(\mathbf{x}) = -\frac{1}{2}\mathbf{x}^\top \Sigma^{-1} \mathbf{x} + \text{const}, \quad \nabla_{\mathbf{x}} \log p(\mathbf{x}) = -\Sigma^{-1} \mathbf{x}.$$

For our problem,

$$s^*(\mathbf{x}, r) \equiv \nabla_{\mathbf{x}} \log p(\mathbf{x} | r) = -\Sigma(r)^{-1} \mathbf{x}. \quad (4)$$

This inverse can be calculated and we get

$$\Sigma(r)^{-1} = I_d + \left(\frac{1}{\sigma_1(r)} - 1\right) \mathbf{e}_1 \mathbf{e}_1^\top. \quad (5)$$

Critical direction: $1/\sigma_1(r) = |r - r_c|^\nu / (1 + |r - r_c|^\nu) \rightarrow 0$ as $r \rightarrow r_c$, so the score component along $\mathbf{e}_1$ vanishes at the bifurcation. **Non-critical directions:** the score is simply $-x_k$ for $k \geq 2$.

2 Student model and loss function

What is student model, write a small paragraph on your approach to this model. What are you going to gain/find out from the model predictions? We adopt a linear-in- $\mathbf{x}$ student with general r -dependence:

$$s_{\theta}(\mathbf{x}, r) = -A(r)\mathbf{x}, \quad A(r) \in \mathbb{R}^{d \times d}. \quad (6)$$

We parameterize $A(r)$ as a degree- $(K-1)$ polynomial in r in some basis $\{\phi_k(r)\}_{k=0}^{K-1}$ (monomials, Chebyshev, Legendre — the choice does not affect the result). For concreteness:

$$A(r) = \sum_{k=0}^{K-1} A_k \phi_k(r), \quad \phi_k(r) = r^k.$$

The trainable parameters are the K matrices $\{A_k\}$, totaling Kd^2 scalar parameters.

The (population) score-matching loss is **add a small schematic**

$$\mathcal{L}(\theta) = \mathbb{E}_{r \sim \rho_{\Delta}} \left[\mathbb{E}_{\mathbf{x}|r} \left[\|s_{\theta}(\mathbf{x}, r) - s^*(\mathbf{x}, r)\|^2 \right] \right], \quad (7)$$

where the training distribution over r has the gap excised:

$$\rho_{\Delta}(r) = \frac{1}{Z_{\Delta}} \mathbf{1}[|r - r_c| > \Delta/2] \cdot \rho_0(r), \quad (8)$$

with ρ_0 a base distribution (we take uniform on $[r_c - L, r_c + L]$) and Z_{Δ} the normalization.

Let $B(r) \equiv A(r) - \Sigma(r)^{-1}$. Then

$$\|s_{\theta}(\mathbf{x}, r) - s^*(\mathbf{x}, r)\|^2 = \|B(r)\mathbf{x}\|^2 = \mathbf{x}^{\top} B(r)^{\top} B(r) \mathbf{x}.$$

For any deterministic symmetric matrix M , the standard identity for a centered Gaussian $\mathbf{x} \sim \mathcal{N}(0, \Sigma)$ is $\mathbb{E}[\mathbf{x}^{\top} M \mathbf{x}] = \text{Tr}(M \Sigma)$. Hence

$$\mathbb{E}_{\mathbf{x}|r} [\|s_{\theta} - s^*\|^2] = \text{Tr}(B(r)^{\top} B(r) \Sigma(r)) = \text{Tr}(B(r) \Sigma(r) B(r)^{\top}).$$

So the loss becomes

$$\mathcal{L}(\theta) = \int \rho_{\Delta}(r) \text{Tr} \left[(A(r) - \Sigma(r)^{-1}) \Sigma(r) (A(r) - \Sigma(r)^{-1})^{\top} \right] dr. \quad (9)$$

2.1 Decoupling the dimensions

We now exploit the diagonal structure of $\Sigma(r)$. In the canonical basis, $\Sigma(r) = \text{diag}(\sigma_1(r), 1, 1, \dots, 1)$. The cost (??) can be expanded in matrix entries. Off-diagonal entries of $A(r)$ contribute only positive terms to the trace, while $\Sigma(r)^{-1}$ has no off-diagonal part. Hence setting off-diagonal entries of $A(r)$ to zero strictly reduces the cost without affecting the diagonal fit. Writing $A(r) = \text{diag}(a_1(r), a_2(r), \dots, a_d(r))$ with each $a_k(r)$ a degree- $(K-1)$ polynomial, the trace splits as

$$\text{Tr}[\dots] = \sum_{k=1}^d (a_k(r) - 1/\sigma_k(r))^2 \sigma_k(r).$$

Non-critical directions ($k \geq 2$). Here $\sigma_k(r) = 1$ and $1/\sigma_k(r) = 1$, so the k -th term is $(a_k(r) - 1)^2$. The polynomial fit $a_k(r) \equiv 1$ (achievable since the constant function belongs to the polynomial basis) gives zero loss. These directions are trivially learned.

Critical direction ($k = 1$). Drop the subscript and call this $a(r) \equiv a_1(r)$. The relevant loss is

$$\mathcal{L}_1(a) = \int \rho_\Delta(r) (a(r) - 1/\sigma_1(r))^2 \sigma_1(r) dr. \quad (10)$$

Define the *target function* and *weight function*:

$$g(r) \equiv \frac{1}{\sigma_1(r)} = \frac{|r - r_c|^\nu}{1 + |r - r_c|^\nu}, \quad w(r) \equiv \rho_\Delta(r) \sigma_1(r). \quad (11)$$

Then

$$\mathcal{L}_1(a) = \int w(r) (a(r) - g(r))^2 dr. \quad (12)$$

This is a *weighted least-squares polynomial fit* of g on the support of ρ_Δ (i.e., on $|r - r_c| > \Delta/2$). **Expand this section.** Note that

- For $|r - r_c| \ll 1$: $g(r) \approx |r - r_c|^\nu$ — the cusp.
- For $|r - r_c| \rightarrow \infty$: $g(r) \rightarrow 1$ — saturates.
- At $r = r_c$: $g(r_c) = 0$.

3 Minimising loss

We henceforth set $r_c = 0$ by relabeling. Expand $a(r) = \sum_{k=0}^{K-1} \theta_k r^k$, with $\boldsymbol{\theta} = (\theta_0, \dots, \theta_{K-1})^\top$ and $\boldsymbol{\phi}(r) = (1, r, r^2, \dots, r^{K-1})^\top$, so

$$a(r) = \boldsymbol{\theta}^\top \boldsymbol{\phi}(r).$$

The loss becomes a quadratic form in $\boldsymbol{\theta}$:

$$\begin{aligned} \mathcal{L}_1 &= \int w(r) (\boldsymbol{\theta}^\top \boldsymbol{\phi}(r) - g(r))^2 dr \\ &= \boldsymbol{\theta}^\top \left[\int w(r) \boldsymbol{\phi}(r) \boldsymbol{\phi}(r)^\top dr \right] \boldsymbol{\theta} - 2\boldsymbol{\theta}^\top \left[\int w(r) g(r) \boldsymbol{\phi}(r) dr \right] + \int w(r) g(r)^2 dr. \end{aligned} \quad (13)$$

Define

$$M_{kl} \equiv \int w(r) r^k r^l dr = \int \rho_\Delta(r) \sigma_1(r) r^{k+l} dr, \quad (14)$$

$$b_k \equiv \int w(r) g(r) r^k dr. \quad (15)$$

Crucially, $\sigma_1(r)g(r) = 1$, so the bias vector simplifies:

$$b_k = \int \rho_\Delta(r) r^k dr. \quad (16)$$

That is, b_k is the k -th moment of the *unweighted* training distribution ρ_Δ .

Setting $\partial \mathcal{L}_1 / \partial \boldsymbol{\theta} = 0$ gives the normal equations $M\boldsymbol{\theta} = \mathbf{b}$, with solution

$$\hat{\boldsymbol{\theta}} = M^{-1} \mathbf{b}. \quad (17)$$

4 Scaling analysis

Place $r_c = 0$. Define a matching scale r_* of order unity that separates inner and outer regions:

- **Inner region:** $\Delta/2 < |r| < r_*$, with $|r|^\nu \ll 1$, so $\sigma_1(r) \approx |r|^{-\nu}$ and $g(r) \approx |r|^\nu$.
- **Outer region:** $r_* < |r| < L$, with $|r|^\nu \gtrsim 1$, so $\sigma_1(r) \rightarrow 1$ and $g(r) \rightarrow 1$.

4.1 Inner-region contributions

Assume $k + l$ even (only nonvanishing cases). The inner contribution to M_{kl} is

$$\begin{aligned} M_{kl}^{\text{in}}(\Delta) &= \int_{\Delta/2 < |r| < r_*} \rho_0(r) \sigma_1(r) r^{k+l} dr \\ &\approx 2\rho_0 \int_{\Delta/2}^{r_*} r^{-\nu} r^{k+l} dr \\ &= \frac{2\rho_0}{k+l-\nu+1} \left[r_*^{k+l-\nu+1} - (\Delta/2)^{k+l-\nu+1} \right]. \end{aligned} \quad (18)$$

For $\nu < 1$ (Feigenbaum case) and $k+l \geq 0$ even, the exponent $k+l-\nu+1 > 0$, so the integral is finite as $\Delta \rightarrow 0$ and the Δ -dependence enters as

$$M_{kl}^{\text{in}}(\Delta) = M_{kl}^{\text{in}}(0) - \frac{2\rho_0}{k+l-\nu+1} (\Delta/2)^{k+l-\nu+1}. \quad (19)$$

Similarly,

$$\begin{aligned} b_k^{\text{in}}(\Delta) &= \int_{\Delta/2 < |r| < r_*} \rho_0(r) r^k dr \\ &\approx \frac{2\rho_0}{k+1} \left[r_*^{k+1} - (\Delta/2)^{k+1} \right]. \end{aligned} \quad (20)$$

4.2 Outer-region contributions

For the outer region, $\sigma_1(r) \rightarrow 1$, $g(r) \rightarrow 1$, $w(r) \rightarrow \rho_0$:

$$M_{kl}^{\text{out}} \approx \rho_0 \int_{r_* < |r| < L} r^{k+l} dr, \quad b_k^{\text{out}} \approx \rho_0 \int_{r_* < |r| < L} r^k dr. \quad (21)$$

Cite other equations and figures as much as possible. Both are Δ -independent at this level of approximation.

4.3 Expansion of $\hat{\theta}(\Delta)$

Combining,

$$M(\Delta) = M_0 + \Delta^{1-\nu} M_1 + O(\Delta^{3-\nu}), \quad (22)$$

$$\mathbf{b}(\Delta) = \mathbf{b}_0 + \Delta \mathbf{b}_1 + O(\Delta^3), \quad (23)$$

where $M_0, M_1, \mathbf{b}_0, \mathbf{b}_1$ are Δ -independent. The corrections in M scale as $\Delta^{1-\nu}$ because the leading inner integrand $\sim r^{-\nu}$ near the gap edge integrates to give a power of the gap that is one greater (boundary contribution). The corrections in $\mathbf{b}$ scale as Δ because the integrand is regular ($\sim r^0$).

Hence

$$\begin{aligned} \hat{\theta}(\Delta) &= M(\Delta)^{-1} \mathbf{b}(\Delta) \\ &= (M_0 + \Delta^{1-\nu} M_1 + \dots)^{-1} (\mathbf{b}_0 + \Delta \mathbf{b}_1 + \dots) \\ &= M_0^{-1} \mathbf{b}_0 + \Delta^{1-\nu} \left[-M_0^{-1} M_1 M_0^{-1} \mathbf{b}_0 \right] + O(\Delta). \end{aligned} \quad (24)$$

So

$$\hat{\theta}_0(\Delta) = \hat{\theta}_0^{(0)} + c_1 \Delta^{1-\nu} + O(\Delta), \quad (25)$$

where $\hat{\theta}_0^{(0)} = \hat{\theta}_0^{(0)}$ is the limiting value at $\Delta \rightarrow 0$, with $\hat{\theta}^{(0)} = M_0^{-1} \mathbf{b}_0$.




Figure 2: **Deterministic scaling properties of design matrix moments and critical parameters near the excised gap ($\nu = 0.35$).** (a) Absolute deviation of the leading moment matrix element $|M_{00}(\Delta) - M_{00}(0)|$ against the gap size Δ . The numerical trend verifies the theoretical inner-region scaling contribution of $\Delta^{1-\nu}$ derived in Eq. (22). (b) Absolute magnitude of the linear correction term $|b_1|$ vs. Δ . The empirical power-law fit ($\sim \Delta^{1.03}$) matches the analytical boundary scaling expectation of $\Delta^{1.00}$ (Eq. 23). (c) Deviation of the leading polynomial coefficient $|\hat{\theta}_0|$ as a function of Δ . The fitted scaling exponent ($\sim \Delta^{0.65}$) perfectly confirms the leading-order expansion $\hat{\theta}_0(\Delta) = \hat{\theta}_0^{(0)} + c_1 \Delta^{1-\nu} + O(\Delta)$ for $1 - \nu = 0.65$ (Eq. 25). Avoid titles, combine all of them in one figures with different colors + labels. Remove $\Delta < 10^{-7}$




Figure 3: **Training and Critical Mean Squared Error (MSE) dependence on gap size Δ for $\nu = 0.75$.** (Left) Train MSE evaluated on the training distribution away from the gap. (Right) Critical MSE evaluated inside the excised gap near r_c across polynomial degrees $K \in \{2, 4, 6, 8, 10, 16\}$. Increasing model capacity K lowers overall approximation error, while the divergence rate as $\Delta \rightarrow 0$ remains governed by the critical exponent $1 - \nu$. **Combine this with previous figure. Reduce aspect ratio. Define mathematical variables - $\mathcal{L}$, $\mathcal{L}_{\text{mse}}$**

5 Finite sampling effects for a deterministic target

Now we move on to analyzing the scaling of θ that solves the following problem:

$$\mathcal{L} = \frac{1}{2} \sum_{\mu=1}^P [w(r_\mu) (\hat{\theta}^T \phi(r_\mu) - y_\mu)^2] + \frac{\lambda}{2} \|\hat{\theta}\|^2 \quad (26)$$

What is the meaning of this loss-function, why is there a λ . Cite relevant references, why is it called the ridge loss. Do a toy-example to show how minimization works and put it as a boxes. The loss function in (??) has been modified to accommodate finite number of data points P .

Differentiating the loss function with respect to $\hat{\theta}$ and setting to zero, we arrive at

$$\begin{aligned} \sum_{\mu=1}^P w(r_\mu) \phi(r_\mu) \phi(r_\mu)^T \hat{\theta} - \sum_{\mu=1}^P w(r_\mu) y_\mu \phi(r_\mu) + \lambda \hat{\theta} &= 0 \\ \Rightarrow \left(\frac{1}{P} \sum_{\mu=1}^P w(r_\mu) \phi(r_\mu) \phi(r_\mu)^T + \frac{\lambda}{P} I \right) \hat{\theta} &= \frac{1}{P} \sum_{\mu=1}^P w(r_\mu) y_\mu \phi(r_\mu) \end{aligned}$$

This motivates the definitions

$$\hat{M} = \frac{1}{P} \sum_{\mu=1}^P w(r_\mu) \phi(r_\mu) \phi(r_\mu)^T \quad (27)$$

and

$$\hat{b} = \frac{1}{P} \sum_{\mu=1}^P w(r_\mu) y_\mu \phi(r_\mu) \quad (28)$$

So the empirical estimator can be compactly written as

$$\hat{\theta}(\Delta, K, P) = (\hat{M} + \frac{\lambda}{P} I)^{-1} \hat{b} \quad (29)$$

$$\hat{b} \in \mathbb{R}^{K \times 1}; \quad \hat{M} \in \mathbb{R}^{K \times K}; \quad \hat{\theta} \in \mathbb{R}^{K \times 1} \quad (30)$$

Assume

$$y_\mu = g(r_\mu) \quad (31)$$

Therefore

$$\hat{b} = \frac{1}{P} \sum_{\mu=1}^P w(r_\mu) g(r_\mu) \phi(r_\mu) \quad (32)$$

and

$$\hat{M} = \frac{1}{P} \sum_{\mu=1}^P w(r_\mu) \phi(r_\mu) \phi(r_\mu)^T \quad (33)$$

Then the estimator is

$$\hat{\theta} = (\hat{M} + \frac{\lambda}{P} I)^{-1} \hat{b} \quad (34)$$

$$\Rightarrow \hat{\theta}(\Delta, K, P) = \left(\frac{1}{P} \sum_{\mu=1}^P w(r_\mu) \phi_\mu \phi_\mu^T + \frac{\lambda}{P} I \right)^{-1} \left(\frac{1}{P} \sum_{\mu=1}^P w(r_\mu) y_\mu \phi_\mu \right) \quad (35)$$

Define

$$M(\Delta, K) = \mathbb{E}_{r \sim \rho_\Delta}[\hat{M}] = \mathbb{E}_{r \sim \rho_\Delta}[w(r) \phi(r) \phi(r)^T] \quad (36)$$

$$b(\Delta, K) = \mathbb{E}_{r \sim \rho_\Delta}[\hat{b}] = \mathbb{E}_{r \sim \rho_\Delta}[w(r) g(r) \phi(r)] \quad (37)$$

As $P \rightarrow \infty$

$$\hat{\theta}(\Delta, K, P) \rightarrow \theta_\infty(\Delta, K) = M^{-1} b \quad (38)$$

which corresponds to the continuum solution obtained in the infinite-sample limit (see eqn (??)).

5.1 Perturbation around infinite sample limit

This has to be expanded. Give as much details as possible. We now introduce the finite data effect as a perturbation to the infinite limit case.

$$\hat{M} = M(\Delta, K) + \delta M \quad (39)$$

and

$$\hat{b} = b(\Delta, K) + \delta b \quad (40)$$

where

$$\delta M = \frac{1}{P} \sum_{\mu=1}^P [w(r_\mu) \phi_\mu \phi_\mu^T - M]; \quad \delta b = \frac{1}{P} \sum_{\mu=1}^P [w(r_\mu) g(r_\mu) \phi_\mu - b]$$

Why and how are you making these claims? The mean and variance of δM and δb are as follows:

$$\mathbb{E}[\delta M] = \mathbb{E}[\delta b] = 0 \quad (41)$$

and

$$\mathbb{E}[\delta M \delta M^\top] = O(P^{-1}), \quad \mathbb{E}[\delta b \delta b^\top] = O(P^{-1}) \quad (42)$$

$$\implies \delta M \sim O(P^{-1/2}), \quad \delta b \sim O(P^{-1/2}) \quad (43)$$

Show that fluctuations are of the $O(P^{-1/2})$. Further we define,

$$A = M + \frac{\lambda}{P} I \quad (44)$$

Then, applying Neumann series expansion to first order in $P^{-1/2}$,

$$\hat{\boldsymbol{\theta}}(\Delta, K, P) = (A + \delta M)^{-1}(b + \delta b) = (A^{-1} - A^{-1}\delta M A^{-1} + O(\delta M^2))(b + \delta b) \quad (45)$$

$$\implies \hat{\boldsymbol{\theta}} = A^{-1}b + A^{-1}\delta b - A^{-1}\delta M A^{-1}b + O(P^{-1}) \quad (46)$$

$$\implies \hat{\boldsymbol{\theta}} = \boldsymbol{\theta} + A^{-1}\delta b - A^{-1}\delta M \boldsymbol{\theta} + O(P^{-1}) \quad (47)$$

$$\implies \hat{\boldsymbol{\theta}}(\Delta, K, P) = \boldsymbol{\theta}(\Delta, K) + \frac{1}{\sqrt{P}}\boldsymbol{\eta}(\Delta, K) + O(P^{-1}) \quad (48)$$

where

$$\boldsymbol{\eta}(\Delta, K) = \sqrt{P}A^{-1}(\delta b - \delta M \boldsymbol{\theta}) \quad (49)$$

The first and second moments of $\boldsymbol{\eta}$ evaluate to:

$$\mathbb{E}[\boldsymbol{\eta}] = \sqrt{P}\left(A^{-1}\mathbb{E}[\delta b] - A^{-1}\mathbb{E}[\delta M]\boldsymbol{\theta}\right) = 0 \quad (50)$$

$$\begin{aligned} \mathbb{E}[\boldsymbol{\eta}\boldsymbol{\eta}^\top] &= PA^{-1}\mathbb{E}\left[(\delta b - \delta M \boldsymbol{\theta})(\delta b - \delta M \boldsymbol{\theta})^\top\right](A^{-1})^\top \\ &= PA^{-1}\mathbb{E}\left[\delta b \delta b^\top - \delta b(\delta M \boldsymbol{\theta})^\top - (\delta M \boldsymbol{\theta})\delta b^\top + (\delta M \boldsymbol{\theta})(\delta M \boldsymbol{\theta})^\top\right]A^{-1} \end{aligned} \quad (51)$$

Since each expectation term in the bracket scales as $O(P^{-1})$, the factor of P cancels, leaving:

$$\mathbb{E}[\boldsymbol{\eta}\boldsymbol{\eta}^\top] = \mathcal{C}(\Delta, K) = O(1) \quad (52)$$

where $\mathcal{C}(\Delta, K)$ is the sampling covariance matrix of the estimator fluctuation.

Using these results, the first and second moments of the estimator $\hat{\boldsymbol{\theta}}(\Delta, K, P)$ are:

$$\mathbb{E}[\hat{\boldsymbol{\theta}}] = \boldsymbol{\theta}(\Delta, K) + O(P^{-1}) \quad (53)$$

$$\begin{aligned} \mathbb{E}[\hat{\boldsymbol{\theta}}\hat{\boldsymbol{\theta}}^\top] &= \mathbb{E}\left[\left(\boldsymbol{\theta} + \frac{1}{\sqrt{P}}\boldsymbol{\eta} + O(P^{-1})\right)\left(\boldsymbol{\theta} + \frac{1}{\sqrt{P}}\boldsymbol{\eta} + O(P^{-1})\right)^\top\right] \\ &= \boldsymbol{\theta}\boldsymbol{\theta}^\top + \frac{1}{\sqrt{P}}\left(\mathbb{E}[\boldsymbol{\eta}]\boldsymbol{\theta}^\top + \boldsymbol{\theta}\mathbb{E}[\boldsymbol{\eta}^\top]\right) + \frac{1}{P}\mathbb{E}[\boldsymbol{\eta}\boldsymbol{\eta}^\top] + O(P^{-3/2}) \\ &= \boldsymbol{\theta}\boldsymbol{\theta}^\top + \frac{1}{P}\mathcal{C}(\Delta, K) + O(P^{-3/2}) \\ &= \boldsymbol{\theta}\boldsymbol{\theta}^\top + O(P^{-1}) \quad \text{to leading order in } P. \end{aligned} \quad (54)$$

Now, incorporating the scaling behavior with respect to the gap parameter Δ from Section ??, where $\theta(\Delta) = \theta_0 + \Delta^{1-\nu}\theta_1 + O(\Delta^{3-\nu})$, we obtain the joint finite-sample and critical-gap scaling for the estimator:

$$\hat{\theta}(\Delta, K, P) = \theta_0 + \theta_1 \Delta^{1-\nu} + \frac{1}{\sqrt{P}} \eta(\Delta, K) + O(\Delta^{3-\nu}, P^{-1}). \quad (55)$$

This explicitly separates the deterministic approximation error induced by the critical gap Δ from the $O(P^{-1/2})$ statistical fluctuations caused by finite dataset size P .




Figure 4: **Validation of Central Limit Theorem scaling and gap-dependent fluctuation amplification for finite sample size P .** (a) Variance of the estimated parameter $\text{Var}[\hat{\theta}_0]$ versus dataset size P over $S = 250$ independent data realizations. The empirical slope $\alpha = -1.03$ confirms the standard $O(P^{-1})$ CLT decay of estimator variance (Eq.??). (b) Scaling of the empirical covariance matrix entry $\mathcal{C}_{00}(\Delta)$ with critical gap Δ . The power-law scaling $\mathcal{C}_{00}(\Delta) \sim \Delta^{-0.49}$ highlights the noise amplification driven by the ill-conditioning of the empirical design matrix $\hat{M}$ as $\Delta \rightarrow 0$.

6 Finite Sampling with noisy target

$$r_\mu \stackrel{i.i.d}{\sim} \rho_\Delta,$$

$$y = g(r_\mu) + \xi_\mu; \quad \xi \sim \mathcal{N}(0, \tau^2),$$

and

$$\phi(r) = (1, r, r^2, \dots, r^{K-1})^T$$