

## Analgesics

<b>Acetaminophen 500 mg tabs</b> Sig: Take 1-2 tabs PO q4-6h prn pain, max 4000mg/day Disp: 30 tabs	<b>Ibuprofen 200 mg tabs</b> Sig: Take 1-2 tabs PO q4-6h prn pain, max 2400mg/day Disp: 30 tabs
<b>Ketorolac 10 mg tabs</b> Sig: Take 1 tab PO q6h prn pain, max 40mg/day Disp: 20 tabs	<b>Acetaminophen 300mg, codeine 30mg, caffeine 15mg tabs</b> Sig: Take 1-2 tabs PO q4-6h prn pain, max 4000mg/day acet Disp: 20 tabs
<b>Acetaminophen 325mg, oxycodone 5mg tabs</b> Sig: Take 1 tab PO q6h prn pain, max 4000mg/day acet Disp: 20 tabs	<b>Acetylsalicylic acid 325mg, oxycodone 5mg tabs</b> Sig: Take 1 tab PO q6h prn pain, max 4000mg/day acet Disp: 20 tabs
<b>Acetaminophen 80mg/mL suspension</b> Sig: Shake well and give 10 mg/kg q4-6h prn pain, max 65mg/kg/day Disp: based on weight	<b>Ibuprofen 20mg/mL suspension</b> Sig: Shake well and give 10mg/kg q6-8h prn pain, max 40mg/kg/day Disp: based on weight

## Antibiotic therapy

<b>Penicillin VK 300 mg tabs</b> Sig: Take 1 tab PO QID for 7 days Disp: 28 tabs	<b>Clindamycin 300 mg caps</b> Sig: Take 1 cap PO QID for 7 days Disp: 28 caps
<b>Amoxicillin 500 mg caps</b> Sig: Take 1 cap PO TID for 7 days Disp: 21 caps	<b>Metronidazole 500 mg tabs</b> Sig: Take 1 tab PO TID for 7 days Disp: 21 tabs
<b>Cephalexin 500 mg tabs</b> Sig: Take 1 tab PO QID for 7 days Disp: 28 tabs	<b>Clarithromycin 500 mg tabs</b> Sig: Take 1 tab PO BID for 7 days Disp: 14 tabs
<b>Amoxicillin 50mg/mL suspension</b> Sig: Shake well and give 25mg/kg TID for 7 days Disp: based on weight	<b>Clindamycin 5mg/mL solution</b> Sig: Give 5mg/kg QID for 7 days Disp: based on weight

## Antibiotic prophylaxis

<b>Amoxicillin 500 mg caps</b> Sig: Take 4 caps 1 hour before procedure Disp: 4 caps	<b>Clindamycin 300 mg caps</b> Sig: Take 2 caps 1 hour before procedure Disp: 2 caps
<b>Cephalexin 500 mg tabs</b> Sig: Take 4 tabs 1 hour before procedure Disp: 4 tabs	
<b>Amoxicillin 50mg/mL suspension</b> Sig: Shake well and give 50mg/kg 1 hour before procedure Disp: based on weight	<b>Clindamycin 5mg/mL solution</b> Sig: Give 20mg/kg 1 hour before procedure Disp: based on weight

## Rinses

<b>Nystatin 100,000 units/mL suspension</b> Sig: Shake well, swish and swallow 5mL QID for 14 days Disp: 300mL	<b>Chlorhexidine 0.12% solution</b> Sig: Swish and spit 15mL BID until finished Disp: 475mL
--	---

## Creams

<b>Acyclovir 5% cream</b> Sig: Apply to affected area QID for 10 days Disp: 30g	<b>Triamcinolone 0.1% cream</b> Sig: Apply to affected area TID Disp: 30g
<b>Nystatin 100,000 units/g cream</b> Sig: Apply to affected area TID Disp: 30g	