

# Lecture 2: Geometry of Linear Equations in 2 and 3 unknowns

M. Krishna Kumar

Principal Research Scientist (Retd.)

DESE, IISc, Bangalore

May 31, 2024

# Recap

- Matrix Representation of system of linear equations

# Recap

- Matrix Representation of system of linear equations
- Some interpretations of matrices in different application domains

# Today's agenda

- Geometry of linear equations involving 2 and 3 unknowns

# Today's agenda

- Geometry of linear equations involving 2 and 3 unknowns
- Solutions to a system of linear equations

# Today's agenda

- Geometry of linear equations involving 2 and 3 unknowns
- Solutions to a system of linear equations
- Existence of solution to system of linear equations

## Example 1

Consider the system of equations

$$l_1 : 2x_1 + x_2 = 3$$

$$l_2 : x_1 + 2x_2 = 3$$

- $l_1$  and  $l_2$  represent two straight lines


## Example 1

Consider the system of equations

$$l_1 : 2x_1 + x_2 = 3$$

$$l_2 : x_1 + 2x_2 = 3$$

- $l_1$  and  $l_2$  represent two straight lines



## Some observations:

- Lines  $l_1$  and  $l_2$  intersect at  $(x_1, x_2) = (1, 1)$ .

## Some observations:

- Lines  $l_1$  and  $l_2$  intersect at  $(x_1, x_2) = (1, 1)$ .
- $(x_1, x_2) = (1, 1)$  -THE solution we are looking for

## Some observations:

- Lines  $l_1$  and  $l_2$  intersect at  $(x_1, x_2) = (1, 1)$ .
- $(x_1, x_2) = (1, 1)$  -THE solution we are looking for
- To solve for  $(x_1, x_2)$  implies find the point(s) of intersection of the two lines,  $l_1$  and  $l_2$ , if any

## Some observations:

- Lines  $l_1$  and  $l_2$  intersect at  $(x_1, x_2) = (1, 1)$ .
- $(x_1, x_2) = (1, 1)$  - **THE** solution we are looking for
- To solve for  $(x_1, x_2)$  implies find the point(s) of intersection of the two lines,  $l_1$  and  $l_2$ , **if any**
- Intersection point - A point that lies on both the lines - Solution to a pair of linear equations in 2 unknowns

## Some observations:

- Lines  $l_1$  and  $l_2$  intersect at  $(x_1, x_2) = (1, 1)$ .
- $(x_1, x_2) = (1, 1)$  - **THE** solution we are looking for
- To solve for  $(x_1, x_2)$  implies find the point(s) of intersection of the two lines,  $l_1$  and  $l_2$ , **if any**
- Intersection point - A point that lies on both the lines - Solution to a pair of linear equations in 2 unknowns
- Only one point of intersection - **UNIQUE solution.**

## Example 2

- 

$$l_1 : 2x_1 + x_2 = 3$$


$$l_2 : 4x_1 + 2x_2 = 6$$

## Example 2

- 

$$l_1 : 2x_1 + x_2 = 3$$

$$l_2 : 4x_1 + 2x_2 = 6$$



## Example 2 - Some Observations

- Two lines overlap

## Example 2 - Some Observations

- Two lines overlap
- $l_1$  and  $l_2$  - intersect at more than one point, in fact, at infinitely many points.

## Example 2 - Some Observations

- Two lines overlap
- $l_1$  and  $l_2$  - intersect at more than one point, in fact, at infinitely many points.
- Every point on  $l_1$  is also there on  $l_2$

## Example 2 - Some Observations

- Two lines overlap
- $l_1$  and  $l_2$  - intersect at more than one point, in fact, at infinitely many points.
- Every point on  $l_1$  is also there on  $l_2$
- Infinitely many solutions

## Example 3

- 

$$l_1 : 2x_1 + x_2 = 3$$


$$l_2 : 2x_1 + x_2 = 8$$

## Example 3

- 

$$l_1 : 2x_1 + x_2 = 3$$

$$l_2 : 2x_1 + x_2 = 8$$



## Example 3 - Some Observations

- $l_1, l_2$  do not intersect at all

## Example 3 - Some Observations

- $l_1, l_2$  do not intersect at all
- We cannot find any point which lies on both the lines simultaneously

## Example 3 - Some Observations

- $l_1, l_2$  do not intersect at all
- We cannot find any point which lies on both the lines simultaneously
- **NO SOLUTION**

# Some Questions to look answers for!

1. When does a system of linear equations have solution?
2. When does a system of linear equations not have solution?
3. If the system of linear equations has solution, how many solutions does it have?

# Consistent & Inconsistent Systems

- **Consistent System** - A system of linear equations is said to be **consistent** if it has one or many solutions

# Consistent & Inconsistent Systems

- **Consistent System** - A system of linear equations is said to be **consistent** if it has one or many solutions
- Example:

$$l_1 : 2x_1 + x_2 = 3$$

$$l_2 : x_1 + 2x_2 = 3$$

# Consistent & Inconsistent Systems

- **Consistent System** - A system of linear equations is said to be **consistent** if it has one or many solutions
- Example:

$$l_1 : 2x_1 + x_2 = 3$$

$$l_2 : x_1 + 2x_2 = 3$$

- Example:

$$l_1 : 2x_1 + x_2 = 3$$

$$l_2 : 4x_1 + 2x_2 = 6$$

# Consistent & Inconsistent Systems

- **Consistent System** - A system of linear equations is said to be **consistent** if it has one or many solutions
- Example:

$$l_1 : 2x_1 + x_2 = 3$$

$$l_2 : x_1 + 2x_2 = 3$$

- Example:

$$l_1 : 2x_1 + x_2 = 3$$

$$l_2 : 4x_1 + 2x_2 = 6$$

- **Inconsistent System** - A system of linear equations is said to be **inconsistent** if it has no solutions.

# Consistent & Inconsistent Systems

- **Consistent System** - A system of linear equations is said to be **consistent** if it has one or many solutions
- Example:

$$l_1 : 2x_1 + x_2 = 3$$

$$l_2 : x_1 + 2x_2 = 3$$

- Example:

$$l_1 : 2x_1 + x_2 = 3$$

$$l_2 : 4x_1 + 2x_2 = 6$$

- **Inconsistent System** - A system of linear equations is said to be **inconsistent** if it has no solutions.

- 

$$l_1 : 2x_1 + x_2 = 3$$

$$l_2 : 2x_1 + x_2 = 8$$

## Some Examples of Consistent Systems - Unique solution case

# Some Examples of Consistent Systems - Unique solution case




Figure: Unique Solution Scenarios

## Some Examples of Consistent Systems - Infinitely many solutions case

# Some Examples of Consistent Systems - Infinitely many solutions case




Figure: Infinitely Many Solutions Scenarios

## Some Examples of Inconsistent Systems

- No solution exists when the lines are parallel or different pairs of lines intersect at different points.

# Some Examples of Inconsistent Systems

- No solution exists when the lines are parallel or different pairs of lines intersect at different points.





Figure: Inconsistent - No Solution Scenarios

# Some Examples of Inconsistent Systems

- No solution exists when the lines are parallel or different pairs of lines intersect at different points.




Figure: Inconsistent - No Solution Scenarios

- Measurement error**

$$l_1 : 2x_1 + x_2 = 3$$

$$l_2 : 2x_1 + x_2 = 8$$

# Some Examples of Inconsistent Systems

- No solution exists when the lines are parallel or different pairs of lines intersect at different points.




Figure: Inconsistent - No Solution Scenarios

- Measurement error**

$$l_1 : 2x_1 + x_2 = 3$$

$$l_2 : 2x_1 + x_2 = 8$$

- Same LHS in both equations, RHS are different - Inconsistency

System of linear equations in 3 unknowns

## Example 4

- 

$$2x + 3y + z = 6$$

$$x - 2y + 3z = 2$$

$$x = 1$$


## Example 4

$$2x + 3y + z = 6$$

$$x - 2y + 3z = 2$$

$$x = 1$$

$$x=1, y=1, z=1$$



## Example 4 - Some Observations

- Point of intersection  $(x, y, z) = (1, 1, 1)$

## Example 4 - Some Observations

- Point of intersection  $(x, y, z) = (1, 1, 1)$
- That is **THE** solution

## Example 5

- 

$$2x + 3y + z = 6$$


$$x - 2y + 3z = 2$$

$$3x + y + 4z = 8$$

## Example 5

## Example 5

*Line of intersection of the 3 planes*



## Example 5 - Some observations

- Intersection of planes in this case - **Line**

## Example 5 - Some observations

- Intersection of planes in this case - [Line](#)
- The parametric form of the line is given by  $(x, y, z) = \left(-\frac{11}{7}t + \frac{18}{7}, \frac{5t}{7} + \frac{2}{7}, t\right)$

## Example 5 - Some observations

- Intersection of planes in this case - **Line**
- The parametric form of the line is given by  $(x, y, z) = \left(-\frac{11}{7}t + \frac{18}{7}, \frac{5t}{7} + \frac{2}{7}, t\right)$
- **Look at where a wall and the adjacent floor meet**

## Example 6

•

$$2x + y + z = -15$$


$$2x + y + z = 20$$

$$2x + y + z = 70$$


(1)

## Some observations about Example 6

## Some observations about Example 6




## Some observations about Example 6




Planes do not intersect at all - **NO SOLUTION**


# Visualizing consistent systems in 3D




# Consistent System with infinitely many solutions



Infinitely many solutions  
(intersection is a line)



Infinitely many solutions  
(planes are all coincident;  
intersection is a plane)




Infinitely many solutions  
(two coincident planes;  
intersection is a line)


Figure: Infinitely Many solutions - Planes intersect in a line<sup>1</sup>

<sup>1</sup>Image Courtesy: Anton, Rorres, Kaul, *Elementary Linear Algebra*, Wiley, 2019, 12th Ed.


# Inconsistent Systems




No solutions  
(three parallel planes;  
no common intersection)



No solutions  
(two parallel planes;  
no common intersection)



No solutions  
(no common intersection)




No solutions  
(two coincident planes  
parallel to the third;  
no common intersection)

Figure: No solutions<sup>2</sup>

<sup>2</sup>Image Courtesy: Anton, Rorres, Kaul, *Elementary Linear Algebra*, Wiley, 2019, 12th Ed.

# Inconsistent Systems



## Some Comments

For a system of linear equations, be it in two equations in two unknowns or three equations in three unknowns, **there can only be**

- unique or exactly one solution

## Some Comments

For a system of linear equations, be it in two equations in two unknowns or three equations in three unknowns, **there can only be**

- unique or exactly one solution
- infinitely many solutions or

## Some Comments

For a system of linear equations, be it in two equations in two unknowns or three equations in three unknowns, **there can only be**

- unique or exactly one solution
- infinitely many solutions or
- no solution at all

# Summary

- Systems of linear equations in 2 unknowns and in 3 unknowns

# Summary

- Systems of linear equations in 2 unknowns and in 3 unknowns
- Geometry of system of equations in two and three unknowns

# Summary

- Systems of linear equations in 2 unknowns and in 3 unknowns
- Geometry of system of equations in two and three unknowns
- We had two important questions, one about the existence of solution to the system of equations and the other about the number of solutions

# Summary

- Systems of linear equations in 2 unknowns and in 3 unknowns
- Geometry of system of equations in two and three unknowns
- We had two important questions, one about the existence of solution to the system of equations and the other about the number of solutions
- Consistent and Inconsistent system of equations

# Summary

- Systems of linear equations in 2 unknowns and in 3 unknowns
- Geometry of system of equations in two and three unknowns
- We had two important questions, one about the existence of solution to the system of equations and the other about the number of solutions
- Consistent and Inconsistent system of equations
- Any system that is consistent has solution, either unique or infinitely many

# Summary

- Systems of linear equations in 2 unknowns and in 3 unknowns
- Geometry of system of equations in two and three unknowns
- We had two important questions, one about the existence of solution to the system of equations and the other about the number of solutions
- Consistent and Inconsistent system of equations
- Any system that is consistent has solution, either unique or infinitely many
- An inconsistent system does not have solution

# Summary

- Systems of linear equations in 2 unknowns and in 3 unknowns
- Geometry of system of equations in two and three unknowns
- We had two important questions, one about the existence of solution to the system of equations and the other about the number of solutions
- Consistent and Inconsistent system of equations
- Any system that is consistent has solution, either unique or infinitely many
- An inconsistent system does not have solution
- **Next Lecture - General system of linear equations, their solutions**

**Thank you!, Namaste**

# Lecture 3: General Linear System of Equations and Matrices

M. Krishna Kumar

Principal Research Scientist (Retd.)

DESE, IISc, Bangalore

# Recap

- Geometry of system of linear equations in 2 and 3 unknowns
- Focus of this lecture - General system of linear equations followed by some matrix definitions

# General System of Linear Equations

- Consider the system of linear equations,

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2$$

$$\vdots \quad \vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m$$

# General System of Linear Equations

- Consider the system of linear equations,

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2$$

$$\vdots \quad \vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m$$

- Matrix Representation:  $A^{m \times n} \mathbf{x}^{n \times 1} = \mathbf{b}^{m \times 1}$

# General System of Linear Equations

- Consider the system of linear equations,

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2$$

$$\vdots \quad \vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m$$

- Matrix Representation:  $A^{m \times n} \mathbf{x}^{n \times 1} = \mathbf{b}^{m \times 1}$
- Double subscript - with the coefficients of the unknowns in all the equations

# General System of Linear Equations

- Consider the system of linear equations,

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2$$

$$\vdots \quad \vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m$$

- Matrix Representation:  $A^{m \times n} \mathbf{x}^{n \times 1} = \mathbf{b}^{m \times 1}$
- Double subscript - with the coefficients of the unknowns in all the equations
- First subscript - equation in which the coefficient occurs,

# General System of Linear Equations

- Consider the system of linear equations,

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2$$

$$\vdots \quad \vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m$$

- Matrix Representation:  $A^{m \times n} \mathbf{x}^{n \times 1} = \mathbf{b}^{m \times 1}$
- Double subscript - with the coefficients of the unknowns in all the equations
- First subscript - equation in which the coefficient occurs,
- Second subscript - unknown that it multiplies

# General System of Linear Equations

- Consider the system of linear equations,

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2$$

$$\vdots \quad \vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m$$

- Matrix Representation:  $A^{m \times n} \mathbf{x}^{n \times 1} = \mathbf{b}^{m \times 1}$
- Double subscript - with the coefficients of the unknowns in all the equations
- First subscript - equation in which the coefficient occurs,
- Second subscript - unknown that it multiplies
- For example:  $a_{34}$  is a coefficient in the third equation and multiplies  $x_4$ .

## Solution to the System of linear equations

- **Particular solution** to this linear system of equations in  $n$  unknowns - Sequence of  $n$  numbers, say  $x_1 = s_1, x_2 = s_2, \dots, x_n = s_n$  that satisfies each and every equation that constitutes the system of equations  $A\mathbf{x} = \mathbf{b}$ .

# Solution to the System of linear equations

- **Particular solution** to this linear system of equations in  $n$  unknowns - Sequence of  $n$  numbers, say  $x_1 = s_1, x_2 = s_2, \dots, x_n = s_n$  that satisfies each and every equation that constitutes the system of equations  $A\mathbf{x} = \mathbf{b}$ .
- Solutions - **n-tuple**

# Solution to the System of linear equations

- **Particular solution** to this linear system of equations in  $n$  unknowns - Sequence of  $n$  numbers, say  $x_1 = s_1, x_2 = s_2, \dots, x_n = s_n$  that satisfies each and every equation that constitutes the system of equations  $A\mathbf{x} = \mathbf{b}$ .
- Solutions - **n-tuple**
- **Complete solution** set for a system of linear equations in  $n$  variables - Set of all  $n$ -tuples that form particular solutions to the system

# Solution to the System of linear equations

- **Particular solution** to this linear system of equations in  $n$  unknowns - Sequence of  $n$  numbers, say  $x_1 = s_1, x_2 = s_2, \dots, x_n = s_n$  that satisfies each and every equation that constitutes the system of equations  $Ax = b$ .
- Solutions -  **$n$ -tuple**
- **Complete solution** set for a system of linear equations in  $n$  variables - Set of all  $n$ -tuples that form particular solutions to the system
- **Question:** Should it always be that the number of equations and the number of unknowns be the same?

# Solution to the System of linear equations

- **Particular solution** to this linear system of equations in  $n$  unknowns - Sequence of  $n$  numbers, say  $x_1 = s_1, x_2 = s_2, \dots, x_n = s_n$  that satisfies each and every equation that constitutes the system of equations  $Ax = b$ .
- Solutions -  **$n$ -tuple**
- **Complete solution** set for a system of linear equations in  $n$  variables - Set of all  $n$ -tuples that form particular solutions to the system
- **Question:** Should it always be that the number of equations and the number of unknowns be the same?
- The answer is **NEED NOT BE**.

## Example 1

Consider the following system of equations:

$$\begin{aligned} 2x_1 + x_2 &= 3 \\ x_1 + 2x_2 &= 3 \\ x_1 + 3x_2 &= 4 \\ 4x_1 + 7x_2 &= 11 \end{aligned} \tag{1}$$

Observe the following

- 4 linear equations involving only 2 unknowns

## Example 1

Consider the following system of equations:

$$\begin{aligned} 2x_1 + x_2 &= 3 \\ x_1 + 2x_2 &= 3 \\ x_1 + 3x_2 &= 4 \\ 4x_1 + 7x_2 &= 11 \end{aligned} \tag{1}$$

Observe the following

- 4 linear equations involving only 2 unknowns
- The number of equations is more than the number of unknowns, i.e.,  $m > n$

## Example 1

Consider the following system of equations:

$$\begin{aligned} 2x_1 + x_2 &= 3 \\ x_1 + 2x_2 &= 3 \\ x_1 + 3x_2 &= 4 \\ 4x_1 + 7x_2 &= 11 \end{aligned} \tag{1}$$

Observe the following

- 4 linear equations involving only 2 unknowns
- The number of equations is more than the number of unknowns, i.e.,  $m > n$
- Tall matrix - **overdetermined** system

# Overdetermined System

- More constraints than the unknown
- May or may not have solution to the system of equations

# Geometry of overdetermined system with unique solution

# Geometry of overdetermined system with unique solution

Example 1:

$$2x_1 + x_2 = 3$$

$$x_1 + 2x_2 = 3$$

$$x_1 + 3x_2 = 4$$

$$4x_1 + 7x_2 = 11$$

# Geometry of overdetermined system with unique solution

Example 1:

$$2x_1 + x_2 = 3$$

$$x_1 + 2x_2 = 3$$

$$x_1 + 3x_2 = 4$$

$$4x_1 + 7x_2 = 11$$




Figure: OverDetermined System with Unique solution. Solution is  $x_1 = 1, x_2 = 1$ .

# Infinitely Many Solutions

- Will an overdetermined system always have a unique solution?

# Infinitely Many Solutions

- Will an overdetermined system always have a unique solution?
- Number of constraints are more than the unknown, it is natural to expect no solution

# Infinitely Many Solutions

- Will an overdetermined system always have a unique solution?
- Number of constraints are more than the unknown, it is natural to expect no solution
- Example:

# Infinitely Many Solutions

- Will an overdetermined system always have a unique solution?
- Number of constraints are more than the unknown, it is natural to expect no solution
- Example:

$$2x_1 + x_2 = 3$$

$$4x_1 + 2x_2 = 6$$

$$3x_1 + 1.5x_2 = 4.5$$

$$8x_1 + 4x_2 = 12$$

# Infinitely Many Solutions

# Infinitely Many Solutions

Example 2

$$2x_1 + x_2 = 3$$

$$4x_1 + 2x_2 = 6$$

$$3x_1 + 1.5x_2 = 4.5$$

$$8x_1 + 4x_2 = 12$$

# Infinitely Many Solutions


Example 2

$$2x_1 + x_2 = 3$$

$$4x_1 + 2x_2 = 6$$

$$3x_1 + 1.5x_2 = 4.5$$

$$8x_1 + 4x_2 = 12$$



# Infinitely Many Solutions

Example 2

$$2x_1 + x_2 = 3$$

$$4x_1 + 2x_2 = 6$$

$$3x_1 + 1.5x_2 = 4.5$$

$$8x_1 + 4x_2 = 12$$




Figure: OverDetermined System with Infinitely many solutions.

# No Solutions

- As a next step, we look at system which has constraints that are impossible to be met simultaneously

$$2x_1 + x_2 = 3$$

$$x_1 + 2x_2 = 3$$

$$x_1 + 3x_2 = 5$$

$$4x_1 + 7x_2 = 4$$

## No Solution case

## No Solution case

Overdetermined system with  
no solution

$$2x_1 + x_2 = 3$$

$$x_1 + 2x_2 = 3$$

$$x_1 + 3x_2 = 5$$

$$4x_1 + 7x_2 = 4$$

# No Solution case


Overdetermined system with no solution

$$2x_1 + x_2 = 3$$

$$x_1 + 2x_2 = 3$$

$$x_1 + 3x_2 = 5$$

$$4x_1 + 7x_2 = 4$$



# No Solution case

Overdetermined system with no solution

$$2x_1 + x_2 = 3$$

$$x_1 + 2x_2 = 3$$

$$x_1 + 3x_2 = 5$$

$$4x_1 + 7x_2 = 4$$




Figure: OverDetermined System with No solutions. Every pair of lines intersect at different points. Together, they do not intersect at any common point.

# Overdetermined Systems

An over determined system ( $m > n$ ) can have

- unique solution or

# Overdetermined Systems

An over determined system ( $m > n$ ) can have

- unique solution or
- infinitely many solutions or

# Overdetermined Systems

An over determined system ( $m > n$ ) can have

- unique solution or
- infinitely many solutions or
- no solution - (Inconsistent case)

# Overdetermined Systems

An over determined system ( $m > n$ ) can have

- unique solution or
- infinitely many solutions or
- no solution - (Inconsistent case)
- Later, from the perspective of vector spaces, what it means to say solution (unique or infinitely many) exists and no solution exists!

# Overdetermined Systems

An over determined system ( $m > n$ ) can have

- unique solution or
- infinitely many solutions or
- no solution - (Inconsistent case)
- Later, from the perspective of vector spaces, what it means to say solution (unique or infinitely many) exists and no solution exists!
- Question: What happens when we have fewer equations than the number of variables?

# Underdetermined System

$$2x_1 + 3x_2 + 4x_3 = 0$$

$$x_1 + 2x_2 + x_3 = 3$$

- 3 unknowns and only 2 equations

# Underdetermined System

$$2x_1 + 3x_2 + 4x_3 = 0$$

$$x_1 + 2x_2 + x_3 = 3$$

- 3 unknowns and only 2 equations
- Represents a system with too few observations for too many variables

# Underdetermined System

$$\begin{aligned}2x_1 + 3x_2 + 4x_3 &= 0 \\x_1 + 2x_2 + x_3 &= 3\end{aligned}$$

- 3 unknowns and only 2 equations
- Represents a system with too few observations for too many variables
- Suppose  $n$  denotes the number of unknowns, and  $m$  the number of equations, we see here that  $m < n$ .

# Underdetermined System

$$\begin{aligned}2x_1 + 3x_2 + 4x_3 &= 0 \\x_1 + 2x_2 + x_3 &= 3\end{aligned}$$

- 3 unknowns and only 2 equations
- Represents a system with too few observations for too many variables
- Suppose  $n$  denotes the number of unknowns, and  $m$  the number of equations, we see here that  $m < n$ .
- System of equations with more unknowns and fewer equations - underdetermined system

# Underdetermined System

$$\begin{aligned}2x_1 + 3x_2 + 4x_3 &= 0 \\x_1 + 2x_2 + x_3 &= 3\end{aligned}$$

- 3 unknowns and only 2 equations
- Represents a system with too few observations for too many variables
- Suppose  $n$  denotes the number of unknowns, and  $m$  the number of equations, we see here that  $m < n$ .
- **System of equations with more unknowns and fewer equations - underdetermined system**
- What can we say about the solution to the system of equations of an underdetermined system?

# Underdetermined System with infinitely many solutions

$$2x_1 + 3x_2 + 4x_3 = 0$$

$$x_1 + 2x_2 + x_3 = 3$$

- The equations correspond to two planes

# Underdetermined System with infinitely many solutions

$$2x_1 + 3x_2 + 4x_3 = 0$$

$$x_1 + 2x_2 + x_3 = 3$$

- The equations correspond to two planes
- Two planes either intersect in a line or do not intersect at all

# Underdetermined System with infinitely many solutions

# Underdetermined System with infinitely many solutions


$$2x_1 + 3x_2 + 4x_3 = 0$$

$$x_1 + 2x_2 + x_3 = 3$$

# Underdetermined System with infinitely many solutions

$$2x_1 + 3x_2 + 4x_3 = 0$$

$$x_1 + 2x_2 + x_3 = 3$$



# Underdetermined System with infinitely many solutions

$$\begin{aligned}2x_1 + 3x_2 + 4x_3 &= 0 \\x_1 + 2x_2 + x_3 &= 3\end{aligned}$$




Figure: Underdetermined System with Infinitely many solutions. The two planes intersect in a line

## Example 5

Underdetermined system - inconsistent or infinitely many solutions. Never a unique solution

## Example 5

Here is an example of an underdetermined system with no solution

$$2x_1 + 3x_2 + 4x_3 = 0$$

$$2x_1 + 3x_2 + 4x_3 = 3$$


Underdetermined system - inconsistent or infinitely many solutions. Never a unique solution

## Example 5

Here is an example of an underdetermined system with no solution

$$2x_1 + 3x_2 + 4x_3 = 0$$

$$2x_1 + 3x_2 + 4x_3 = 3$$



Underdetermined system - inconsistent or infinitely many solutions. Never a unique solution

## Example 5

Here is an example of an underdetermined system with no solution

$$2x_1 + 3x_2 + 4x_3 = 0$$

$$2x_1 + 3x_2 + 4x_3 = 3$$




Figure: Underdetermined System with No solutions. The two planes are parallel

Underdetermined system - inconsistent or infinitely many solutions. Never a unique solution

## Matrices

# Matrices

- A typical matrix is defined as a rectangular array of numbers arranged in  $m$  rows and  $n$  columns

# Matrices

- A typical matrix is defined as a rectangular array of numbers arranged in  $m$  rows and  $n$  columns

$$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}$$

# Matrices

- A typical matrix is defined as a rectangular array of numbers arranged in  $m$  rows and  $n$  columns

$$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}$$

- $A$  :  $m$  by  $n$  real matrix if all its entries are real numbers

## Some examples of matrices - You all know it, still

- $m < n$  - wide matrix. Example:

$$A = \begin{pmatrix} 1 & 0 & 2 & 5.38 \\ 2.1987 & 3 & -1 & 4 \end{pmatrix} \text{ 2 by 4 matrix .}$$

## Some examples of matrices - You all know it, still

- $m < n$  - wide matrix. Example:

$$A = \begin{pmatrix} 1 & 0 & 2 & 5.38 \\ 2.1987 & 3 & -1 & 4 \end{pmatrix} \text{ 2 by 4 matrix .}$$

- $m > n$  - tall matrix. Example

$$B = \begin{pmatrix} 1 & 0 \\ 2 & 5.38 \\ 2.1987 & \sqrt{2} \\ 3 & -1 \\ 4 & 7 \end{pmatrix} \text{ 5 by 2 matrix - Tall Matrix.}$$

## Some examples of matrices - You all know it, still

- $m < n$  - wide matrix. Example:

$$A = \begin{pmatrix} 1 & 0 & 2 & 5.38 \\ 2.1987 & 3 & -1 & 4 \end{pmatrix} \text{ 2 by 4 matrix .}$$

- $m > n$  - tall matrix. Example

$$B = \begin{pmatrix} 1 & 0 \\ 2 & 5.38 \\ 2.1987 & \sqrt{2} \\ 3 & -1 \\ 4 & 7 \end{pmatrix} \text{ 5 by 2 matrix - Tall Matrix.}$$

- $m = n$  - square matrix. Example:

$$C = \begin{pmatrix} 1 & 0 & 2 \\ 2.1987 & \sqrt{2} & 7 \\ 3 & -17 & 0 \end{pmatrix} \text{ 3 by 3 matrix.}$$

- A Matrix with only one row is called a row vector or a row matrix. Example:

$$D = \begin{pmatrix} 1 & 0 & 2 \end{pmatrix} \quad 1 \text{ by } 3 \text{ matrix.}$$

- Any matrix - collection of  $m$  row vectors
- A matrix with only one column is called a column vector or a column matrix. Example:

$$E = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} \quad 3 \text{ by } 1 \text{ matrix}$$

- Any matrix - collection of  $n$  column vectors
- The 3-by-3 matrix  $I = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$  is called an 3 by 3 IDENTITY matrix An  $n \times n$  identity matrix is denoted by  $\mathcal{I}_{n \times n}$  or simply  $\mathcal{I}$

# Diagonal matrix

- A diagonal matrix is an  $n$  by  $n$  matrix with entries above and below the main diagonal are 0, that is a matrix is a diagonal matrix if and only if it is square and  $d_{i,j} = 0$ , for  $i \neq j$ .

Example:  $F = \begin{pmatrix} 4 & 0 & 0 & 0 \\ 0 & 7.2 & 0 & 0 \\ 0 & 0 & 0.123 & 0 \\ 0 & 0 & 0 & \pi \end{pmatrix}$

- Two matrices  $A$  and  $B$  are said to be **EQUAL** if and only if they have the same size and all of their corresponding entries are the same.  $A = \begin{pmatrix} 1 & 0 & 2 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}$ , Not equal matrices though their entries are the same

# Triangular Matrices

- A matrix is said to be upper triangular  $U$ , if it is a square matrix with all entries below the main diagonal equal to 0. An upper triangular matrix has its entries equal to zero at

all those positions  $a_{i,j}$  where  $i > j$ . Example:  $U = \begin{pmatrix} 1 & 3 & 2 \\ 0 & 2 & -1 \\ 0 & 0 & \pi \end{pmatrix}$

- A matrix is said to be lower triangular  $L$ , if it is a square matrix with all entries above the main diagonal equal to 0. A lower triangular matrix has its entries equal to zero at all those positions  $a_{i,j}$  where  $i < j$ .

Example:  $L = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 2 & 0 \\ \pi & -5 & 23 \end{pmatrix}$

- A matrix with all its entries is called a zero matrix. A zero matrix can be square or rectangular.

# Summary

- In this lecture, we looked at general system of linear equations.
- We defined what an overdetermined system is. These are systems that have more equations than the unknowns.
- We also looked at underdetermined system as the system with fewer equations than unknowns.
- We looked at the visualizations for such systems
- We defined various matrices.
- In the next lecture, we will look at operations on matrices.

Namaste!!!