

Warm Up Exercise

Solve the system:

$$\begin{aligned}x - 2z &= 0 \\x + 4y - 4z - w &= 0 \\x + 2y - 2w &= 0 \\y - z &= 0\end{aligned}$$

by

- ▶ Writing down the appropriate augmented matrix.
- ▶ Reducing the augmented matrix by Gaussian elimination to RREF.
- ▶ Identifying the solution set.

Warm Up Exercise

- ▶ Augmented matrix:

$$\begin{array}{rcrcrcrcrcrcl} x & - & 2z & = & 0 & & & & & \\ x & + & 4y & - & 4z & - & w & = & 0 & \\ & x & + & 2y & - & 2w & = & 0 & & \\ & & y & - & z & = & 0 & & & \end{array} \Rightarrow \left[\begin{array}{cccc|c} 1 & 0 & -2 & 0 & 0 \\ 1 & 4 & -4 & -1 & 0 \\ 1 & 2 & 0 & -2 & 0 \\ 0 & 1 & -1 & 0 & 0 \end{array} \right]$$

Warm Up Exercise

- ▶ Augmented matrix:

$$\begin{array}{rcrcrcrcrcrcl} x & - & 2z & = & 0 & & & & & \\ x & + & 4y & - & 4z & - & w & = & 0 & \\ x & + & 2y & - & 2w & = & 0 & & & \\ y & - & z & = & 0 & & & & & \end{array} \Rightarrow \left[\begin{array}{cccc|c} 1 & 0 & -2 & 0 & 0 \\ 1 & 4 & -4 & -1 & 0 \\ 1 & 2 & 0 & -2 & 0 \\ 0 & 1 & -1 & 0 & 0 \end{array} \right]$$

- ▶ RREF:

$$\left[\begin{array}{cccc|c} 1 & 0 & -2 & 0 & 0 \\ 1 & 4 & -4 & -1 & 0 \\ 1 & 2 & 0 & -2 & 0 \\ 0 & 1 & -1 & 0 & 0 \end{array} \right] \sim \left[\begin{array}{cccc|c} 1 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1/2 & 0 \\ 0 & 0 & 1 & -1/2 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

Warm Up Exercise

► Sln set:

$$\left[\begin{array}{cccc|c} 1 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1/2 & 0 \\ 0 & 0 & 1 & -1/2 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \Rightarrow \begin{array}{rcl} x - w & = & 0 \\ y - w/2 & = & 0 \\ z - w/2 & = & 0 \\ 0 & = & 0 \end{array}$$

Warm Up Exercise

► Sln set:

$$\left[\begin{array}{cccc|c} 1 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1/2 & 0 \\ 0 & 0 & 1 & -1/2 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \Rightarrow \begin{array}{rcl} x - w & = & 0 \\ y - w/2 & = & 0 \\ z - w/2 & = & 0 \\ 0 & = & 0 \end{array} \Rightarrow \begin{array}{rcl} x & = & w \\ y & = & w/2 \\ z & = & w/2 \\ w & = & w \end{array}$$

Warm Up Exercise

► Sln set:

$$\left[\begin{array}{cccc|c} 1 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1/2 & 0 \\ 0 & 0 & 1 & -1/2 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \Rightarrow \begin{array}{rcl} x - w & = & 0 \\ y - w/2 & = & 0 \\ z - w/2 & = & 0 \\ 0 & = & 0 \end{array} \Rightarrow \begin{array}{rcl} x & = & w \\ y & = & w/2 \\ z & = & w/2 \\ w & = & w \end{array}$$

In vector form:

$$\begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = w \begin{bmatrix} 1 \\ 1/2 \\ 1/2 \\ 1 \end{bmatrix}$$

Parameterized Form

$$\left\{ \begin{bmatrix} t \\ t/2 \\ t/2 \\ t \end{bmatrix} : t \in \mathbb{R} \right\}$$

Where t is a parameter.

Parameterized Form

$$\left\{ \begin{bmatrix} t \\ t/2 \\ t/2 \\ t \end{bmatrix} : t \in \mathbb{R} \right\}$$

Where t is a parameter.

Note:

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \\ w(t) \end{bmatrix} = t \begin{bmatrix} 1 \\ 1/2 \\ 1/2 \\ 1 \end{bmatrix}$$

is a line through the origin in the direction of $[1, 1/2, 1/2, 1]^T$ in \mathbb{R}^4 .

Some Things We Know So Far

1. Systems of eqns \leftrightarrow Matrix equation $Ax = b$.

Some Things We Know So Far

1. Systems of eqns \leftrightarrow Matrix equation $Ax = b$.
 - ▶ Has unique solution $x = A^{-1}b$ if A is nonsingular/invertible.

Some Things We Know So Far

1. Systems of eqns \leftrightarrow Matrix equation $Ax = b$.
 - ▶ Has unique solution $x = A^{-1}b$ if A is nonsingular/invertible.
 - ▶ May have no solution or an infinite number of solutions otherwise!

Some Things We Know So Far

1. Systems of eqns \leftrightarrow Matrix equation $Ax = b$.
 - ▶ Has unique solution $x = A^{-1}b$ if A is nonsingular/invertible.
 - ▶ May have no solution or an infinite number of solutions otherwise!
2. Systems of eqns \leftrightarrow augmented matrices.

Some Things We Know So Far

1. Systems of eqns \leftrightarrow Matrix equation $Ax = b$.
 - ▶ Has unique solution $x = A^{-1}b$ if A is nonsingular/invertible.
 - ▶ May have no solution or an infinite number of solutions otherwise!
2. Systems of eqns \leftrightarrow augmented matrices.
 - ▶ Solve via Gaussian Elimination/Row Reduction/Gauss-Jordan procedure.

Some Things We Know So Far

1. Systems of eqns \leftrightarrow Matrix equation $Ax = b$.
 - ▶ Has unique solution $x = A^{-1}b$ if A is nonsingular/invertible.
 - ▶ May have no solution or an infinite number of solutions otherwise!
2. Systems of eqns \leftrightarrow augmented matrices.
 - ▶ Solve via Gaussian Elimination/Row Reduction/Gauss-Jordan procedure.
 - ▶ Elementary row operations:

Some Things We Know So Far

1. Systems of eqns \leftrightarrow Matrix equation $Ax = b$.
 - ▶ Has unique solution $x = A^{-1}b$ if A is nonsingular/invertible.
 - ▶ May have no solution or an infinite number of solutions otherwise!
2. Systems of eqns \leftrightarrow augmented matrices.
 - ▶ Solve via Gaussian Elimination/Row Reduction/Gauss-Jordan procedure.
 - ▶ Elementary row operations:
 - ▶ Can be represented by (left) matrix multiplication with **elementary** matrices (E).

Some Things We Know So Far

1. Systems of eqns \leftrightarrow Matrix equation $Ax = b$.
 - ▶ Has unique solution $x = A^{-1}b$ if A is nonsingular/invertible.
 - ▶ May have no solution or an infinite number of solutions otherwise!
2. Systems of eqns \leftrightarrow augmented matrices.
 - ▶ Solve via Gaussian Elimination/Row Reduction/Gauss-Jordan procedure.
 - ▶ Elementary row operations:
 - ▶ Can be represented by (left) matrix multiplication with **elementary** matrices (E).
 - ▶ Are invertible. So the corresponding elementary matrices are non-singular. Also: the inverse matrices are elementary!

Some Things We Know So Far

1. Systems of eqns \leftrightarrow Matrix equation $Ax = b$.
 - ▶ Has unique solution $x = A^{-1}b$ if A is nonsingular/invertible.
 - ▶ May have no solution or an infinite number of solutions otherwise!
2. Systems of eqns \leftrightarrow augmented matrices.
 - ▶ Solve via Gaussian Elimination/Row Reduction/Gauss-Jordan procedure.
 - ▶ Elementary row operations:
 - ▶ Can be represented by (left) matrix multiplication with **elementary** matrices (E).
 - ▶ Are invertible. So the corresponding elementary matrices are non-singular. Also: the inverse matrices are elementary!

Question: What relationship is there between singularity (invertibility) and elementary row operations (elementary matrix multiplication)?

Row equivalence

Recall that matrices A and B are **row equivalent** if there are elementary matrices E_1, \dots, E_k and

$$A = E_k E_{k-1} \cdots E_1 B.$$

Row equivalence

Recall that matrices A and B are **row equivalent** if there are elementary matrices E_1, \dots, E_k and

$$A = E_k E_{k-1} \cdots E_1 B.$$

Notes:

1. Row equivalence is *symmetric*.

Row equivalence

Recall that matrices A and B are **row equivalent** if there are elementary matrices E_1, \dots, E_k and

$$A = E_k E_{k-1} \cdots E_1 B.$$

Notes:

1. Row equivalence is *symmetric*. If $A \sim B$ then $B \sim A$ since

$$B = E_1^{-1} \cdots E_k^{-1} A$$

Row equivalence

Recall that matrices A and B are **row equivalent** if there are elementary matrices E_1, \dots, E_k and

$$A = E_k E_{k-1} \cdots E_1 B.$$

Notes:

1. Row equivalence is *symmetric*. If $A \sim B$ then $B \sim A$ since

$$B = E_1^{-1} \cdots E_k^{-1} A$$

and each E_i^{-1} is also elementary.

Row equivalence

Recall that matrices A and B are **row equivalent** if there are elementary matrices E_1, \dots, E_k and

$$A = E_k E_{k-1} \cdots E_1 B.$$

Notes:

1. Row equivalence is *symmetric*. If $A \sim B$ then $B \sim A$ since

$$B = E_1^{-1} \cdots E_k^{-1} A$$

and each E_i^{-1} is also elementary.

2. Row equivalent augmented matrices represent systems of equations with identical solution sets.

Theory (1)

Lemma

Let $A \in \mathbb{R}^{n \times n}$. If the only solution to $Ax = 0$ is the trivial solution, then A is row equivalent to the identity.

Theory (1)

Lemma

Let $A \in \mathbb{R}^{n \times n}$. If the only solution to $Ax = 0$ is the trivial solution, then A is row equivalent to the identity.

Proof.

Row reduce A to an $n \times n$ matrix B in RREF:

$$[A|0] \sim [B|0].$$

Theory (1)

Lemma

Let $A \in \mathbb{R}^{n \times n}$. If the only solution to $Ax = 0$ is the trivial solution, then A is row equivalent to the identity.

Proof.

Row reduce A to an $n \times n$ matrix B in RREF:

$$[A|0] \sim [B|0].$$

$Ax = 0$ and $Bx = 0$ have the same solution set (they are row equivalent). So the augmented matrix $[B|0]$ gives:

$$[B|0] \Leftrightarrow \begin{array}{l} x_1 = 0 \\ x_2 = 0 \\ \vdots \\ x_n = 0 \end{array}$$

Theory (1)

Lemma

Let $A \in \mathbb{R}^{n \times n}$. If the only solution to $Ax = 0$ is the trivial solution, then A is row equivalent to the identity.

Proof.

Row reduce A to an $n \times n$ matrix B in RREF:

$$[A|0] \sim [B|0].$$

$Ax = 0$ and $Bx = 0$ have the same solution set (they are row equivalent). So the augmented matrix $[B|0]$ gives:

$$[B|0] \Leftrightarrow \begin{array}{l} x_1 = 0 \\ x_2 = 0 \\ \vdots \\ x_n = 0 \end{array} \Leftrightarrow [I|0]$$

Hence $B = I$ (RREF is unique).

Theory (2)

Theorem

A matrix A is nonsingular $\Leftrightarrow A = \prod_{i=1}^k E_i$.

Theory (2)

Theorem

A matrix A is nonsingular $\Leftrightarrow A = \prod_{i=1}^k E_i$.

Proof.

(\Leftarrow):

Theory (2)

Theorem

A matrix A is nonsingular $\Leftrightarrow A = \prod_{i=1}^k E_i$.

Proof.

(\Leftarrow): E_i are all invertible. So

$$A = E_1 E_2 \cdots E_k \Rightarrow A^{-1} = (E_k)^{-1} \cdots (E_1)^{-1}$$

and A is not singular.

Theory (2)

Theorem

A matrix A is nonsingular $\Leftrightarrow A = \prod_{i=1}^k E_i$.

Proof.

(\Leftarrow) : E_i are all invertible. So

$$A = E_1 E_2 \cdots E_k \Rightarrow A^{-1} = (E_k)^{-1} \cdots (E_1)^{-1}$$

and A is not singular.

(\Rightarrow) :

Theory (2)

Theorem

A matrix A is nonsingular $\Leftrightarrow A = \prod_{i=1}^k E_i$.

Proof.

(\Leftarrow): E_i are all invertible. So

$$A = E_1 E_2 \cdots E_k \Rightarrow A^{-1} = (E_k)^{-1} \cdots (E_1)^{-1}$$

and A is not singular.

(\Rightarrow): If A^{-1} exists, then $Ax = 0$ has unique solution $A^{-1}0 = 0$.

Theory (2)

Theorem

A matrix A is nonsingular $\Leftrightarrow A = \prod_{i=1}^k E_i$.

Proof.

(\Leftarrow): E_i are all invertible. So

$$A = E_1 E_2 \cdots E_k \Rightarrow A^{-1} = (E_k)^{-1} \cdots (E_1)^{-1}$$

and A is not singular.

(\Rightarrow): If A^{-1} exists, then $Ax = 0$ has unique solution $A^{-1}0 = 0$. So A is row equivalent to I

Theory (2)

Theorem

A matrix A is nonsingular $\Leftrightarrow A = \prod_{i=1}^k E_i$.

Proof.

(\Leftarrow): E_i are all invertible. So

$$A = E_1 E_2 \cdots E_k \Rightarrow A^{-1} = (E_k)^{-1} \cdots (E_1)^{-1}$$

and A is not singular.

(\Rightarrow): If A^{-1} exists, then $Ax = 0$ has unique solution $A^{-1}0 = 0$. So A is row equivalent to I So there exist E_i so that

$$A = E_1 E_2 \cdots E_k I = E_1 E_2 \cdots E_k.$$



Summary

All of the following statements are logically equivalent:

- ▶ A is nonsingular.
- ▶ $Ax = 0$ has only the trivial solution.
- ▶ $Ax = b$ has a unique solution for every b .
- ▶ A is row equivalent to the identity matrix.
- ▶ A is a product of elementary matrices

Application: A Way to Compute Inverses

Suppose that A is invertible.

Application: A Way to Compute Inverses

Suppose that A is invertible. Then we know that A is row equivalent to I :

$$E_k \cdots E_2 E_1 A = I$$

Application: A Way to Compute Inverses

Suppose that A is invertible. Then we know that A is row equivalent to I :

$$E_k \cdots E_2 E_1 A = I \Rightarrow E_k \cdots E_2 E_1 = A^{-1}.$$

Application: A Way to Compute Inverses

Suppose that A is invertible. Then we know that A is row equivalent to I :

$$E_k \cdots E_2 E_1 A = I \Rightarrow E_k \cdots E_2 E_1 = A^{-1}.$$

Therefore the partitioned/augmented matrix $[A|I] \sim [I|A^{-1}]$ since

$$E_k \cdots E_2 E_1 [A|I] = [I|A^{-1}].$$

Application: A Way to Compute Inverses

Suppose that A is invertible. Then we know that A is row equivalent to I :

$$E_k \cdots E_2 E_1 A = I \Rightarrow E_k \cdots E_2 E_1 = A^{-1}.$$

Therefore the partitioned/augmented matrix $[A|I] \sim [I|A^{-1}]$ since

$$E_k \cdots E_2 E_1 [A|I] = [I|A^{-1}].$$

Conclusion: the collection of row operations which transform A to I is the same as the row operations which transform I to A^{-1} .

Example

Compute

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}^{-1}.$$

Example

Compute

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}^{-1}.$$

Write as

$$\left[\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 3 & 4 & 0 & 1 \end{array} \right]$$

Example

Compute

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}^{-1}.$$

Write as

$$\left[\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 3 & 4 & 0 & 1 \end{array} \right] \sim \left[\begin{array}{cc|cc} 1 & 0 & -2 & 1 \\ 0 & 1 & 3/2 & -1/2 \end{array} \right]$$

Example

Compute

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}^{-1}.$$

Write as

$$\left[\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 3 & 4 & 0 & 1 \end{array} \right] \sim \left[\begin{array}{cc|cc} 1 & 0 & -2 & 1 \\ 0 & 1 & 3/2 & -1/2 \end{array} \right]$$

So

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}^{-1} = \begin{bmatrix} -2 & 1 \\ 3/2 & -1/2 \end{bmatrix}$$

Exercises

Compute the following (if they exist):

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 2 & 3 \\ 1 & 2 & 4 \end{bmatrix}^{-1} \quad \text{and} \quad \begin{bmatrix} 1 & 2 & -3 \\ 1 & -2 & 1 \\ 5 & -2 & -3 \end{bmatrix}^{-1}$$

Results

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 2 & 3 \\ 1 & 2 & 4 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & -1 & 0 \\ 3/2 & 1/2 & -3/2 \\ -1 & 0 & 1 \end{bmatrix}$$

Results

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 2 & 3 \\ 1 & 2 & 4 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & -1 & 0 \\ 3/2 & 1/2 & -3/2 \\ -1 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 & -3 \\ 1 & -2 & 1 \\ 5 & -2 & -3 \end{bmatrix} \text{ is singular } (2r_1 + 3r_2 = r_3).$$