

Inner Products

Let V be a vector space. An *inner product* on V is a “pairing” of vectors:

$$(\cdot, \cdot) : V \times V \rightarrow F$$

which maps a pair of vectors to an element of the field. It must satisfy the properties

1. $(u, u) \geq 0$ and $(u, u) = 0 \Leftrightarrow u = 0$.
2. $(u, v) = (v, u)$.
3. $(u, v + w) = (u, v) + (u, w)$.
4. $c(u, v) = (cu, v) = (cv, u)$.

Inner Products

Let V be a vector space. An *inner product* on V is a “pairing” of vectors:

$$(\cdot, \cdot) : V \times V \rightarrow F$$

which maps a pair of vectors to an element of the field. It must satisfy the properties

1. $(u, u) \geq 0$ and $(u, u) = 0 \Leftrightarrow u = 0$.
2. $(u, v) = (v, u)$.
3. $(u, v + w) = (u, v) + (u, w)$.
4. $c(u, v) = (cu, v) = (cv, u)$.

Two standard examples:

- Dot product in \mathbb{R}^n .

Inner Products

Let V be a vector space. An *inner product* on V is a “pairing” of vectors:

$$(\cdot, \cdot) : V \times V \rightarrow F$$

which maps a pair of vectors to an element of the field. It must satisfy the properties

1. $(u, u) \geq 0$ and $(u, u) = 0 \Leftrightarrow u = 0$.
2. $(u, v) = (v, u)$.
3. $(u, v + w) = (u, v) + (u, w)$.
4. $c(u, v) = (cu, v) = (cv, u)$.

Two standard examples:

- ▶ Dot product in \mathbb{R}^n .
- ▶ $\int_a^b fg \, dx$ on integrable functions.

Geometry

Inner products allow us to do geometry:

- ▶ The **norm** of v is

$$\|v\| := \sqrt{(v, v)}.$$

- ▶ The **distance** between vectors v and w is

$$d(v, w) := \|v - w\|.$$

- ▶ The **angle** between vectors v and w is

$$\cos^{-1} \frac{(v, w)}{\|v\| \|w\|}.$$

Geometry

Inner products allow us to do geometry:

- ▶ The **norm** of v is

$$\|v\| := \sqrt{(v, v)}.$$

- ▶ The **distance** between vectors v and w is

$$d(v, w) := \|v - w\|.$$

- ▶ The **angle** between vectors v and w is

$$\cos^{-1} \frac{(v, w)}{\|v\| \|w\|}.$$

Exercise:

Show that $\frac{1}{\sqrt{\pi}} \sin(x)$ is a unit vector using the inner product

$$\int_0^{2\pi} fg \, dx.$$

More Geometry

Inner products allow us to do geometry in places which do not seem “natural.”

More Geometry

Inner products allow us to do geometry in places which do not seem “natural.” Two important geometry theorems:

Theorem (Cauchy-Schwartz)

For any two vectors in an inner product space V we have

$$|(u, v)| \leq \|u\| \|v\|.$$

More Geometry

Inner products allow us to do geometry in places which do not seem “natural.” Two important geometry theorems:

Theorem (Cauchy-Schwartz)

For any two vectors in an inner product space V we have

$$|(u, v)| \leq \|u\| \|v\|.$$

Theorem (Triangle Inequality)

For any two vectors in an inner product space V we have

$$\|u + v\| \leq \|u\| + \|v\|.$$

More Geometry

Inner products allow us to do geometry in places which do not seem “natural.” Two important geometry theorems:

Theorem (Cauchy-Schwartz)

For any two vectors in an inner product space V we have

$$|(u, v)| \leq \|u\| \|v\|.$$

Theorem (Triangle Inequality)

For any two vectors in an inner product space V we have

$$\|u + v\| \leq \|u\| + \|v\|.$$

The point is that these results rely **ONLY** on the properties of the inner products and not **HOW** you compute them or **WHAT** the vectors represent!

Orthogonal and Orthonormal Sets

Definition

An **orthogonal set** of vectors is a set

$$\{v_1, v_2, \dots, v_n\}$$

with the property that

$$(v_i, v_j) = 0, \quad i \neq j.$$

Orthogonal and Orthonormal Sets

Definition

An **orthogonal set** of vectors is a set

$$\{v_1, v_2, \dots, v_n\}$$

with the property that

$$(v_i, v_j) = 0, \quad i \neq j.$$

If, in addition, we have the property

$$(v_i, v_i) = 1$$

then the set is called **orthonormal**.

Orthogonal and Orthonormal Sets

Definition

An **orthogonal set** of vectors is a set

$$\{v_1, v_2, \dots, v_n\}$$

with the property that

$$(v_i, v_j) = 0, \quad i \neq j.$$

If, in addition, we have the property

$$(v_i, v_i) = 1$$

then the set is called **orthonormal**. That is, it is an orthogonal set of unit vectors.

Orthogonal Implies Linear Independence

Theorem

A finite orthogonal set of non-zero vectors is linearly independent.

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$.

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$. We want to show that $a_i = 0$.

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$. We want to show that $a_i = 0$. Take the inner product

$$\left(v_j, \sum_{i=1}^n a_i v_i \right).$$

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$. We want to show that $a_i = 0$. Take the inner product

$$\left(v_j, \sum_{i=1}^n a_i v_i \right).$$

Note that this must be zero.

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$. We want to show that $a_i = 0$. Take the inner product

$$\left(v_j, \sum_{i=1}^n a_i v_i \right).$$

Note that this must be zero.

Also we have

$$\left(v_j, \sum_{i=1}^n a_i v_i \right) = \sum_{i=1}^n a_i (v_j, v_i)$$

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$. We want to show that $a_i = 0$. Take the inner product

$$\left(v_j, \sum_{i=1}^n a_i v_i \right).$$

Note that this must be zero.

Also we have

$$\left(v_j, \sum_{i=1}^n a_i v_i \right) = \sum_{i=1}^n a_i (v_j, v_i) = a_j (v_j, v_j).$$

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$. We want to show that $a_j = 0$. Take the inner product

$$\left(v_j, \sum_{i=1}^n a_i v_i \right).$$

Note that this must be zero.

Also we have

$$\left(v_j, \sum_{i=1}^n a_i v_i \right) = \sum_{i=1}^n a_i (v_j, v_i) = a_j (v_j, v_j).$$

So we have $0 = a_j (v_j, v_j)$.

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$. We want to show that $a_j = 0$. Take the inner product

$$\left(v_j, \sum_{i=1}^n a_i v_i \right).$$

Note that this must be zero.

Also we have

$$\left(v_j, \sum_{i=1}^n a_i v_i \right) = \sum_{i=1}^n a_i (v_j, v_i) = a_j (v_j, v_j).$$

So we have $0 = a_j (v_j, v_j)$. Now, $(v_j, v_j) \neq 0$ therefore $a_j = 0$. □

How to write coordinates in an orthogonal basis

The proof offers an interesting result:

How to write coordinates in an orthogonal basis

The proof offers an interesting result: Let $\{v_i\}$ be a (nonempty), finite, orthogonal set.

How to write coordinates in an orthogonal basis

The proof offers an interesting result: Let $\{v_i\}$ be a (nonempty), finite, orthogonal set. Then it is linearly independent and a basis for $\text{span}\{v_i\}$.

How to write coordinates in an orthogonal basis

The proof offers an interesting result: Let $\{v_i\}$ be a (nonempty), finite, orthogonal set. Then it is linearly independent and a basis for $\text{span}\{v_i\}$. Let $x \in \text{span}\{v_i\}$.

How to write coordinates in an orthogonal basis

The proof offers an interesting result: Let $\{v_i\}$ be a (nonempty), finite, orthogonal set. Then it is linearly independent and a basis for $\text{span}\{v_i\}$. Let $x \in \text{span}\{v_i\}$. So

$$x = \sum_{i=1}^n a_i v_i.$$

How to write coordinates in an orthogonal basis

The proof offers an interesting result: Let $\{v_i\}$ be a (nonempty), finite, orthogonal set. Then it is linearly independent and a basis for $\text{span}\{v_i\}$. Let $x \in \text{span}\{v_i\}$. So

$$x = \sum_{i=1}^n a_i v_i.$$

Note that from the proof:

$$\left(v_j, \sum_{i=1}^n a_i v_i \right) = a_j (v_j, v_j) = a_j \|v_j\|^2.$$

How to write coordinates in an orthogonal basis

The proof offers an interesting result: Let $\{v_i\}$ be a (nonempty), finite, orthogonal set. Then it is linearly independent and a basis for $\text{span}\{v_i\}$. Let $x \in \text{span}\{v_i\}$. So

$$x = \sum_{i=1}^n a_i v_i.$$

Note that from the proof:

$$\left(v_j, \sum_{i=1}^n a_i v_i \right) = a_j (v_j, v_j) = a_j \|v_j\|^2.$$

So:

$$a_j = \frac{(v_j, x)}{\|v_j\|^2}.$$

How to write coordinates in an orthogonal basis

The proof offers an interesting result: Let $\{v_i\}$ be a (nonempty), finite, orthogonal set. Then it is linearly independent and a basis for $\text{span}\{v_i\}$. Let $x \in \text{span}\{v_i\}$. So

$$x = \sum_{i=1}^n a_i v_i.$$

Note that from the proof:

$$\left(v_j, \sum_{i=1}^n a_i v_i \right) = a_j (v_j, v_j) = a_j \|v_j\|^2.$$

So:

$$a_j = \frac{(v_j, x)}{\|v_j\|^2}.$$

This means that *vector coordinates* can be expressed in terms of inner products!

Orthonormal Vectors

If our set of vectors is orthonormal, then we can do better:

$$a_j = \frac{(v_j, x)}{\|v_j\|^2} = (v_j, x).$$

Orthonormal Vectors

If our set of vectors is orthonormal, then we can do better:

$$a_j = \frac{(v_j, x)}{\|v_j\|^2} = (v_j, x).$$

In fact, orthonormal vectors can reduce all inner products to a “dot product”!

Orthonormal Vectors

If our set of vectors is orthonormal, then we can do better:

$$a_j = \frac{(v_j, x)}{\|v_j\|^2} = (v_j, x).$$

In fact, orthonormal vectors can reduce all inner products to a “dot product”!

Theorem

Let $V = \text{span}(S)$ where $S = \{v_1, \dots, v_n\}$ is an orthonormal set.

Let $x, y \in V$ and write them as

$$x = a_1 v_1 + \dots + a_n v_n = \sum_{i=1}^n a_i v_i, \quad y = b_1 v_1 + \dots + b_n v_n = \sum_{i=1}^n b_i v_i.$$

Then

$$(x, y) = \sum_{i=1}^n a_i b_i.$$

Proof

To prove, let's compute (x, y) :

Proof

To prove, let's compute (x, y) :

$$(x, y) = \left(\sum_{i=1}^n a_i v_i, \sum_{j=1}^n b_j v_j \right)$$

Proof

To prove, let's compute (x, y) :

$$\begin{aligned}(x, y) &= \left(\sum_{i=1}^n a_i v_i, \sum_{j=1}^n b_j v_j \right) \\ &= \sum_{i=1}^n \sum_{j=1}^n a_i b_j (v_i, v_j)\end{aligned}$$

Proof

To prove, let's compute (x, y) :

$$\begin{aligned}(x, y) &= \left(\sum_{i=1}^n a_i v_i, \sum_{j=1}^n b_j v_j \right) \\&= \sum_{i=1}^n \sum_{j=1}^n a_i b_j (v_i, v_j) \\&= \sum_{i=1}^n a_i b_i (v_i, v_i)\end{aligned}$$

Proof

To prove, let's compute (x, y) :

$$\begin{aligned}(x, y) &= \left(\sum_{i=1}^n a_i v_i, \sum_{j=1}^n b_j v_j \right) \\&= \sum_{i=1}^n \sum_{j=1}^n a_i b_j (v_i, v_j) \\&= \sum_{i=1}^n a_i b_i (v_i, v_i) \\&= \sum_{i=1}^n a_i b_i.\end{aligned}$$

Summary and question

So far, we have many nice properties IF we have an orthonormal basis.

Summary and question

So far, we have many nice properties IF we have an orthonormal basis.

Question: given a vector space, can we always find an orthonormal basis?

Summary and question

So far, we have many nice properties IF we have an orthonormal basis.

Question: given a vector space, can we always find an orthonormal basis? YES!

The Gram-Schmidt Process

Theorem (Gram-Schmidt)

Let V be a (nontrivial) vector space of dimension m with an inner product. Then V has an orthonormal basis.

The Gram-Schmidt Process

Theorem (Gram-Schmidt)

Let V be a (nontrivial) vector space of dimension m with an inner product. Then V has an orthonormal basis.

The proof of this is constructive and the method is known as the *Gram-Schmidt* process.

The Gram-Schmidt Process

Theorem (Gram-Schmidt)

Let V be a (nontrivial) vector space of dimension m with an inner product. Then V has an orthonormal basis.

The proof of this is constructive and the method is known as the *Gram-Schmidt* process. The method is worth knowing over the actual result of the theorem!

The Gram-Schmidt Process

Let $\{w_1, w_2, \dots, w_n\}$ be a basis for V .

The Gram-Schmidt Process

Let $\{w_1, w_2, \dots, w_n\}$ be a basis for V . The goal is the following:

The Gram-Schmidt Process

Let $\{w_1, w_2, \dots, w_n\}$ be a basis for V . The goal is the following:

Construct a list of orthonormal vectors $\{v_1, \dots, v_n\}$ so that

$$\text{span}\{v_1\} = \text{span}\{w_1\}$$

The Gram-Schmidt Process

Let $\{w_1, w_2, \dots, w_n\}$ be a basis for V . The goal is the following:

Construct a list of orthonormal vectors $\{v_1, \dots, v_n\}$ so that

$$\text{span}\{v_1\} = \text{span}\{w_1\}$$

$$\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$$

The Gram-Schmidt Process

Let $\{w_1, w_2, \dots, w_n\}$ be a basis for V . The goal is the following:

Construct a list of orthonormal vectors $\{v_1, \dots, v_n\}$ so that

$$\text{span}\{v_1\} = \text{span}\{w_1\}$$

$$\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$$

$$\text{span}\{v_1, v_2, v_3\} = \text{span}\{w_1, w_2, w_3\}$$

$$\vdots$$

The Gram-Schmidt Process

Let $\{w_1, w_2, \dots, w_n\}$ be a basis for V . The goal is the following:

Construct a list of orthonormal vectors $\{v_1, \dots, v_n\}$ so that

$$\text{span}\{v_1\} = \text{span}\{w_1\}$$

$$\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$$

$$\text{span}\{v_1, v_2, v_3\} = \text{span}\{w_1, w_2, w_3\}$$

$$\vdots$$

To begin, set $v_1 = \frac{1}{\|w_1\|} w_1$.

The Gram-Schmidt Process

Let $\{w_1, w_2, \dots, w_n\}$ be a basis for V . The goal is the following:

Construct a list of orthonormal vectors $\{v_1, \dots, v_n\}$ so that

$$\text{span}\{v_1\} = \text{span}\{w_1\}$$

$$\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$$

$$\text{span}\{v_1, v_2, v_3\} = \text{span}\{w_1, w_2, w_3\}$$

$$\vdots$$

To begin, set $v_1 = \frac{1}{\|w_1\|} w_1$.

Clearly, v_1 is a unit vector and $\text{span}\{v_1\} = \text{span}\{w_1\}$.

The Gram-Schmidt Process

Next, we want to find a vector v_2 so that $\{v_1, v_2\}$ is orthonormal and $\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$.

The Gram-Schmidt Process

Next, we want to find a vector v_2 so that $\{v_1, v_2\}$ is orthonormal and $\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$.

First note that $\text{span}\{w_1, w_2\} = \text{span}\{v_1, w_2\}$.

The Gram-Schmidt Process

Next, we want to find a vector v_2 so that $\{v_1, v_2\}$ is orthonormal and $\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$.

First note that $\text{span}\{w_1, w_2\} = \text{span}\{v_1, w_2\}$.

So if we can find v_2 so that $w_2 \in \text{span}\{v_1, v_2\}$ then we will have $\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$.

The Gram-Schmidt Process

Next, we want to find a vector v_2 so that $\{v_1, v_2\}$ is orthonormal and $\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$.

First note that $\text{span}\{w_1, w_2\} = \text{span}\{v_1, w_2\}$.

So if we can find v_2 so that $w_2 \in \text{span}\{v_1, v_2\}$ then we will have $\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$.

So write $w_2 = av_1 + bv_2$

The Gram-Schmidt Process

Next, we want to find a vector v_2 so that $\{v_1, v_2\}$ is orthonormal and $\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$.

First note that $\text{span}\{w_1, w_2\} = \text{span}\{v_1, w_2\}$.

So if we can find v_2 so that $w_2 \in \text{span}\{v_1, v_2\}$ then we will have $\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$.

So write $w_2 = av_1 + bv_2$ and consider the inner product:

$$(v_1, w_2) = a(v_1, v_1) + b(v_1, v_2)$$

The Gram-Schmidt Process

Next, we want to find a vector v_2 so that $\{v_1, v_2\}$ is orthonormal and $\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$.

First note that $\text{span}\{w_1, w_2\} = \text{span}\{v_1, w_2\}$.

So if we can find v_2 so that $w_2 \in \text{span}\{v_1, v_2\}$ then we will have $\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$.

So write $w_2 = av_1 + bv_2$ and consider the inner product:

$$(v_1, w_2) = a(v_1, v_1) + b(v_1, v_2) = a.$$

Therefore $bv_2 = w_2 - (v_1, w_2)v_1$.

The Gram-Schmidt Process

Next, we want to find a vector v_2 so that $\{v_1, v_2\}$ is orthonormal and $\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$.

First note that $\text{span}\{w_1, w_2\} = \text{span}\{v_1, w_2\}$.

So if we can find v_2 so that $w_2 \in \text{span}\{v_1, v_2\}$ then we will have $\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$.

So write $w_2 = av_1 + bv_2$ and consider the inner product:

$$(v_1, w_2) = a(v_1, v_1) + b(v_1, v_2) = a.$$

Therefore $bv_2 = w_2 - (v_1, w_2)v_1$. Take the norm to find

$$\|bv_2\| = \|w_2 - (v_1, w_2)v_1\|$$

The Gram-Schmidt Process

Next, we want to find a vector v_2 so that $\{v_1, v_2\}$ is orthonormal and $\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$.

First note that $\text{span}\{w_1, w_2\} = \text{span}\{v_1, w_2\}$.

So if we can find v_2 so that $w_2 \in \text{span}\{v_1, v_2\}$ then we will have $\text{span}\{v_1, v_2\} = \text{span}\{w_1, w_2\}$.

So write $w_2 = av_1 + bv_2$ and consider the inner product:

$$(v_1, w_2) = a(v_1, v_1) + b(v_1, v_2) = a.$$

Therefore $bv_2 = w_2 - (v_1, w_2)v_1$. Take the norm to find

$$\|bv_2\| = \|w_2 - (v_1, w_2)v_1\| \Rightarrow b = \|w_2 - (v_1, w_2)v_1\|.$$

The Gram-Schmidt Process

In general: Let $\{v_1, \dots, v_n\}$ be an ON set so that

$$\text{span}\{v_1, \dots, v_n\} = \text{span}\{w_1, \dots, w_n\}.$$

The Gram-Schmidt Process

In general: Let $\{v_1, \dots, v_n\}$ be an ON set so that

$$\text{span}\{v_1, \dots, v_n\} = \text{span}\{w_1, \dots, w_n\}.$$

In order to find a vector v_{n+1} so that $\{v_1, \dots, v_{n+1}\}$ is ON and

$$\text{span}\{v_1, \dots, v_{n+1}\} = \text{span}\{w_1, \dots, w_{n+1}\}$$

Compute a unit vector in the direction of

$$w_{n+1} - \sum_{i=1}^n (v_i, w_{n+1}) v_i.$$

Exercise

Compute an ON basis for the subset of \mathbb{R}^3 spanned by

$$\begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$