

Some Special Matrices

An $n \times n$ matrix A is **diagonal** if $a_{ij} = 0$ when $i \neq j$.

A matrix S is **scalar** if it is diagonal and all of its diagonal elements are equal.

The $n \times n$ **identity** matrix I_n is a diagonal matrix with only 1s on the diagonal.

An $n \times n$ matrix U is **upper triangular** if $u_{ij} = 0$ when $i > j$.

An $n \times n$ matrix L is **lower triangular** if $\ell_{ij} = 0$ when $i < j$.

Give an example of each type of matrix in $\mathbb{R}^{3 \times 3}$.

Properties of Matrix Operations

We know a few matrix operations now:

Addition, Scalar multiplication, Transposition, and Multiplication.

Some of these operations exhibit interesting properties.

Properties of Matrix Operations

We know a few matrix operations now:

Addition, Scalar multiplication, Transposition, and Multiplication.

Some of these operations exhibit interesting properties.

(And some don't.)

Properties of Matrix Addition

Matrix addition works the way that you expect. Assume that A, B, C are all matrices of appropriate sizes.

Properties of Matrix Addition

Matrix addition works the way that you expect. Assume that A, B, C are all matrices of appropriate sizes.

- ▶ It is commutative:

Properties of Matrix Addition

Matrix addition works the way that you expect. Assume that A, B, C are all matrices of appropriate sizes.

- ▶ It is commutative: $A + B = B + A$.

Properties of Matrix Addition

Matrix addition works the way that you expect. Assume that A, B, C are all matrices of appropriate sizes.

- ▶ It is commutative: $A + B = B + A$.
- ▶ It is associative:

Properties of Matrix Addition

Matrix addition works the way that you expect. Assume that A, B, C are all matrices of appropriate sizes.

- ▶ It is commutative: $A + B = B + A$.
- ▶ It is associative: $(A + B) + C = A + (B + C)$.

Properties of Matrix Addition

Matrix addition works the way that you expect. Assume that A, B, C are all matrices of appropriate sizes.

- ▶ It is commutative: $A + B = B + A$.
- ▶ It is associative: $(A + B) + C = A + (B + C)$.
- ▶ It has an identity:

Properties of Matrix Addition

Matrix addition works the way that you expect. Assume that A, B, C are all matrices of appropriate sizes.

- ▶ It is commutative: $A + B = B + A$.
- ▶ It is associative: $(A + B) + C = A + (B + C)$.
- ▶ It has an identity: $A + 0 = A$.

Properties of Matrix Addition

Matrix addition works the way that you expect. Assume that A, B, C are all matrices of appropriate sizes.

- ▶ It is commutative: $A + B = B + A$.
- ▶ It is associative: $(A + B) + C = A + (B + C)$.
- ▶ It has an identity: $A + 0 = A$.

0 is a *matrix* where all entries are the real number 0 .

Properties of Matrix Addition

Matrix addition works the way that you expect. Assume that A, B, C are all matrices of appropriate sizes.

- ▶ It is commutative: $A + B = B + A$.
- ▶ It is associative: $(A + B) + C = A + (B + C)$.
- ▶ It has an identity: $A + 0 = A$.
0 is a *matrix* where all entries are the real number 0.
- ▶ It has additive inverses:

Properties of Matrix Addition

Matrix addition works the way that you expect. Assume that A, B, C are all matrices of appropriate sizes.

- ▶ It is commutative: $A + B = B + A$.
- ▶ It is associative: $(A + B) + C = A + (B + C)$.
- ▶ It has an identity: $A + 0 = A$.
0 is a *matrix* where all entries are the real number 0.
- ▶ It has additive inverses: $A + (-A) = 0$.

Properties of Matrix Addition

Matrix addition works the way that you expect. Assume that A, B, C are all matrices of appropriate sizes.

- ▶ It is commutative: $A + B = B + A$.
- ▶ It is associative: $(A + B) + C = A + (B + C)$.
- ▶ It has an identity: $A + 0 = A$.

0 is a *matrix* where all entries are the real number 0 .

- ▶ It has additive inverses: $A + (-A) = 0$.
 $-A$ is the matrix $-1 * A$.

Properties of Matrix Addition

Matrix addition works the way that you expect. Assume that A, B, C are all matrices of appropriate sizes.

- ▶ It is commutative: $A + B = B + A$.
- ▶ It is associative: $(A + B) + C = A + (B + C)$.
- ▶ It has an identity: $A + 0 = A$.

0 is a *matrix* where all entries are the real number 0 .

- ▶ It has additive inverses: $A + (-A) = 0$.
 $-A$ is the matrix $-1 * A$.

(Note: This makes $\mathbb{R}^{n \times m}$ a *group* under addition.)

Properties of Matrix Addition

Matrix addition works the way that you expect. Assume that A, B, C are all matrices of appropriate sizes.

- ▶ It is commutative: $A + B = B + A$.
- ▶ It is associative: $(A + B) + C = A + (B + C)$.
- ▶ It has an identity: $A + 0 = A$.

0 is a *matrix* where all entries are the real number 0 .

- ▶ It has additive inverses: $A + (-A) = 0$.
 $-A$ is the matrix $-1 * A$.

(Note: This makes $\mathbb{R}^{n \times m}$ a *group* under addition.)

Summary: Matrix Addition is pretty normal.

Proof Interlude

All of these properties are true because the definition of addition of matrices works well with the properties of addition for real number already.

Proof Interlude

All of these properties are true because the definition of addition of matrices works well with the properties of addition for real number already. To illustrate this, let's prove the commutative property:

Proof Interlude

All of these properties are true because the definition of addition of matrices works well with the properties of addition for real number already. To illustrate this, let's prove the commutative property:

Commutative Property of Matrix Addition.

Let $A = [a_{ij}]$ and $B = [b_{ij}]$ both be $n \times m$ real matrices.

Proof Interlude

All of these properties are true because the definition of addition of matrices works well with the properties of addition for real number already. To illustrate this, let's prove the commutative property:

Commutative Property of Matrix Addition.

Let $A = [a_{ij}]$ and $B = [b_{ij}]$ both be $n \times m$ real matrices.

Then

$$A + B := [a_{ij} + b_{ij}] = [b_{ij} + a_{ij}] =: B + A.$$

Proof Interlude

All of these properties are true because the definition of addition of matrices works well with the properties of addition for real number already. To illustrate this, let's prove the commutative property:

Commutative Property of Matrix Addition.

Let $A = [a_{ij}]$ and $B = [b_{ij}]$ both be $n \times m$ real matrices.

Then

$$A + B := [a_{ij} + b_{ij}] = [b_{ij} + a_{ij}] =: B + A.$$

The left and right equalities hold by definition.

Proof Interlude

All of these properties are true because the definition of addition of matrices works well with the properties of addition for real number already. To illustrate this, let's prove the commutative property:

Commutative Property of Matrix Addition.

Let $A = [a_{ij}]$ and $B = [b_{ij}]$ both be $n \times m$ real matrices.

Then

$$A + B := [a_{ij} + b_{ij}] = [b_{ij} + a_{ij}] =: B + A.$$

The left and right equalities hold by definition. The center equality holds by the properties of addition of real numbers. \square

Proof Interlude

All of these properties are true because the definition of addition of matrices works well with the properties of addition for real number already. To illustrate this, let's prove the commutative property:

Commutative Property of Matrix Addition.

Let $A = [a_{ij}]$ and $B = [b_{ij}]$ both be $n \times m$ real matrices.

Then

$$A + B := [a_{ij} + b_{ij}] = [b_{ij} + a_{ij}] =: B + A.$$

The left and right equalities hold by definition. The center equality holds by the properties of addition of real numbers. \square

The idea is:

Proof Interlude

All of these properties are true because the definition of addition of matrices works well with the properties of addition for real number already. To illustrate this, let's prove the commutative property:

Commutative Property of Matrix Addition.

Let $A = [a_{ij}]$ and $B = [b_{ij}]$ both be $n \times m$ real matrices.

Then

$$A + B := [a_{ij} + b_{ij}] = [b_{ij} + a_{ij}] =: B + A.$$

The left and right equalities hold by definition. The center equality holds by the properties of addition of real numbers. \square

The idea is:

Matrices break down into components

Proof Interlude

All of these properties are true because the definition of addition of matrices works well with the properties of addition for real number already. To illustrate this, let's prove the commutative property:

Commutative Property of Matrix Addition.

Let $A = [a_{ij}]$ and $B = [b_{ij}]$ both be $n \times m$ real matrices.

Then

$$A + B := [a_{ij} + b_{ij}] = [b_{ij} + a_{ij}] =: B + A.$$

The left and right equalities hold by definition. The center equality holds by the properties of addition of real numbers. \square

The idea is:

Matrices break down into components
which are real numbers

Proof Interlude

All of these properties are true because the definition of addition of matrices works well with the properties of addition for real number already. To illustrate this, let's prove the commutative property:

Commutative Property of Matrix Addition.

Let $A = [a_{ij}]$ and $B = [b_{ij}]$ both be $n \times m$ real matrices.

Then

$$A + B := [a_{ij} + b_{ij}] = [b_{ij} + a_{ij}] =: B + A.$$

The left and right equalities hold by definition. The center equality holds by the properties of addition of real numbers. \square

The idea is:

Matrices break down into components
which are real numbers
which commute when added.

Properties of Scalar Multiplication

Let $r, s \in \mathbb{R}$ and A, B matrices of appropriate sizes. Then for scalar multiplication

- ▶ $(rs)A = r(sA).$

Properties of Scalar Multiplication

Let $r, s \in \mathbb{R}$ and A, B matrices of appropriate sizes. Then for scalar multiplication

- ▶ $(rs)A = r(sA)$. (An associativity property.)

Properties of Scalar Multiplication

Let $r, s \in \mathbb{R}$ and A, B matrices of appropriate sizes. Then for scalar multiplication

- ▶ $(rs)A = r(sA)$. (An associativity property.)
- ▶ $(r + s)A = rA + sA$.

Properties of Scalar Multiplication

Let $r, s \in \mathbb{R}$ and A, B matrices of appropriate sizes. Then for scalar multiplication

- ▶ $(rs)A = r(sA)$. (An associativity property.)
- ▶ $(r + s)A = rA + sA$. (A distributive property.)

Properties of Scalar Multiplication

Let $r, s \in \mathbb{R}$ and A, B matrices of appropriate sizes. Then for scalar multiplication

- ▶ $(rs)A = r(sA)$. (An associativity property.)
- ▶ $(r + s)A = rA + sA$. (A distributive property.)
- ▶ $r(A + B) = rA + rB$.

Properties of Scalar Multiplication

Let $r, s \in \mathbb{R}$ and A, B matrices of appropriate sizes. Then for scalar multiplication

- ▶ $(rs)A = r(sA)$. (An associativity property.)
- ▶ $(r + s)A = rA + sA$. (A distributive property.)
- ▶ $r(A + B) = rA + rB$. (Another distributive property.)

Properties of Scalar Multiplication

Let $r, s \in \mathbb{R}$ and A, B matrices of appropriate sizes. Then for scalar multiplication

- ▶ $(rs)A = r(sA)$. (An associativity property.)
- ▶ $(r + s)A = rA + sA$. (A distributive property.)
- ▶ $r(A + B) = rA + rB$. (Another distributive property.)
- ▶ $1A = A$.

Properties of Scalar Multiplication

Let $r, s \in \mathbb{R}$ and A, B matrices of appropriate sizes. Then for scalar multiplication

- ▶ $(rs)A = r(sA)$. (An associativity property.)
- ▶ $(r + s)A = rA + sA$. (A distributive property.)
- ▶ $r(A + B) = rA + rB$. (Another distributive property.)
- ▶ $1A = A$. (Duh.)

Properties of Scalar Multiplication

Let $r, s \in \mathbb{R}$ and A, B matrices of appropriate sizes. Then for scalar multiplication

- ▶ $(rs)A = r(sA)$. (An associativity property.)
- ▶ $(r + s)A = rA + sA$. (A distributive property.)
- ▶ $r(A + B) = rA + rB$. (Another distributive property.)
- ▶ $1A = A$. (Duh.)

Note: These properties along with those from matrix addition make $\mathbb{R}^{n \times m}$ a(n abstract) **vector space**.

Properties of Scalar Multiplication

Let $r, s \in \mathbb{R}$ and A, B matrices of appropriate sizes. Then for scalar multiplication

- ▶ $(rs)A = r(sA)$. (An associativity property.)
- ▶ $(r + s)A = rA + sA$. (A distributive property.)
- ▶ $r(A + B) = rA + rB$. (Another distributive property.)
- ▶ $1A = A$. (Duh.)

Note: These properties along with those from matrix addition make $\mathbb{R}^{n \times m}$ a(n abstract) **vector space**.

Summary: Scalar multiplication also works the way that you expect.

Properties of Scalar Multiplication

Let $r, s \in \mathbb{R}$ and A, B matrices of appropriate sizes. Then for scalar multiplication

- ▶ $(rs)A = r(sA)$. (An associativity property.)
- ▶ $(r + s)A = rA + sA$. (A distributive property.)
- ▶ $r(A + B) = rA + rB$. (Another distributive property.)
- ▶ $1A = A$. (Duh.)

Note: These properties along with those from matrix addition make $\mathbb{R}^{n \times m}$ a(n abstract) **vector space**.

Summary: Scalar multiplication also works the way that you expect.

The reason, again, is that the component-wise definition of scalar multiplication agrees with the standard multiplicative properties of the real numbers.

Properties of Transposition

Transposition has some interesting properties worth noting. There is no real analog to these in the real numbers, since transposition depends on the 2-D nature of the matrix.

Properties of Transposition

Transposition has some interesting properties worth noting. There is no real analog to these in the real numbers, since transposition depends on the 2-D nature of the matrix. Here r is a real number and A, B are both appropriately sized matrices.

- ▶ $(A^T)^T = A.$

Properties of Transposition

Transposition has some interesting properties worth noting. There is no real analog to these in the real numbers, since transposition depends on the 2-D nature of the matrix. Here r is a real number and A, B are both appropriately sized matrices.

- ▶ $(A^T)^T = A$. You could say that transposition is its own inverse.

Properties of Transposition

Transposition has some interesting properties worth noting. There is no real analog to these in the real numbers, since transposition depends on the 2-D nature of the matrix. Here r is a real number and A, B are both appropriately sized matrices.

- ▶ $(A^T)^T = A$. You could say that transposition is its own inverse.
- ▶ $(A + B)^T = A^T + B^T$.

Properties of Transposition

Transposition has some interesting properties worth noting. There is no real analog to these in the real numbers, since transposition depends on the 2-D nature of the matrix. Here r is a real number and A, B are both appropriately sized matrices.

- ▶ $(A^T)^T = A$. You could say that transposition is its own inverse.
- ▶ $(A + B)^T = A^T + B^T$. Transposition “distributes.”

Properties of Transposition

Transposition has some interesting properties worth noting. There is no real analog to these in the real numbers, since transposition depends on the 2-D nature of the matrix. Here r is a real number and A, B are both appropriately sized matrices.

- ▶ $(A^T)^T = A$. You could say that transposition is its own inverse.
- ▶ $(A + B)^T = A^T + B^T$. Transposition “distributes.”
- ▶ $(rA)^T = rA^T$

Properties of Transposition

Transposition has some interesting properties worth noting. There is no real analog to these in the real numbers, since transposition depends on the 2-D nature of the matrix. Here r is a real number and A, B are both appropriately sized matrices.

- ▶ $(A^T)^T = A$. You could say that transposition is its own inverse.
- ▶ $(A + B)^T = A^T + B^T$. Transposition “distributes.”
- ▶ $(rA)^T = rA^T$ Scalar multiplication “factors” out of transposition.

Properties of Transposition

Transposition has some interesting properties worth noting. There is no real analog to these in the real numbers, since transposition depends on the 2-D nature of the matrix. Here r is a real number and A, B are both appropriately sized matrices.

- ▶ $(A^T)^T = A$. You could say that transposition is its own inverse.
- ▶ $(A + B)^T = A^T + B^T$. Transposition “distributes.”
- ▶ $(rA)^T = rA^T$ Scalar multiplication “factors” out of transposition.

These last two properties show that transposition is a **linear** operation.

Exercise Break

Compute the following:

1.

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 4 & -6 \\ -2 & 3 \end{bmatrix}$$

2.

$$\begin{bmatrix} 1 & 3 & 2 \\ 2 & -1 & 3 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 2 & 2 \\ 3 & -1 \end{bmatrix}$$

3.

$$\begin{bmatrix} 0 & 2 & 3 \\ 1 & 2 & -1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & -1 \\ 2 & 3 \end{bmatrix}$$

Results:

1.

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 4 & -6 \\ -2 & 3 \end{bmatrix} =$$

Results:

1.

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 4 & -6 \\ -2 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

Results:

1.

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 4 & -6 \\ -2 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}. \quad \text{Weird.}$$

Results:

1.

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 4 & -6 \\ -2 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}. \quad \text{Weird.}$$

2.

$$\begin{bmatrix} 1 & 3 & 2 \\ 2 & -1 & 3 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 2 & 2 \\ 3 & -1 \end{bmatrix} =$$

Results:

1.

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 4 & -6 \\ -2 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}. \quad \text{Weird.}$$

2.

$$\begin{bmatrix} 1 & 3 & 2 \\ 2 & -1 & 3 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 2 & 2 \\ 3 & -1 \end{bmatrix} = \begin{bmatrix} 12 & 5 \\ 7 & -3 \end{bmatrix}$$

Results:

1.

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 4 & -6 \\ -2 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}. \quad \text{Weird.}$$

2.

$$\begin{bmatrix} 1 & 3 & 2 \\ 2 & -1 & 3 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 2 & 2 \\ 3 & -1 \end{bmatrix} = \begin{bmatrix} 12 & 5 \\ 7 & -3 \end{bmatrix}$$

3.

$$\begin{bmatrix} 0 & 2 & 3 \\ 1 & 2 & -1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & -1 \\ 2 & 3 \end{bmatrix} =$$

Results:

1.

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 4 & -6 \\ -2 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}. \quad \text{Weird.}$$

2.

$$\begin{bmatrix} 1 & 3 & 2 \\ 2 & -1 & 3 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 2 & 2 \\ 3 & -1 \end{bmatrix} = \begin{bmatrix} 12 & 5 \\ 7 & -3 \end{bmatrix}$$

3.

$$\begin{bmatrix} 0 & 2 & 3 \\ 1 & 2 & -1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & -1 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} 12 & 7 \\ 5 & -3 \end{bmatrix}.$$

Results:

1.

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 4 & -6 \\ -2 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}. \quad \text{Weird.}$$

2.

$$\begin{bmatrix} 1 & 3 & 2 \\ 2 & -1 & 3 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 2 & 2 \\ 3 & -1 \end{bmatrix} = \begin{bmatrix} 12 & 5 \\ 7 & -3 \end{bmatrix}$$

3.

$$\begin{bmatrix} 0 & 2 & 3 \\ 1 & 2 & -1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & -1 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} 12 & 7 \\ 5 & -3 \end{bmatrix}. \quad \text{Neat.}$$

The “Neat” Property of Transposition

From our examples, it appears that $(AB)^T = B^T A^T$. Let's prove it:

The “Neat” Property of Transposition

From our examples, it appears that $(AB)^T = B^T A^T$. Let's prove it:

Proof.

Our goal is to show that the (i,j) -th element of $(AB)^T$ is the same as the (i,j) -th element of $B^T A^T$.

The “Neat” Property of Transposition

From our examples, it appears that $(AB)^T = B^T A^T$. Let's prove it:

Proof.

Our goal is to show that the (i,j) -th element of $(AB)^T$ is the same as the (i,j) -th element of $B^T A^T$. Note that:

$$\left((AB)^T \right)_{ij}$$

The “Neat” Property of Transposition

From our examples, it appears that $(AB)^T = B^T A^T$. Let's prove it:

Proof.

Our goal is to show that the (i, j) -th element of $(AB)^T$ is the same as the (i, j) -th element of $B^T A^T$. Note that:

$$\left((AB)^T\right)_{ij} = (AB)_{ji}$$

The “Neat” Property of Transposition

From our examples, it appears that $(AB)^T = B^T A^T$. Let's prove it:

Proof.

Our goal is to show that the (i, j) -th element of $(AB)^T$ is the same as the (i, j) -th element of $B^T A^T$. Note that:

$$\left((AB)^T\right)_{ij} = (AB)_{ji} = \sum_{k=1}^{\ell} a_{jk} b_{ki}$$

The “Neat” Property of Transposition

From our examples, it appears that $(AB)^T = B^T A^T$. Let's prove it:

Proof.

Our goal is to show that the (i, j) -th element of $(AB)^T$ is the same as the (i, j) -th element of $B^T A^T$. Note that:

$$\left((AB)^T\right)_{ij} = (AB)_{ji} = \sum_{k=1}^{\ell} a_{jk} b_{ki}$$

and

$$\left(B^T A^T\right)_{ij}$$

The “Neat” Property of Transposition

From our examples, it appears that $(AB)^T = B^T A^T$. Let's prove it:

Proof.

Our goal is to show that the (i,j) -th element of $(AB)^T$ is the same as the (i,j) -th element of $B^T A^T$. Note that:

$$\left((AB)^T\right)_{ij} = (AB)_{ji} = \sum_{k=1}^{\ell} a_{jk} b_{ki}$$

and

$$\left(B^T A^T\right)_{ij} = \sum_{k=1}^{\ell} b_{ik}^T a_{kj}^T$$

The “Neat” Property of Transposition

From our examples, it appears that $(AB)^T = B^T A^T$. Let's prove it:

Proof.

Our goal is to show that the (i,j) -th element of $(AB)^T$ is the same as the (i,j) -th element of $B^T A^T$. Note that:

$$\left((AB)^T\right)_{ij} = (AB)_{ji} = \sum_{k=1}^{\ell} a_{jk} b_{ki}$$

and

$$\left(B^T A^T\right)_{ij} = \sum_{k=1}^{\ell} b_{ik}^T a_{kj}^T = \sum_{k=1}^{\ell} b_{ki} a_{jk}.$$



Properties of Matrix Multiplication

Let A, B, C be appropriately sized matrices and r a scalar.

Properties of Matrix Multiplication

Let A, B, C be appropriately sized matrices and r a scalar.

We have some “good” properties:

Properties of Matrix Multiplication

Let A, B, C be appropriately sized matrices and r a scalar.

We have some “good” properties:

- ▶ $IA = AI = A.$

Properties of Matrix Multiplication

Let A, B, C be appropriately sized matrices and r a scalar.

We have some “good” properties:

- ▶ $IA = AI = A$. I is a multiplicative identity.
- ▶ $A(BC) = (AB)C$.

Properties of Matrix Multiplication

Let A, B, C be appropriately sized matrices and r a scalar.

We have some “good” properties:

- ▶ $IA = AI = A$. I is a multiplicative identity.
- ▶ $A(BC) = (AB)C$. Associative.

Properties of Matrix Multiplication

Let A, B, C be appropriately sized matrices and r a scalar.

We have some “good” properties:

- ▶ $IA = AI = A$. I is a multiplicative identity.
- ▶ $A(BC) = (AB)C$. Associative.
- ▶ $(rA)B = r(AB)$.

Properties of Matrix Multiplication

Let A, B, C be appropriately sized matrices and r a scalar.

We have some “good” properties:

- ▶ $IA = AI = A$. I is a multiplicative identity.
- ▶ $A(BC) = (AB)C$. Associative.
- ▶ $(rA)B = r(AB)$. Plays nice with scalar mult.

Properties of Matrix Multiplication

Let A, B, C be appropriately sized matrices and r a scalar.

We have some “good” properties:

- ▶ $IA = AI = A$. I is a multiplicative identity.
- ▶ $A(BC) = (AB)C$. Associative.
- ▶ $(rA)B = r(AB)$. Plays nice with scalar mult.
- ▶ $(A + B)C = AC + BC$ and $C(A + B) = CA + CB$.

Properties of Matrix Multiplication

Let A, B, C be appropriately sized matrices and r a scalar.

We have some “good” properties:

- ▶ $IA = AI = A$. I is a multiplicative identity.
- ▶ $A(BC) = (AB)C$. Associative.
- ▶ $(rA)B = r(AB)$. Plays nice with scalar mult.
- ▶ $(A + B)C = AC + BC$ and $C(A + B) = CA + CB$. Distributive.

Properties of Matrix Multiplication

Let A, B, C be appropriately sized matrices and r a scalar.

We have some “good” properties:

- ▶ $IA = AI = A$. I is a multiplicative identity.
- ▶ $A(BC) = (AB)C$. Associative.
- ▶ $(rA)B = r(AB)$. Plays nice with scalar mult.
- ▶ $(A + B)C = AC + BC$ and $C(A + B) = CA + CB$.
Distributive.

But why do we have two distributive laws?

Properties of Matrix Multiplication

Let A, B, C be appropriately sized matrices and r a scalar.

We have some “good” properties:

- ▶ $IA = AI = A$. I is a multiplicative identity.
- ▶ $A(BC) = (AB)C$. Associative.
- ▶ $(rA)B = r(AB)$. Plays nice with scalar mult.
- ▶ $(A + B)C = AC + BC$ and $C(A + B) = CA + CB$. Distributive.

But why do we have two distributive laws?

Because there are also some “not nice” properties:

- ▶ AB need not be equal to BA !

Properties of Matrix Multiplication

Let A, B, C be appropriately sized matrices and r a scalar.

We have some “good” properties:

- ▶ $IA = AI = A$. I is a multiplicative identity.
- ▶ $A(BC) = (AB)C$. Associative.
- ▶ $(rA)B = r(AB)$. Plays nice with scalar mult.
- ▶ $(A + B)C = AC + BC$ and $C(A + B) = CA + CB$.
Distributive.

But why do we have two distributive laws?

Because there are also some “not nice” properties:

- ▶ AB need not be equal to BA !
- ▶ $AB = 0$ does not mean that one of A or B is 0 !

Properties of Matrix Multiplication

Let A, B, C be appropriately sized matrices and r a scalar.

We have some “good” properties:

- ▶ $IA = AI = A$. I is a multiplicative identity.
- ▶ $A(BC) = (AB)C$. Associative.
- ▶ $(rA)B = r(AB)$. Plays nice with scalar mult.
- ▶ $(A + B)C = AC + BC$ and $C(A + B) = CA + CB$. Distributive.

But why do we have two distributive laws?

Because there are also some “not nice” properties:

- ▶ AB need not be equal to BA !
- ▶ $AB = 0$ does not mean that one of A or B is 0!
- ▶ $AB = AC$ even if $B \neq C$!

Properties of Matrix Multiplication

Let A, B, C be appropriately sized matrices and r a scalar.

We have some “good” properties:

- ▶ $IA = AI = A$. I is a multiplicative identity.
- ▶ $A(BC) = (AB)C$. Associative.
- ▶ $(rA)B = r(AB)$. Plays nice with scalar mult.
- ▶ $(A + B)C = AC + BC$ and $C(A + B) = CA + CB$. Distributive.

But why do we have two distributive laws?

Because there are also some “not nice” properties:

- ▶ AB need not be equal to BA !
- ▶ $AB = 0$ does not mean that one of A or B is 0!
- ▶ $AB = AC$ even if $B \neq C$!

The last two properties are a little troublesome! In general we cannot cancel/divide/etc.!

Why no cancellation rule makes things interesting!

What's the big deal about not being able to cancel?

Why no cancellation rule makes things interesting!

What's the big deal about not being able to cancel?

Recall that a linear system can be represented in the form

$$Ax = b$$

Where A and b are known and x is an unknown.

Why no cancellation rule makes things interesting!

What's the big deal about not being able to cancel?

Recall that a linear system can be represented in the form

$$Ax = b$$

Where A and b are known and x is an unknown. “Solving” the linear system is the same as finding the value of the vector x .

Why no cancellation rule makes things interesting!

What's the big deal about not being able to cancel?

Recall that a linear system can be represented in the form

$$Ax = b$$

Where A and b are known and x is an unknown. “Solving” the linear system is the same as finding the value of the vector x .

Now suppose we *know* a vector v so that

$$Av = b.$$

Why no cancellation rule makes things interesting!

What's the big deal about not being able to cancel?

Recall that a linear system can be represented in the form

$$Ax = b$$

Where A and b are known and x is an unknown. “Solving” the linear system is the same as finding the value of the vector x .

Now suppose we *know* a vector v so that

$$Av = b.$$

What we want to say is that

$$Ax = Av$$

Why no cancellation rule makes things interesting!

What's the big deal about not being able to cancel?

Recall that a linear system can be represented in the form

$$Ax = b$$

Where A and b are known and x is an unknown. “Solving” the linear system is the same as finding the value of the vector x .

Now suppose we *know* a vector v so that

$$Av = b.$$

What we want to say is that

$$Ax = Av \Rightarrow x = v!$$

So *if* we could cancel then we would know that $x = v$.

Why no cancellation rule makes things interesting!

What's the big deal about not being able to cancel?

Recall that a linear system can be represented in the form

$$Ax = b$$

Where A and b are known and x is an unknown. “Solving” the linear system is the same as finding the value of the vector x .

Now suppose we *know* a vector v so that

$$Av = b.$$

What we want to say is that

$$Ax = Av \Rightarrow x = v!$$

So *if* we could cancel then we would know that $x = v$. That is: a solution *exists* and is *unique*!

No cancellation \Leftrightarrow bad solution sets

The lack of a cancellation law is exactly the reason why linear systems can have no solution or an infinite number of solutions!
(And vice versa.)

Summary

- ▶ Matrix addition and scalar multiplication: Behaves well. “Vector space”.
- ▶ Transposition: Is novel. “Linear”.
- ▶ Multiplication: Has some issues. Be aware of them! No cancellation law! Makes “solving” an issue.
- ▶ Proving properties is not too rough. Reduce operation to the component definitions.