

# Exercises

Compute the following determinants directly from the definition:

$$\begin{vmatrix} 1 & 2 & 3 \\ 0 & 1 & 1 \\ 0 & 1 & 2 \end{vmatrix} \quad \text{and} \quad \begin{vmatrix} 1 & 0 & 0 \\ 2 & 1 & 1 \\ 3 & 1 & 2 \end{vmatrix}.$$

# Exercises

Compute the following determinants directly from the definition:

$$\begin{vmatrix} 1 & 2 & 3 \\ 0 & 1 & 1 \\ 0 & 1 & 2 \end{vmatrix} \quad \text{and} \quad \begin{vmatrix} 1 & 0 & 0 \\ 2 & 1 & 1 \\ 3 & 1 & 2 \end{vmatrix}.$$

Result: They are both 1.

# Exercises

Compute the following determinants directly from the definition:

$$\begin{vmatrix} 1 & 2 & 3 \\ 0 & 1 & 1 \\ 0 & 1 & 2 \end{vmatrix} \quad \text{and} \quad \begin{vmatrix} 1 & 0 & 0 \\ 2 & 1 & 1 \\ 3 & 1 & 2 \end{vmatrix}.$$

Result: They are both 1. Note that the matrices are transposes.

# Determinant Definition

Recall that the determinant is a function:

$$\det : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$$

so that

1.  $\det(I) = 1$ .
2.  $\det$  is *antisymmetric* on the rows of  $A$ .
3.  $\det$  is *linear* on each row of  $A$ .

# Determinant Definition

Recall that the determinant is a function:

$$\det : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$$

so that

1.  $\det(I) = 1$ .
2.  $\det$  is *antisymmetric* on the rows of  $A$ .
3.  $\det$  is *linear* on each row of  $A$ .

These properties can be applied to the elementary row operations:

Old Matrix	Operation	New Matrix	Det Prop
$A$	Switch two rows of $A$	$B$	$ B  = - A $
$A$	Scale one row of $A$ by $k$	$B$	$ B  = k A $
$A$	Add/replace rows of $A$	$B$	$ B  =  A $

# Immediate Results:

## Theorem

*Let  $A$  be an  $n \times n$  matrix then*

- 1. If  $A$  is diagonal, upper triangular, or lower triangular then*

$$\det(A) = \prod_{i=1}^n a_{ii}.$$

# Immediate Results:

## Theorem

*Let  $A$  be an  $n \times n$  matrix then*

- 1. If  $A$  is diagonal, upper triangular, or lower triangular then*

$$\det(A) = \prod_{i=1}^n a_{ii}.$$

- 2. If two rows of  $A$  are the same, then  $\det(A) = 0$ .*

# Immediate Results:

## Theorem

Let  $A$  be an  $n \times n$  matrix then

1. If  $A$  is diagonal, upper triangular, or lower triangular then

$$\det(A) = \prod_{i=1}^n a_{ii}.$$

2. If two rows of  $A$  are the same, then  $\det(A) = 0$ .
  - 2.1 If  $A$  has a row of zeros, then  $|A| = 0$ .



# Immediate Results:

## Theorem

Let  $A$  be an  $n \times n$  matrix then

1. If  $A$  is diagonal, upper triangular, or lower triangular then

$$\det(A) = \prod_{i=1}^n a_{ii}.$$

2. If two rows of  $A$  are the same, then  $\det(A) = 0$ .
  - 2.1 If  $A$  has a row of zeros, then  $|A| = 0$ .
  - 2.2 If  $A$  is singular then  $|A| = 0$ .

# Immediate Results:

## Theorem

Let  $A$  be an  $n \times n$  matrix then

1. If  $A$  is diagonal, upper triangular, or lower triangular then

$$\det(A) = \prod_{i=1}^n a_{ii}.$$

2. If two rows of  $A$  are the same, then  $\det(A) = 0$ .

2.1 If  $A$  has a row of zeros, then  $|A| = 0$ .

2.2 If  $A$  is singular then  $|A| = 0$ .

(1) tells us that if we row reduce  $A$  to a triangular form, we can easily compute the determinant.

# Immediate Results:

## Theorem

Let  $A$  be an  $n \times n$  matrix then

1. If  $A$  is diagonal, upper triangular, or lower triangular then

$$\det(A) = \prod_{i=1}^n a_{ii}.$$

2. If two rows of  $A$  are the same, then  $\det(A) = 0$ .

2.1 If  $A$  has a row of zeros, then  $|A| = 0$ .

2.2 If  $A$  is singular then  $|A| = 0$ .

(1) tells us that if we row reduce  $A$  to a triangular form, we can easily compute the determinant.

One goal today is to strengthen (2.2) into an 'if and only if' statement.

# Determinants and Elementary Matrices

Recall that each elementary row operation can be represented by *left* multiplication by some matrix  $E$ :

Operation	Example $E$	Determinant
-----------	-------------	-------------

# Determinants and Elementary Matrices

Recall that each elementary row operation can be represented by *left* multiplication by some matrix  $E$ :

Operation	Example $E$	Determinant
Switch Rows	$E = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$ E  = -1$

# Determinants and Elementary Matrices

Recall that each elementary row operation can be represented by *left* multiplication by some matrix  $E$ :

Operation	Example $E$	Determinant
Switch Rows	$E = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$ E  = -1$
Scale row	$E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$ E  = k$

# Determinants and Elementary Matrices

Recall that each elementary row operation can be represented by *left* multiplication by some matrix  $E$ :

Operation	Example $E$	Determinant
Switch Rows	$E = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$ E  = -1$
Scale row	$E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$ E  = k$
Add rows	$E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$	$ E  = 1$

# Determinants and Elementary Matrices

Recall that each elementary row operation can be represented by *left* multiplication by some matrix  $E$ :

Operation	Example $E$	Determinant
Switch Rows	$E = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$ E  = -1$
Scale row	$E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & 1 \end{bmatrix}$	$ E  = k$
Add rows	$E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$	$ E  = 1$

## Lemma

*Let  $A$  be a matrix and  $E$  be an elementary matrix. Then*

$$\det(EA) = \det(E) \det(A).$$



# Result from Lemma

## Theorem

*A is non-singular*  $\Rightarrow \det(A) \neq 0$ .

# Result from Lemma

## Theorem

*A is non-singular*  $\Rightarrow \det(A) \neq 0$ .

## Proof.

Suppose *A* is non-singular.

# Result from Lemma

## Theorem

*A is non-singular*  $\Rightarrow \det(A) \neq 0$ .

## Proof.

Suppose  $A$  is non-singular. Then there exist elementary row operation matrices  $E_i$  so that

$$I = E_1 E_2 E_3 \cdots E_n A$$

# Result from Lemma

## Theorem

*A is non-singular*  $\Rightarrow \det(A) \neq 0$ .

## Proof.

Suppose  $A$  is non-singular. Then there exist elementary row operation matrices  $E_i$  so that

$$\begin{aligned} I &= E_1 E_2 E_3 \cdots E_n A \\ \det(I) &= \det(E_1 \cdots E_n A) \end{aligned}$$

# Result from Lemma

## Theorem

*A is non-singular*  $\Rightarrow \det(A) \neq 0$ .

## Proof.

Suppose  $A$  is non-singular. Then there exist elementary row operation matrices  $E_i$  so that

$$\begin{aligned} I &= E_1 E_2 E_3 \cdots E_n A \\ \det(I) &= \det(E_1 \cdots E_n A) \\ 1 &= \det(E_1) \cdots \det(E_n) \det(A) \end{aligned}$$

# Result from Lemma

## Theorem

*A is non-singular*  $\Rightarrow \det(A) \neq 0$ .

## Proof.

Suppose  $A$  is non-singular. Then there exist elementary row operation matrices  $E_i$  so that

$$\begin{aligned} I &= E_1 E_2 E_3 \cdots E_n A \\ \det(I) &= \det(E_1 \cdots E_n A) \\ 1 &= \det(E_1) \cdots \det(E_n) \det(A) \quad (\text{from lemma}) \end{aligned}$$

# Result from Lemma

## Theorem

*A is non-singular*  $\Rightarrow \det(A) \neq 0$ .

## Proof.

Suppose  $A$  is non-singular. Then there exist elementary row operation matrices  $E_i$  so that

$$\begin{aligned} I &= E_1 E_2 E_3 \cdots E_n A \\ \det(I) &= \det(E_1 \cdots E_n A) \\ 1 &= \det(E_1) \cdots \det(E_n) \det(A) \quad (\text{from lemma}) \\ 1 &= \underbrace{\det(E_1) \cdots \det(E_n)}_{\text{all non-zero}} \det(A) \end{aligned}$$

# Result from Lemma

## Theorem

*A is non-singular*  $\Rightarrow \det(A) \neq 0$ .

## Proof.

Suppose  $A$  is non-singular. Then there exist elementary row operation matrices  $E_i$  so that

$$\begin{aligned} I &= E_1 E_2 E_3 \cdots E_n A \\ \det(I) &= \det(E_1 \cdots E_n A) \\ 1 &= \det(E_1) \cdots \det(E_n) \det(A) \quad (\text{from lemma}) \\ 1 &= \underbrace{\det(E_1) \cdots \det(E_n)}_{\text{all non-zero}} \det(A) \end{aligned}$$

Therefore  $\det(A) \neq 0$ .





$$A \text{ singular} \Leftrightarrow |A| = 0$$

Combining our results from the last lecture and today, we have the following:

$$A \text{ singular} \Leftrightarrow |A| = 0$$

Combining our results from the last lecture and today, we have the following:

$$A \text{ singular} \Leftrightarrow |A| = 0.$$

$$A \text{ singular} \Leftrightarrow |A| = 0$$

Combining our results from the last lecture and today, we have the following:

$$A \text{ singular} \Leftrightarrow |A| = 0.$$

(or  $A$  non-singular  $\Leftrightarrow |A| \neq 0$ .)

$$A \text{ singular} \Leftrightarrow |A| = 0$$

Combining our results from the last lecture and today, we have the following:

$$A \text{ singular} \Leftrightarrow |A| = 0.$$

(or  $A$  non-singular  $\Leftrightarrow |A| \neq 0$ .)

This has direct bearing on solving systems of equations  $Ax = b$ :

- ▶ If  $|A| \neq 0$  then is  $Ax = b$  consistent? What types of solutions?

$$A \text{ singular} \Leftrightarrow |A| = 0$$

Combining our results from the last lecture and today, we have the following:

$$A \text{ singular} \Leftrightarrow |A| = 0.$$

(or  $A$  non-singular  $\Leftrightarrow |A| \neq 0$ .)

This has direct bearing on solving systems of equations  $Ax = b$ :

- ▶ If  $|A| \neq 0$  then is  $Ax = b$  consistent? What types of solutions?
  - ▶ Yes, with unique solutions!

$$A \text{ singular} \Leftrightarrow |A| = 0$$

Combining our results from the last lecture and today, we have the following:

$$A \text{ singular} \Leftrightarrow |A| = 0.$$

(or  $A$  non-singular  $\Leftrightarrow |A| \neq 0$ .)

This has direct bearing on solving systems of equations  $Ax = b$ :

- ▶ If  $|A| \neq 0$  then is  $Ax = b$  consistent? What types of solutions?
  - ▶ Yes, with unique solutions!
- ▶ If  $|A| = 0$  then is  $Ax = b$  consistent? What types of solutions?

# $A$ singular $\Leftrightarrow |A| = 0$

Combining our results from the last lecture and today, we have the following:

$$A \text{ singular} \Leftrightarrow |A| = 0.$$
$$(\text{or } A \text{ non-singular} \Leftrightarrow |A| \neq 0.)$$

This has direct bearing on solving systems of equations  $Ax = b$ :

- ▶ If  $|A| \neq 0$  then is  $Ax = b$  consistent? What types of solutions?
  - ▶ Yes, with unique solutions!
- ▶ If  $|A| = 0$  then is  $Ax = b$  consistent? What types of solutions?
  - ▶ Possibly inconsistent. If consistent, has infinite solutions.

# $A$ singular $\Leftrightarrow |A| = 0$

Combining our results from the last lecture and today, we have the following:

$$A \text{ singular} \Leftrightarrow |A| = 0.$$

(or  $A$  non-singular  $\Leftrightarrow |A| \neq 0$ .)

This has direct bearing on solving systems of equations  $Ax = b$ :

- ▶ If  $|A| \neq 0$  then is  $Ax = b$  consistent? What types of solutions?
  - ▶ Yes, with unique solutions!
- ▶ If  $|A| = 0$  then is  $Ax = b$  consistent? What types of solutions?
  - ▶ Possibly inconsistent. If consistent, has infinite solutions.
- ▶ If  $|A| = 0$  then is  $Ax = 0$  consistent? What types of solutions?



# $A$ singular $\Leftrightarrow |A| = 0$

Combining our results from the last lecture and today, we have the following:

$$A \text{ singular} \Leftrightarrow |A| = 0.$$

(or  $A$  non-singular  $\Leftrightarrow |A| \neq 0$ .)

This has direct bearing on solving systems of equations  $Ax = b$ :

- ▶ If  $|A| \neq 0$  then is  $Ax = b$  consistent? What types of solutions?
  - ▶ Yes, with unique solutions!
- ▶ If  $|A| = 0$  then is  $Ax = b$  consistent? What types of solutions?
  - ▶ Possibly inconsistent. If consistent, has infinite solutions.
- ▶ If  $|A| = 0$  then is  $Ax = 0$  consistent? What types of solutions?
  - ▶ Consistent with infinite solutions.

## Lemma Result (2)

### Theorem

*Let  $A$  and  $B$  be  $n \times n$  matrices. Then*

$$\det(AB) = \det(A) \det(B)$$

## Lemma Result (2)

### Theorem

*Let  $A$  and  $B$  be  $n \times n$  matrices. Then*

$$\det(AB) = \det(A) \det(B)$$

### Proof.

**Case 1**  $A$  is singular. Then so is  $AB$ .

## Lemma Result (2)

### Theorem

*Let  $A$  and  $B$  be  $n \times n$  matrices. Then*

$$\det(AB) = \det(A) \det(B)$$

### Proof.

**Case 1**  $A$  is singular. Then so is  $AB$ . The result reduces to  $0 = 0$  which is true.

## Lemma Result (2)

### Theorem

*Let  $A$  and  $B$  be  $n \times n$  matrices. Then*

$$\det(AB) = \det(A) \det(B)$$

### Proof.

**Case 1**  $A$  is singular. Then so is  $AB$ . The result reduces to  $0 = 0$  which is true.

**Case 2**  $A$  is non-singular.

## Lemma Result (2)

### Theorem

*Let  $A$  and  $B$  be  $n \times n$  matrices. Then*

$$\det(AB) = \det(A) \det(B)$$

### Proof.

**Case 1**  $A$  is singular. Then so is  $AB$ . The result reduces to  $0 = 0$  which is true.

**Case 2**  $A$  is non-singular. Then  $A = E_1 \cdots E_n$ .

## Lemma Result (2)

### Theorem

Let  $A$  and  $B$  be  $n \times n$  matrices. Then

$$\det(AB) = \det(A) \det(B)$$

### Proof.

**Case 1**  $A$  is singular. Then so is  $AB$ . The result reduces to  $0 = 0$  which is true.

**Case 2**  $A$  is non-singular. Then  $A = E_1 \cdots E_n$ . So

$$\det(AB) = \det(E_1 \cdots E_n B)$$

## Lemma Result (2)

### Theorem

Let  $A$  and  $B$  be  $n \times n$  matrices. Then

$$\det(AB) = \det(A) \det(B)$$

### Proof.

**Case 1**  $A$  is singular. Then so is  $AB$ . The result reduces to  $0 = 0$  which is true.

**Case 2**  $A$  is non-singular. Then  $A = E_1 \cdots E_n$ . So

$$\begin{aligned} \det(AB) &= \det(E_1 \cdots E_n B) \\ &= \det(E_1) \cdots \det(E_n) \det(B) \end{aligned}$$



## Lemma Result (2)

### Theorem

Let  $A$  and  $B$  be  $n \times n$  matrices. Then

$$\det(AB) = \det(A) \det(B)$$

### Proof.

**Case 1**  $A$  is singular. Then so is  $AB$ . The result reduces to  $0 = 0$  which is true.

**Case 2**  $A$  is non-singular. Then  $A = E_1 \cdots E_n$ . So

$$\begin{aligned} \det(AB) &= \det(E_1 \cdots E_n B) \\ &= \det(E_1) \cdots \det(E_n) \det(B) \quad (\text{from lemma}) \end{aligned}$$

## Lemma Result (2)

### Theorem

Let  $A$  and  $B$  be  $n \times n$  matrices. Then

$$\det(AB) = \det(A) \det(B)$$

### Proof.

**Case 1**  $A$  is singular. Then so is  $AB$ . The result reduces to  $0 = 0$  which is true.

**Case 2**  $A$  is non-singular. Then  $A = E_1 \cdots E_n$ . So

$$\begin{aligned} \det(AB) &= \det(E_1 \cdots E_n B) \\ &= \det(E_1) \cdots \det(E_n) \det(B) \quad (\text{from lemma}) \\ &= \det(E_1 \cdots E_n) \det(B) \end{aligned}$$

## Lemma Result (2)

### Theorem

Let  $A$  and  $B$  be  $n \times n$  matrices. Then

$$\det(AB) = \det(A) \det(B)$$

### Proof.

**Case 1**  $A$  is singular. Then so is  $AB$ . The result reduces to  $0 = 0$  which is true.

**Case 2**  $A$  is non-singular. Then  $A = E_1 \cdots E_n$ . So

$$\begin{aligned} \det(AB) &= \det(E_1 \cdots E_n B) \\ &= \det(E_1) \cdots \det(E_n) \det(B) \quad (\text{from lemma}) \\ &= \det(E_1 \cdots E_n) \det(B) \quad (\text{lemma again}) \end{aligned}$$

## Lemma Result (2)

### Theorem

Let  $A$  and  $B$  be  $n \times n$  matrices. Then

$$\det(AB) = \det(A) \det(B)$$

### Proof.

**Case 1**  $A$  is singular. Then so is  $AB$ . The result reduces to  $0 = 0$  which is true.

**Case 2**  $A$  is non-singular. Then  $A = E_1 \cdots E_n$ . So

$$\begin{aligned} \det(AB) &= \det(E_1 \cdots E_n B) \\ &= \det(E_1) \cdots \det(E_n) \det(B) \quad (\text{from lemma}) \\ &= \det(E_1 \cdots E_n) \det(B) \quad (\text{lemma again}) \\ &= \det(A) \det(B). \end{aligned}$$

$$|AB| = |A||B|$$

Some nice applications of this result:

- ▶  $\det(AB) = \det(BA)$ .

$$|AB| = |A||B|$$

Some nice applications of this result:

- ▶  $\det(AB) = \det(BA)$ .
  - ▶ Since  $\det(AB) = \det(A)\det(B) = \det(B)\det(A) = \det(BA)$ .

$$|AB| = |A||B|$$

Some nice applications of this result:

- ▶  $\det(AB) = \det(BA)$ .
  - ▶ Since  $\det(AB) = \det(A)\det(B) = \det(B)\det(A) = \det(BA)$ .
  - ▶ Note: If  $B = E$  is an elementary operation, then we have *column operations effect the value of the determinant the same as row operations!*

$$|AB| = |A||B|$$

Some nice applications of this result:

- ▶  $\det(AB) = \det(BA)$ .
  - ▶ Since  $\det(AB) = \det(A)\det(B) = \det(B)\det(A) = \det(BA)$ .
  - ▶ Note: If  $B = E$  is an elementary operation, then we have *column operations effect the value of the determinant the same as row operations!*
- ▶  $\det(A) = \det(A^T)$ .



$$|AB| = |A||B|$$

Some nice applications of this result:

- ▶  $\det(AB) = \det(BA)$ .
  - ▶ Since  $\det(AB) = \det(A)\det(B) = \det(B)\det(A) = \det(BA)$ .
  - ▶ Note: If  $B = E$  is an elementary operation, then we have *column operations effect the value of the determinant the same as row operations!*
- ▶  $\det(A) = \det(A^T)$ .
  - ▶ To prove, write  $A = E_1 \cdots E_n$  and take the transpose of both sides.

$$|AB| = |A||B|$$

Some nice applications of this result:

- ▶  $\det(AB) = \det(BA)$ .
  - ▶ Since  $\det(AB) = \det(A)\det(B) = \det(B)\det(A) = \det(BA)$ .
  - ▶ Note: If  $B = E$  is an elementary operation, then we have *column operations effect the value of the determinant the same as row operations!*
- ▶  $\det(A) = \det(A^T)$ .
  - ▶ To prove, write  $A = E_1 \cdots E_n$  and take the transpose of both sides.
- ▶  $|A^{-1}| = |A|^{-1}$ .

$$|AB| = |A||B|$$

Some nice applications of this result:

- ▶  $\det(AB) = \det(BA)$ .
  - ▶ Since  $\det(AB) = \det(A)\det(B) = \det(B)\det(A) = \det(BA)$ .
  - ▶ Note: If  $B = E$  is an elementary operation, then we have *column operations effect the value of the determinant the same as row operations!*
- ▶  $\det(A) = \det(A^T)$ .
  - ▶ To prove, write  $A = E_1 \cdots E_n$  and take the transpose of both sides.
- ▶  $|A^{-1}| = |A|^{-1}$ .
  - ▶ Exercise: Prove this using  $AA^{-1} = I$ .

# Final Application of the lemma

# Final Application of the lemma

## Theorem

*The determinant is unique!*

# Final Application of the lemma

## Theorem

*The determinant is unique!*

How do we prove this?

# Final Application of the lemma

## Theorem

*The determinant is unique!*

How do we prove this?

Suppose there are two functions  $(\det_1)$  and  $(\det_2)$  which satisfy all the properties of the determinant.

# Final Application of the lemma

## Theorem

*The determinant is unique!*

How do we prove this?

Suppose there are two functions  $(\det_1)$  and  $(\det_2)$  which satisfy all the properties of the determinant. We need to show that  $\det_1(A) = \det_2(A)$  for every matrix  $A$ .



# Proof

Proof.

Case 1  $A$  is singular.

# Proof

Proof.

Case 1  $A$  is singular. Then  $\det_1(A) = \det_2(A) = 0$ .

# Proof

Proof.

Case 1  $A$  is singular. Then  $\det_1(A) = \det_2(A) = 0$ . So the two functions agree on all singular matrices.

Case 2  $A$  is non-singular.

# Proof

Proof.

**Case 1**  $A$  is singular. Then  $\det_1(A) = \det_2(A) = 0$ . So the two functions agree on all singular matrices.

**Case 2**  $A$  is non-singular. Then  $A = E_1 \cdots E_n$  for some elementary matrices  $E_i$ .

# Proof

Proof.

**Case 1**  $A$  is singular. Then  $\det_1(A) = \det_2(A) = 0$ . So the two functions agree on all singular matrices.

**Case 2**  $A$  is non-singular. Then  $A = E_1 \cdots E_n$  for some elementary matrices  $E_i$ . We have  $\det_1(E_i) = \det_2(E_i)$ .

# Proof

Proof.

**Case 1**  $A$  is singular. Then  $\det_1(A) = \det_2(A) = 0$ . So the two functions agree on all singular matrices.

**Case 2**  $A$  is non-singular. Then  $A = E_1 \cdots E_n$  for some elementary matrices  $E_i$ . We have  $\det_1(E_i) = \det_2(E_i)$ . Therefore:

$$\begin{aligned}\det_1(A) &= \det_1(E_1 \cdots E_n) \\ &= \det_1(E_1) \cdots \det_1(E_n) \\ &= \det_2(E_1) \cdots \det_2(E_n) \\ &= \det_2(E_1 \cdots E_n) \\ &= \det_2(A)\end{aligned}$$



# Why Uniqueness is Important

Now we know: if a computation satisfies all of the properties of the determinant, then it must be the determinant!

# Why Uniqueness is Important

Now we know: if a computation satisfies all of the properties of the determinant, then it must be the determinant!

Next: We will look at a computational interpretation for the determinant!