

# Review: Matrix Transformation Maps

Let  $A \in \mathbb{R}^{m \times n}$ . Then we can construct a map:

$$T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m \quad \text{by} \quad T_A(x) = Ax.$$

# Review: Matrix Transformation Maps

Let  $A \in \mathbb{R}^{m \times n}$ . Then we can construct a map:

$$T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m \quad \text{by} \quad T_A(x) = Ax.$$

Question: How does the (non)singularity of  $A$  affect the map  $T_A$ ?

# Review: Matrix Transformation Maps

Let  $A \in \mathbb{R}^{m \times n}$ . Then we can construct a map:

$$T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m \quad \text{by} \quad T_A(x) = Ax.$$

Question: How does the (non)singularity of  $A$  affect the map  $T_A$ ?

This question is closely related to whether or not  $T_A$  is an isomorphism!

# Review: Matrix Transformation Maps

Let  $A \in \mathbb{R}^{m \times n}$ . Then we can construct a map:

$$T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m \quad \text{by} \quad T_A(x) = Ax.$$

Question: How does the (non)singularity of  $A$  affect the map  $T_A$ ?

This question is closely related to whether or not  $T_A$  is an isomorphism! (that is: 1-1 and onto).

# Review: Matrix Transformation Maps

Let  $A \in \mathbb{R}^{m \times n}$ . Then we can construct a map:

$$T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m \quad \text{by} \quad T_A(x) = Ax.$$

Question: How does the (non)singularity of  $A$  affect the map  $T_A$ ?

This question is closely related to whether or not  $T_A$  is an isomorphism! (that is: 1-1 and onto).

In order to study these questions we will investigate two sets:

# Review: Matrix Transformation Maps

Let  $A \in \mathbb{R}^{m \times n}$ . Then we can construct a map:

$$T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m \quad \text{by} \quad T_A(x) = Ax.$$

Question: How does the (non)singularity of  $A$  affect the map  $T_A$ ?

This question is closely related to whether or not  $T_A$  is an isomorphism! (that is: 1-1 and onto).

In order to study these questions we will investigate two sets:

- The collection of vectors in  $\mathbb{R}^n$  which map to zero.

# Review: Matrix Transformation Maps

Let  $A \in \mathbb{R}^{m \times n}$ . Then we can construct a map:

$$T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m \quad \text{by} \quad T_A(x) = Ax.$$

Question: How does the (non)singularity of  $A$  affect the map  $T_A$ ?

This question is closely related to whether or not  $T_A$  is an isomorphism! (that is: 1-1 and onto).

In order to study these questions we will investigate two sets:

- ▶ The collection of vectors in  $\mathbb{R}^n$  which map to zero.
  - ▶ We will see how this tells us if the map is 1-1.

# Review: Matrix Transformation Maps

Let  $A \in \mathbb{R}^{m \times n}$ . Then we can construct a map:

$$T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m \quad \text{by} \quad T_A(x) = Ax.$$

Question: How does the (non)singularity of  $A$  affect the map  $T_A$ ?

This question is closely related to whether or not  $T_A$  is an isomorphism! (that is: 1-1 and onto).

In order to study these questions we will investigate two sets:

- ▶ The collection of vectors in  $\mathbb{R}^n$  which map to zero.
  - ▶ We will see how this tells us if the map is 1-1.
- ▶ The collection of vectors in  $\mathbb{R}^m$  which are mapped to.



# Review: Matrix Transformation Maps

Let  $A \in \mathbb{R}^{m \times n}$ . Then we can construct a map:

$$T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m \quad \text{by} \quad T_A(x) = Ax.$$

Question: How does the (non)singularity of  $A$  affect the map  $T_A$ ?

This question is closely related to whether or not  $T_A$  is an isomorphism! (that is: 1-1 and onto).

In order to study these questions we will investigate two sets:

- ▶ The collection of vectors in  $\mathbb{R}^n$  which map to zero.
  - ▶ We will see how this tells us if the map is 1-1.
- ▶ The collection of vectors in  $\mathbb{R}^m$  which are mapped to.
  - ▶ This will help us learn if the map is onto.

# Null Space/Kernel

As before, let  $A$  be an  $m \times n$  matrix and  $T_A$  its associated transformation.

# Null Space/Kernel

As before, let  $A$  be an  $m \times n$  matrix and  $T_A$  its associated transformation.

## Definition

The **null space** of  $A$  is the set

$$\text{null}(A) = \{x \in \mathbb{R}^n : Ax = 0\}.$$

# Null Space/Kernel

As before, let  $A$  be an  $m \times n$  matrix and  $T_A$  its associated transformation.

## Definition

The **null space** of  $A$  is the set

$$\text{null}(A) = \{x \in \mathbb{R}^n : Ax = 0\}.$$

The **kernel** of  $T_A$  is the set

$$\ker(T_A) = \{x \in \mathbb{R}^n : T_A(x) = 0\}$$

# Null Space/Kernel

As before, let  $A$  be an  $m \times n$  matrix and  $T_A$  its associated transformation.

## Definition

The **null space** of  $A$  is the set

$$\text{null}(A) = \{x \in \mathbb{R}^n : Ax = 0\}.$$

The **kernel** of  $T_A$  is the set

$$\ker(T_A) = \{x \in \mathbb{R}^n : T_A(x) = 0\}$$

It should be clear that

$$\ker(T_A) = \text{null}(A).$$

# Null Space/Kernel

As before, let  $A$  be an  $m \times n$  matrix and  $T_A$  its associated transformation.

## Definition

The **null space** of  $A$  is the set

$$\text{null}(A) = \{x \in \mathbb{R}^n : Ax = 0\}.$$

The **kernel** of  $T_A$  is the set

$$\ker(T_A) = \{x \in \mathbb{R}^n : T_A(x) = 0\}$$

It should be clear that

$$\ker(T_A) = \text{null}(A).$$

*Kernel* is the more robust definition and we will need it later. For now, we will work only with the null space.

# Null Space = Homogeneous slns!

This is not the first time that we have met the null space of a matrix, though.

# Null Space = Homogeneous slns!

This is not the first time that we have met the null space of a matrix, though.

The null space of  $A$  is exactly the set of solutions to  $Ax = 0$ .



# Null Space = Homogeneous slns!

This is not the first time that we have met the null space of a matrix, though.

The null space of  $A$  is exactly the set of solutions to  $Ax = 0$ .  
Therefore the null space is the set of solutions to the homogeneous linear problem!

# Null Space = Homogeneous slns!

This is not the first time that we have met the null space of a matrix, though.

The null space of  $A$  is exactly the set of solutions to  $Ax = 0$ . Therefore the null space is the set of solutions to the homogeneous linear problem! In particular: it is a subspace of  $\mathbb{R}^n$  and therefore has a basis!

## Exercise

Find the null space of the following matrix. In particular, express it as the span of appropriate basis vectors.

$$\begin{bmatrix} 1 & -2 & 0 & 3 & -1 \\ 3 & 2 & 8 & 1 & 4 \\ 2 & 3 & 7 & 2 & 3 \\ -1 & 2 & 0 & 4 & -3 \end{bmatrix}$$

## Exercise

Find the null space of the following matrix. In particular, express it as the span of appropriate basis vectors.

$$\begin{bmatrix} 1 & -2 & 0 & 3 & -1 \\ 3 & 2 & 8 & 1 & 4 \\ 2 & 3 & 7 & 2 & 3 \\ -1 & 2 & 0 & 4 & -3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

## Exercise

Find the null space of the following matrix. In particular, express it as the span of appropriate basis vectors.

$$\begin{bmatrix} 1 & -2 & 0 & 3 & -1 \\ 3 & 2 & 8 & 1 & 4 \\ 2 & 3 & 7 & 2 & 3 \\ -1 & 2 & 0 & 4 & -3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Solution:

## Exercise

Find the null space of the following matrix. In particular, express it as the span of appropriate basis vectors.

$$\begin{bmatrix} 1 & -2 & 0 & 3 & -1 \\ 3 & 2 & 8 & 1 & 4 \\ 2 & 3 & 7 & 2 & 3 \\ -1 & 2 & 0 & 4 & -3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Solution:

$$a = -2c - e$$

$$b = -c - e$$

$$c = c$$

$$d = e$$

$$e = e$$

## Exercise

Find the null space of the following matrix. In particular, express it as the span of appropriate basis vectors.

$$\begin{bmatrix} 1 & -2 & 0 & 3 & -1 \\ 3 & 2 & 8 & 1 & 4 \\ 2 & 3 & 7 & 2 & 3 \\ -1 & 2 & 0 & 4 & -3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Solution:

$$\begin{aligned} a &= -2c - e \\ b &= -c - e \\ c &= c \\ d &= e \\ e &= e \end{aligned} \quad \Leftrightarrow \quad \text{null}(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ -1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ -1 \\ 0 \\ 1 \\ 1 \end{bmatrix} \right\}.$$

## Exercise

Find the null space of the following matrix. In particular, express it as the span of appropriate basis vectors.

$$\begin{bmatrix} 1 & -2 & 0 & 3 & -1 \\ 3 & 2 & 8 & 1 & 4 \\ 2 & 3 & 7 & 2 & 3 \\ -1 & 2 & 0 & 4 & -3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Solution:

$$\begin{aligned} a &= -2c - e \\ b &= -c - e \\ c &= c \\ d &= e \\ e &= e \end{aligned} \quad \Leftrightarrow \quad \text{null}(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ -1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ -1 \\ 0 \\ 1 \\ 1 \end{bmatrix} \right\}.$$

It has dimension 2.



# Column Space/Range

Again, let  $A$  be an  $m \times n$  matrix and  $T_A$  its associated transformation.

# Column Space/Range

Again, let  $A$  be an  $m \times n$  matrix and  $T_A$  its associated transformation.

## Definition

The **column space** of  $A$  is the set of all possible linear combinations of the columns of the matrix  $A$ :

# Column Space/Range

Again, let  $A$  be an  $m \times n$  matrix and  $T_A$  its associated transformation.

## Definition

The **column space** of  $A$  is the set of all possible linear combinations of the columns of the matrix  $A$ :

$$\text{col}(A) = \left\{ \sum_{i=1}^n x_i a_i : x_i \in \mathbb{R}, A = [a_1 \ a_2 \ \cdots \ a_n] \right\}.$$

# Column Space/Range

Again, let  $A$  be an  $m \times n$  matrix and  $T_A$  its associated transformation.

## Definition

The **column space** of  $A$  is the set of all possible linear combinations of the columns of the matrix  $A$ :

$$\text{col}(A) = \left\{ \sum_{i=1}^n x_i a_i : x_i \in \mathbb{R}, A = [a_1 \ a_2 \ \cdots \ a_n] \right\}.$$

Note that this is a subset of  $\mathbb{R}^m$ .

# Column Space/Range

Again, let  $A$  be an  $m \times n$  matrix and  $T_A$  its associated transformation.

## Definition

The **column space** of  $A$  is the set of all possible linear combinations of the columns of the matrix  $A$ :

$$\text{col}(A) = \left\{ \sum_{i=1}^n x_i a_i : x_i \in \mathbb{R}, A = [a_1 \ a_2 \ \cdots \ a_n] \right\}.$$

Note that this is a subset of  $\mathbb{R}^m$ .

The **range** of  $T_A$  is the set of elements in  $\mathbb{R}^m$  which are mapped to:

# Column Space/Range

Again, let  $A$  be an  $m \times n$  matrix and  $T_A$  its associated transformation.

## Definition

The **column space** of  $A$  is the set of all possible linear combinations of the columns of the matrix  $A$ :

$$\text{col}(A) = \left\{ \sum_{i=1}^n x_i a_i : x_i \in \mathbb{R}, A = [a_1 \ a_2 \ \cdots \ a_n] \right\}.$$

Note that this is a subset of  $\mathbb{R}^m$ .

The **range** of  $T_A$  is the set of elements in  $\mathbb{R}^m$  which are mapped to:

$$\text{range}(T_A) = \{y \in \mathbb{R}^m : \text{there is an } x \in \mathbb{R}^n \text{ so that } T_A(x) = y\}$$

# Column Space of $A$ is Range of $T_A$

To see this, just recall that:

$$\sum_{i=1}^n x_i a_i = \begin{bmatrix} a_1 & a_2 & \cdots & a_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = Ax$$

## Column Space of $A$ is Range of $T_A$

To see this, just recall that:

$$\sum_{i=1}^n x_i a_i = \begin{bmatrix} a_1 & a_2 & \cdots & a_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = Ax$$

As before: range is the more general object, but for now we concentrate on column space.



## Column Space of $A$ is Range of $T_A$

To see this, just recall that:

$$\sum_{i=1}^n x_i a_i = \begin{bmatrix} a_1 & a_2 & \cdots & a_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = Ax$$

As before: range is the more general object, but for now we concentrate on column space.

Since the column space of a matrix is the span of the columns, we know that it must be a subspace of  $\mathbb{R}^m$ .

## Column Space of $A$ is Range of $T_A$

To see this, just recall that:

$$\sum_{i=1}^n x_i a_i = \begin{bmatrix} a_1 & a_2 & \cdots & a_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = Ax$$

As before: range is the more general object, but for now we concentrate on column space.

Since the column space of a matrix is the span of the columns, we know that it must be a subspace of  $\mathbb{R}^m$ . Therefore it has a basis!

## Column Space of $A$ is Range of $T_A$

To see this, just recall that:

$$\sum_{i=1}^n x_i a_i = \begin{bmatrix} a_1 & a_2 & \cdots & a_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = Ax$$

As before: range is the more general object, but for now we concentrate on column space.

Since the column space of a matrix is the span of the columns, we know that it must be a subspace of  $\mathbb{R}^m$ . Therefore it has a basis!

In fact, we already know how to find the basis for  $\text{col}(A)$ !

## Column Space of $A$ is Range of $T_A$

To see this, just recall that:

$$\sum_{i=1}^n x_i a_i = \begin{bmatrix} a_1 & a_2 & \cdots & a_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = Ax$$

As before: range is the more general object, but for now we concentrate on column space.

Since the column space of a matrix is the span of the columns, we know that it must be a subspace of  $\mathbb{R}^m$ . Therefore it has a basis!

In fact, we already know how to find the basis for  $\text{col}(A)$ ! It is the exact same problem as finding a basis for the span of a collection of vectors!

## Exercise

Find the column space of the following matrix. In particular, express it as the span of appropriate basis vectors.

$$\begin{bmatrix} 1 & -2 & 0 & 3 & -1 \\ 3 & 2 & 8 & 1 & 4 \\ 2 & 3 & 7 & 2 & 3 \\ -1 & 2 & 0 & 4 & -3 \end{bmatrix}$$

## Exercise

Find the column space of the following matrix. In particular, express it as the span of appropriate basis vectors.

$$\begin{bmatrix} 1 & -2 & 0 & 3 & -1 \\ 3 & 2 & 8 & 1 & 4 \\ 2 & 3 & 7 & 2 & 3 \\ -1 & 2 & 0 & 4 & -3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

## Exercise

Find the column space of the following matrix. In particular, express it as the span of appropriate basis vectors.

$$\begin{bmatrix} 1 & -2 & 0 & 3 & -1 \\ 3 & 2 & 8 & 1 & 4 \\ 2 & 3 & 7 & 2 & 3 \\ -1 & 2 & 0 & 4 & -3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Solution:

We can take as the basis the vectors corresponding to columns 1, 2, 4:

## Exercise

Find the column space of the following matrix. In particular, express it as the span of appropriate basis vectors.

$$\begin{bmatrix} 1 & -2 & 0 & 3 & -1 \\ 3 & 2 & 8 & 1 & 4 \\ 2 & 3 & 7 & 2 & 3 \\ -1 & 2 & 0 & 4 & -3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Solution:

We can take as the basis the vectors corresponding to columns 1, 2, 4:

$$\text{col}(A) = \text{span} \left\{ \begin{bmatrix} 1 \\ 3 \\ 2 \\ -1 \end{bmatrix}, \begin{bmatrix} -2 \\ 2 \\ 3 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ 2 \\ 4 \end{bmatrix} \right\}$$



## Exercise

Find the column space of the following matrix. In particular, express it as the span of appropriate basis vectors.

$$\begin{bmatrix} 1 & -2 & 0 & 3 & -1 \\ 3 & 2 & 8 & 1 & 4 \\ 2 & 3 & 7 & 2 & 3 \\ -1 & 2 & 0 & 4 & -3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Solution:

We can take as the basis the vectors corresponding to columns 1, 2, 4:

$$\text{col}(A) = \text{span} \left\{ \begin{bmatrix} 1 \\ 3 \\ 2 \\ -1 \end{bmatrix}, \begin{bmatrix} -2 \\ 2 \\ 3 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ 2 \\ 4 \end{bmatrix} \right\}$$

Note that the dimension of the column space is 3.

# Nullity and Rank

## Definition

The **nullity** of  $A$  is the dimension of the null space of  $A$ .

# Nullity and Rank

## Definition

The **nullity** of  $A$  is the dimension of the null space of  $A$ .

The **rank** of  $A$  is the dimension of the column space of  $A$ .

# Nullity and Rank

## Definition

The **nullity** of  $A$  is the dimension of the null space of  $A$ .

The **rank** of  $A$  is the dimension of the column space of  $A$ .

## Theorem (Dimension Theorem)

*Let  $A$  be an  $m \times n$  matrix. Then*

$$\text{rank } A + \text{nullity } A = n.$$

## Proof.

Reduce  $A$  to RREF.

# Nullity and Rank

## Definition

The **nullity** of  $A$  is the dimension of the null space of  $A$ .

The **rank** of  $A$  is the dimension of the column space of  $A$ .

## Theorem (Dimension Theorem)

*Let  $A$  be an  $m \times n$  matrix. Then*

$$\text{rank } A + \text{nullity } A = n.$$

## Proof.

Reduce  $A$  to RREF. Then the  $n$  columns are split into two groups: Those which describe the basis for the column space and those which give the null space. □

## Example

The  $5 \times 5$  matrix

$$\begin{bmatrix} 1 & 1 & 4 & 1 & 2 \\ 0 & 1 & 2 & 1 & 1 \\ 0 & 0 & 0 & 1 & 2 \\ 1 & -1 & 0 & 0 & 2 \\ 2 & 1 & 6 & 0 & 1 \end{bmatrix}$$

# Example

The  $5 \times 5$  matrix

$$\begin{bmatrix} 1 & 1 & 4 & 1 & 2 \\ 0 & 1 & 2 & 1 & 1 \\ 0 & 0 & 0 & 1 & 2 \\ 1 & -1 & 0 & 0 & 2 \\ 2 & 1 & 6 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 2 & 0 & -1 \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

## Example

The  $5 \times 5$  matrix

$$\begin{bmatrix} 1 & 1 & 4 & 1 & 2 \\ 0 & 1 & 2 & 1 & 1 \\ 0 & 0 & 0 & 1 & 2 \\ 1 & -1 & 0 & 0 & 2 \\ 2 & 1 & 6 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 2 & 0 & -1 \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

So  $\text{rank } A = 3$  and  $\text{nullity } A = 2$ .



## Example

The  $5 \times 5$  matrix

$$\begin{bmatrix} 1 & 1 & 4 & 1 & 2 \\ 0 & 1 & 2 & 1 & 1 \\ 0 & 0 & 0 & 1 & 2 \\ 1 & -1 & 0 & 0 & 2 \\ 2 & 1 & 6 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 2 & 0 & -1 \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

So  $\text{rank } A = 3$  and  $\text{nullity } A = 2$ . And of course,  $3 + 2 = 5$ .

# Summary

Both column space (rank) and null space (nullity) give us new ways to describe prior results:

# Summary

Both column space (rank) and null space (nullity) give us new ways to describe prior results:

## Theorem

*For a matrix  $A \in \mathbb{R}^{n \times n}$  the following statements are ALL EQUIVALENT:*

1. *A is nonsingular.*

# Summary

Both column space (rank) and null space (nullity) give us new ways to describe prior results:

## Theorem

*For a matrix  $A \in \mathbb{R}^{n \times n}$  the following statements are ALL EQUIVALENT:*

- 1.  $A$  is nonsingular.*
- 2.  $Ax = 0$  has only the trivial solution.*

# Summary

Both column space (rank) and null space (nullity) give us new ways to describe prior results:

## Theorem

*For a matrix  $A \in \mathbb{R}^{n \times n}$  the following statements are ALL EQUIVALENT:*

- 1.  $A$  is nonsingular.*
- 2.  $Ax = 0$  has only the trivial solution.*
- 3.  $A$  is equivalent to  $I$  (via elementary matrix operations).*

# Summary

Both column space (rank) and null space (nullity) give us new ways to describe prior results:

## Theorem

*For a matrix  $A \in \mathbb{R}^{n \times n}$  the following statements are ALL EQUIVALENT:*

- 1.  $A$  is nonsingular.*
- 2.  $Ax = 0$  has only the trivial solution.*
- 3.  $A$  is equivalent to  $I$  (via elementary matrix operations).*
- 4. For every  $b$  there is a unique solution to  $Ax = b$ .*

# Summary

Both column space (rank) and null space (nullity) give us new ways to describe prior results:

## Theorem

*For a matrix  $A \in \mathbb{R}^{n \times n}$  the following statements are ALL EQUIVALENT:*

- 1.  $A$  is nonsingular.*
- 2.  $Ax = 0$  has only the trivial solution.*
- 3.  $A$  is equivalent to  $I$  (via elementary matrix operations).*
- 4. For every  $b$  there is a unique solution to  $Ax = b$ .*
- 5.  $A$  can be written as a product of elementary matrices.*

# Summary

Both column space (rank) and null space (nullity) give us new ways to describe prior results:

## Theorem

*For a matrix  $A \in \mathbb{R}^{n \times n}$  the following statements are ALL EQUIVALENT:*

- 1.  $A$  is nonsingular.*
- 2.  $Ax = 0$  has only the trivial solution.*
- 3.  $A$  is equivalent to  $I$  (via elementary matrix operations).*
- 4. For every  $b$  there is a unique solution to  $Ax = b$ .*
- 5.  $A$  can be written as a product of elementary matrices.*
- 6.  $|A| \neq 0$ .*



# Summary

Both column space (rank) and null space (nullity) give us new ways to describe prior results:

## Theorem

*For a matrix  $A \in \mathbb{R}^{n \times n}$  the following statements are ALL EQUIVALENT:*

- 1.  $A$  is nonsingular.*
- 2.  $Ax = 0$  has only the trivial solution.*
- 3.  $A$  is equivalent to  $I$  (via elementary matrix operations).*
- 4. For every  $b$  there is a unique solution to  $Ax = b$ .*
- 5.  $A$  can be written as a product of elementary matrices.*
- 6.  $|A| \neq 0$ .*
- 7. The rank of  $A$  is  $n$ .*

# Summary

Both column space (rank) and null space (nullity) give us new ways to describe prior results:

## Theorem

*For a matrix  $A \in \mathbb{R}^{n \times n}$  the following statements are ALL EQUIVALENT:*

- 1.  $A$  is nonsingular.*
- 2.  $Ax = 0$  has only the trivial solution.*
- 3.  $A$  is equivalent to  $I$  (via elementary matrix operations).*
- 4. For every  $b$  there is a unique solution to  $Ax = b$ .*
- 5.  $A$  can be written as a product of elementary matrices.*
- 6.  $|A| \neq 0$ .*
- 7. The rank of  $A$  is  $n$ .*
- 8. The nullity of  $A$  is 0.*

# Summary

Both column space (rank) and null space (nullity) give us new ways to describe prior results:

## Theorem

*For a matrix  $A \in \mathbb{R}^{n \times n}$  the following statements are ALL EQUIVALENT:*

- 1.  $A$  is nonsingular.*
- 2.  $Ax = 0$  has only the trivial solution.*
- 3.  $A$  is equivalent to  $I$  (via elementary matrix operations).*
- 4. For every  $b$  there is a unique solution to  $Ax = b$ .*
- 5.  $A$  can be written as a product of elementary matrices.*
- 6.  $|A| \neq 0$ .*
- 7. The rank of  $A$  is  $n$ .*
- 8. The nullity of  $A$  is 0.*
- 9. The columns of  $A$  are linearly independent.*

# Final Comment

It is possible to also construct something called the *row space* of a matrix.

# Final Comment

It is possible to also construct something called the *row space* of a matrix. It turns out that the dimension of the row space is ALSO the rank of the matrix!

It is possible to also construct something called the *row space* of a matrix. It turns out that the dimension of the row space is ALSO the rank of the matrix! Therefore we could also add:

10. The rows of  $A$  are linearly independent.