Exercise

Find all solutions to Ax = 0 for

$$A = \begin{bmatrix} 1 & 1 & 4 & 1 & 2 \\ 0 & 1 & 2 & 1 & 1 \\ 0 & 0 & 0 & 1 & 2 \\ 1 & -1 & 0 & 0 & 2 \\ 2 & 1 & 6 & 0 & 1 \end{bmatrix}$$

Solution:

Use row reduction to show:

Solution:

Use row reduction to show:

Therefore

$$x_1 = -2x_3 - x_5$$

 $x_2 = -2x_3 + x_5$
 $x_3 = x_3$
 $x_4 = -2x_5$
 $x_5 = x_5$

Parametric Form of Solution

This can be written as

Parametric Form of Solution

This can be written as

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = s \begin{bmatrix} -2 \\ -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + t \begin{bmatrix} -1 \\ 1 \\ 0 \\ -2 \\ 1 \end{bmatrix}$$

Null Space

Definition

The **Null Space** of a matrix A is the set

$$N(A) := \{x \in \mathbb{R}^n : Ax = 0\}$$

Null Space

Definition

The **Null Space** of a matrix *A* is the set

$$N(A) := \{x \in \mathbb{R}^n : Ax = 0\}$$

That is, the null space is exactly the set of solutions to the homogeneous problem!

Null Space

Definition

The **Null Space** of a matrix A is the set

$$N(A) := \{x \in \mathbb{R}^n : Ax = 0\}$$

That is, the null space is exactly the set of solutions to the homogeneous problem!

Based on our exercise, we can write

$$N(A) = \operatorname{span} \left\{ \begin{bmatrix} -2 \\ -2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ 0 \\ -2 \\ 1 \end{bmatrix} \right\}.$$

It is not so hard to show that the null space of a matrix is a subspace

It is not so hard to show that the null space of a matrix is a subspace (in fact, it is a HW problem I assigned).

It is not so hard to show that the null space of a matrix is a subspace (in fact, it is a HW problem I assigned). Therefore it has a *basis* and *dimension*.

It is not so hard to show that the null space of a matrix is a subspace (in fact, it is a HW problem I assigned). Therefore it has a *basis* and *dimension*.

Definition

The dimension of the null space of a matrix A is called the **nullity** of A.

It is not so hard to show that the null space of a matrix is a subspace (in fact, it is a HW problem I assigned). Therefore it has a *basis* and *dimension*.

Definition

The dimension of the null space of a matrix A is called the **nullity** of A.

From our example, the basis of N(A) is

$$\left\{ \begin{bmatrix} -2\\ -2\\ 1\\ 0\\ 0 \end{bmatrix}, \begin{bmatrix} -1\\ 1\\ 0\\ -2\\ 1 \end{bmatrix} \right\}.$$

It is not so hard to show that the null space of a matrix is a subspace (in fact, it is a HW problem I assigned). Therefore it has a *basis* and *dimension*.

Definition

The dimension of the null space of a matrix A is called the **nullity** of A.

From our example, the basis of N(A) is

$$\left\{ \begin{bmatrix} -2\\ -2\\ 1\\ 0\\ 0 \end{bmatrix}, \begin{bmatrix} -1\\ 1\\ 0\\ -2\\ 1 \end{bmatrix} \right\}.$$

The nullity of A is 2.

Exercise

Exercise

Find a basis for the null space (and therefore also the nullity) for the following matrices:

$$A = \begin{bmatrix} 1 & 2 & 1 & 2 & 1 \\ 1 & 2 & 2 & 1 & 2 \\ 2 & 4 & 3 & 3 & 3 \\ 0 & 0 & 1 & -1 & 1 \end{bmatrix}$$

$$B = \begin{bmatrix} 1 & 0 & 2 \\ 2 & 1 & 3 \\ 3 & 1 & 2 \end{bmatrix}$$

For A:

$$\left\{ \begin{bmatrix} -2\\1\\0\\0\\0\end{bmatrix}, \begin{bmatrix} -3\\0\\1\\1\\0\end{bmatrix} \right\}$$

For A:

$$\left\{ \begin{bmatrix} -2\\1\\0\\0\\0 \end{bmatrix}, \begin{bmatrix} -3\\0\\1\\1\\0 \end{bmatrix} \right\}$$

And the nullity is 2.

For A:

$$\left\{ \begin{bmatrix} -2\\1\\0\\0\\0 \end{bmatrix}, \begin{bmatrix} -3\\0\\1\\1\\0 \end{bmatrix} \right\}$$

And the nullity is 2.

For B:

For A:

$$\left\{ \begin{bmatrix} -2\\1\\0\\0\\0 \end{bmatrix}, \begin{bmatrix} -3\\0\\1\\1\\0 \end{bmatrix} \right\}$$

And the nullity is 2.

For *B*: The null space is *trivial*!

For A:

$$\left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -3 \\ 0 \\ 1 \\ 1 \\ 0 \end{bmatrix} \right\}$$

And the nullity is 2.

For B: The null space is *trivial*! It has no basis and the nullity is 0.

Let A be a matrix and b a vector.

Let A be a matrix and b a vector. Let x be a solution to Ax = b.

Let A be a matrix and b a vector. Let x be a solution to Ax = b. Let $y \in N(A)$.

Let A be a matrix and b a vector. Let x be a solution to Ax = b. Let $y \in N(A)$. Then:

$$A(x+y)$$

Let A be a matrix and b a vector. Let x be a solution to Ax = b. Let $y \in N(A)$.

$$A(x+y)=Ax+Ay$$

Let A be a matrix and b a vector. Let x be a solution to Ax = b. Let $y \in N(A)$.

Then:

$$A(x+y) = Ax + Ay = b + 0$$

Let A be a matrix and b a vector.

Let x be a solution to Ax = b.

Let $y \in N(A)$.

Then:

$$A(x+y) = Ax + Ay = b + 0 = b.$$

Let A be a matrix and b a vector.

Let x be a solution to Ax = b.

Let $y \in N(A)$.

Then:

$$A(x+y) = Ax + Ay = b + 0 = b.$$

Let A be a matrix and b a vector.

Let x be a solution to Ax = b.

Let $y \in N(A)$.

Then:

$$A(x+y) = Ax + Ay = b + 0 = b.$$

$$A(x_1 - x_2)$$

Let A be a matrix and b a vector.

Let x be a solution to Ax = b.

Let $y \in N(A)$.

Then:

$$A(x+y) = Ax + Ay = b + 0 = b.$$

$$A(x_1-x_2)=Ax_1-Ax_2$$

Let A be a matrix and b a vector.

Let x be a solution to Ax = b.

Let $y \in N(A)$.

Then:

$$A(x+y) = Ax + Ay = b + 0 = b.$$

$$A(x_1-x_2) = Ax_1 - Ax_2 = b - b$$

Let A be a matrix and b a vector.

Let x be a solution to Ax = b.

Let $y \in N(A)$.

Then:

$$A(x+y) = Ax + Ay = b + 0 = b.$$

$$A(x_1-x_2)=Ax_1-Ax_2=b-b=0.$$

Let A be a matrix and b a vector.

Let x be a solution to Ax = b.

Let $y \in N(A)$.

Then:

$$A(x+y) = Ax + Ay = b + 0 = b.$$

Moreover: If x_1 and x_2 are both solutions to Ax = b then

$$A(x_1-x_2)=Ax_1-Ax_2=b-b=0.$$

So $x_1 - x_2$ is in the null space!

Let A be a matrix and b a vector.

Let x be a solution to Ax = b.

Let $y \in N(A)$.

Then:

$$A(x+y) = Ax + Ay = b + 0 = b.$$

Moreover: If x_1 and x_2 are both solutions to Ax = b then

$$A(x_1-x_2)=Ax_1-Ax_2=b-b=0.$$

So $x_1 - x_2$ is in the null space! Summary:

Let A be a matrix and b a vector.

Let x be a solution to Ax = b.

Let $y \in N(A)$.

Then:

$$A(x+y) = Ax + Ay = b + 0 = b.$$

Moreover: If x_1 and x_2 are both solutions to Ax = b then

$$A(x_1-x_2)=Ax_1-Ax_2=b-b=0.$$

So $x_1 - x_2$ is in the null space! Summary:

Solutions to a linear system differ by a vector in the null space.

Let A be a matrix and b a vector.

Let x be a solution to Ax = b.

Let $y \in N(A)$.

Then:

$$A(x+y) = Ax + Ay = b + 0 = b.$$

Moreover: If x_1 and x_2 are both solutions to Ax = b then

$$A(x_1-x_2)=Ax_1-Ax_2=b-b=0.$$

So $x_1 - x_2$ is in the null space! Summary:

- Solutions to a linear system differ by a vector in the null space.
- Solutions to linear systems are the null space shifted by a solution vector.

Consider the linear system

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 4 & -6 \\ 3 & 6 & -9 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}$$

Consider the linear system

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 4 & -6 \\ 3 & 6 & -9 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}$$

What is the null space of the system matrix?

Consider the linear system

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 4 & -6 \\ 3 & 6 & -9 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}$$

What is the null space of the system matrix?

$$N(A) = \operatorname{span} \left\{ \begin{bmatrix} -2\\1\\0 \end{bmatrix}, \begin{bmatrix} 3\\0\\1 \end{bmatrix} \right\}.$$

Consider the linear system

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 4 & -6 \\ 3 & 6 & -9 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}$$

What is the null space of the system matrix?

$$N(A) = \operatorname{span} \left\{ \begin{bmatrix} -2\\1\\0 \end{bmatrix}, \begin{bmatrix} 3\\0\\1 \end{bmatrix} \right\}.$$

Note also that $\begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}$ is a solution.

Example Continued

Therefore the set of all solutions to the problem is

$$\left\{ \begin{bmatrix} 2\\0\\0 \end{bmatrix} + s \begin{bmatrix} -2\\1\\0 \end{bmatrix} + t \begin{bmatrix} 3\\0\\1 \end{bmatrix} : s,t \in \mathbb{R} \right\}$$

Example Continued

Therefore the set of all solutions to the problem is

$$\left\{ \begin{bmatrix} 2\\0\\0 \end{bmatrix} + s \begin{bmatrix} -2\\1\\0 \end{bmatrix} + t \begin{bmatrix} 3\\0\\1 \end{bmatrix} : s,t \in \mathbb{R} \right\}$$

This is exactly the parametric form for a plane through the point (2,0,0)! That is, it is the plane "shifted" to (2,0,0).