

Exercises

Let $A, B \in \mathbb{R}^{n \times n}$ and $x, y \in \mathbb{R}^n$. For each of the following determine if the statement is true or false (and try to give a reason why):

1. A linear system of three equations can have exactly three different solutions.
2. Suppose none of the entries of A or B are 0. Then $AB \neq 0$.
3. $Ax \in \mathbb{R}^n$.
4. $A + A^T$ is symmetric.
5. Ax is a linear combination of the columns of A .
6. If $Ax = 0$ then $x = 0$.
7. Homogenous linear systems are consistent.
8. If $Ax = Ay$ then $x = y$.

Exercises

Let $A, B \in \mathbb{R}^{n \times n}$ and $x, y \in \mathbb{R}^n$. For each of the following determine if the statement is true or false (and try to give a reason why):

1. A linear system of three equations can have exactly three different solutions.

Exercises

Let $A, B \in \mathbb{R}^{n \times n}$ and $x, y \in \mathbb{R}^n$. For each of the following determine if the statement is true or false (and try to give a reason why):

1. A linear system of three equations can have exactly three different solutions.

FALSE

Exercises

Let $A, B \in \mathbb{R}^{n \times n}$ and $x, y \in \mathbb{R}^n$. For each of the following determine if the statement is true or false (and try to give a reason why):

1. A linear system of three equations can have exactly three different solutions.

FALSE: If a linear system has more than one solution, then it must have an infinite number of solutions.

Exercises

Let $A, B \in \mathbb{R}^{n \times n}$ and $x, y \in \mathbb{R}^n$. For each of the following determine if the statement is true or false (and try to give a reason why):

1. A linear system of three equations can have exactly three different solutions.

FALSE: If a linear system has more than one solution, then it must have an infinite number of solutions.

2. Suppose none of the entries of A or B are 0. Then $AB \neq 0$.

Exercises

Let $A, B \in \mathbb{R}^{n \times n}$ and $x, y \in \mathbb{R}^n$. For each of the following determine if the statement is true or false (and try to give a reason why):

1. A linear system of three equations can have exactly three different solutions.

FALSE: If a linear system has more than one solution, then it must have an infinite number of solutions.

2. Suppose none of the entries of A or B are 0. Then $AB \neq 0$.

FALSE

Exercises

Let $A, B \in \mathbb{R}^{n \times n}$ and $x, y \in \mathbb{R}^n$. For each of the following determine if the statement is true or false (and try to give a reason why):

1. A linear system of three equations can have exactly three different solutions.

FALSE: If a linear system has more than one solution, then it must have an infinite number of solutions.

2. Suppose none of the entries of A or B are 0. Then $AB \neq 0$.

FALSE: Let $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$.

Exercises

Let $A, B \in \mathbb{R}^{n \times n}$ and $x, y \in \mathbb{R}^n$. For each of the following determine if the statement is true or false (and try to give a reason why):

1. A linear system of three equations can have exactly three different solutions.

FALSE: If a linear system has more than one solution, then it must have an infinite number of solutions.

2. Suppose none of the entries of A or B are 0. Then $AB \neq 0$.

FALSE: Let $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$.

3. $Ax \in \mathbb{R}^n$.

Exercises

Let $A, B \in \mathbb{R}^{n \times n}$ and $x, y \in \mathbb{R}^n$. For each of the following determine if the statement is true or false (and try to give a reason why):

1. A linear system of three equations can have exactly three different solutions.

FALSE: If a linear system has more than one solution, then it must have an infinite number of solutions.

2. Suppose none of the entries of A or B are 0. Then $AB \neq 0$.

FALSE: Let $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$.

3. $Ax \in \mathbb{R}^n$.

TRUE

Exercises

Let $A, B \in \mathbb{R}^{n \times n}$ and $x, y \in \mathbb{R}^n$. For each of the following determine if the statement is true or false (and try to give a reason why):

1. A linear system of three equations can have exactly three different solutions.

FALSE: If a linear system has more than one solution, then it must have an infinite number of solutions.

2. Suppose none of the entries of A or B are 0. Then $AB \neq 0$.

FALSE: Let $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$.

3. $Ax \in \mathbb{R}^n$.

TRUE: The number of elements in Ax is the number of rows of $A \in \mathbb{R}^{n \times n}$.

Exercises

4 $A + A^T$ is symmetric.

Exercises

- 4 $A + A^T$ is symmetric.
TRUE

Exercises

4 $A + A^T$ is symmetric.

TRUE:

$$(A + A^T)^T = A^T + (A^T)^T = A^T + A = A + A^T.$$

Therefore $A + A^T$ is equal to its own transpose and must be symmetric.

Exercises

4 $A + A^T$ is symmetric.

TRUE:

$$(A + A^T)^T = A^T + (A^T)^T = A^T + A = A + A^T.$$

Therefore $A + A^T$ is equal to its own transpose and must be symmetric.

5 Ax is a linear combination of the columns of A .

Exercises

- 4 $A + A^T$ is symmetric.

TRUE:

$$(A + A^T)^T = A^T + (A^T)^T = A^T + A = A + A^T.$$

Therefore $A + A^T$ is equal to its own transpose and must be symmetric.

- 5 Ax is a linear combination of the columns of A .

TRUE

Exercises

- 4 $A + A^T$ is symmetric.

TRUE:

$$(A + A^T)^T = A^T + (A^T)^T = A^T + A = A + A^T.$$

Therefore $A + A^T$ is equal to its own transpose and must be symmetric.

- 5 Ax is a linear combination of the columns of A .

TRUE: Write $A = [a_1 \cdots a_n]$ where a_i is the i -th column of A and $x = [x_1 \cdots x_n]^T$. Then

$$Ax = \begin{bmatrix} a_1 & \cdots & a_n \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = \sum_{i=1}^n x_i a_i.$$

6 If $Ax = 0$ then $x = 0$.

6 If $Ax = 0$ then $x = 0$.
FALSE

Exercises

6 If $Ax = 0$ then $x = 0$.

FALSE:

$$\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 0.$$

Exercises

6 If $Ax = 0$ then $x = 0$.

FALSE:

$$\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 0.$$

7 Homogenous linear systems are consistent.

Exercises

6 If $Ax = 0$ then $x = 0$.

FALSE:

$$\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 0.$$

7 Homogenous linear systems are consistent. TRUE

Exercises

6 If $Ax = 0$ then $x = 0$.

FALSE:

$$\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 0.$$

7 Homogenous linear systems are consistent. TRUE: The zero vector is always a solution to $Ax = 0$.

Exercises

6 If $Ax = 0$ then $x = 0$.

FALSE:

$$\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 0.$$

7 Homogenous linear systems are consistent. TRUE: The zero vector is always a solution to $Ax = 0$.

8 If $Ax = Ay$ then $x = y$.

Exercises

6 If $Ax = 0$ then $x = 0$.

FALSE:

$$\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 0.$$

7 Homogenous linear systems are consistent. TRUE: The zero vector is always a solution to $Ax = 0$.

8 If $Ax = Ay$ then $x = y$. FALSE.

More Comments about Linear Systems and Matrices

From last time (and from the warm up exercises 6-8) matrix multiplication does not have a “cancellation law.”

More Comments about Linear Systems and Matrices

From last time (and from the warm up exercises 6-8) matrix multiplication does not have a “cancellation law.” That is:

$$Ax = Ay \not\Rightarrow x = y.$$

More Comments about Linear Systems and Matrices

From last time (and from the warm up exercises 6-8) matrix multiplication does not have a “cancellation law.” That is:

$$Ax = Ay \not\Rightarrow x = y.$$

This manifests itself in the context of solving systems of equations: Some systems do not have solutions/have an infinite number of solutions.

More Comments about Linear Systems and Matrices

From last time (and from the warm up exercises 6-8) matrix multiplication does not have a “cancellation law.” That is:

$$Ax = Ay \not\Rightarrow x = y.$$

This manifests itself in the context of solving systems of equations: Some systems do not have solutions/have an infinite number of solutions. Both are awkward.

More Comments about Linear Systems and Matrices

From last time (and from the warm up exercises 6-8) matrix multiplication does not have a “cancellation law.” That is:

$$Ax = Ay \not\Rightarrow x = y.$$

This manifests itself in the context of solving systems of equations: Some systems do not have solutions/have an infinite number of solutions. Both are awkward.

OTOH, we know that some systems DO have unique solutions. What types of matrices are they associated with?

Nonsingular and Inverse Matrices

Definition

$A \in \mathbb{R}^{n \times n}$ is called **nonsingular** or **invertible** if there exists another matrix $B \in \mathbb{R}^{n \times n}$ so that $AB = I$.

Nonsingular and Inverse Matrices

Definition

$A \in \mathbb{R}^{n \times n}$ is called **nonsingular** or **invertible** if there exists another matrix $B \in \mathbb{R}^{n \times n}$ so that $AB = I$.

1. Recall that I is the identity matrix in $\mathbb{R}^{n \times n}$.

Nonsingular and Inverse Matrices

Definition

$A \in \mathbb{R}^{n \times n}$ is called **nonsingular** or **invertible** if there exists another matrix $B \in \mathbb{R}^{n \times n}$ so that $AB = I$.

1. Recall that I is the identity matrix in $\mathbb{R}^{n \times n}$.
2. We can prove (later) that matrices commute with their inverses: $AB = BA = I$.

Nonsingular and Inverse Matrices

Definition

$A \in \mathbb{R}^{n \times n}$ is called **nonsingular** or **invertible** if there exists another matrix $B \in \mathbb{R}^{n \times n}$ so that $AB = I$.

1. Recall that I is the identity matrix in $\mathbb{R}^{n \times n}$.
2. We can prove (later) that matrices commute with their inverses: $AB = BA = I$.
3. In the above, B is called an **inverse** of A .

Nonsingular and Inverse Matrices

Definition

$A \in \mathbb{R}^{n \times n}$ is called **nonsingular** or **invertible** if there exists another matrix $B \in \mathbb{R}^{n \times n}$ so that $AB = I$.

1. Recall that I is the identity matrix in $\mathbb{R}^{n \times n}$.
2. We can prove (later) that matrices commute with their inverses: $AB = BA = I$.
3. In the above, B is called an **inverse** of A .

Exercise

Show that $\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$ is nonsingular by finding an explicit inverse matrix.

Nonsingular and Inverse Matrices

Definition

$A \in \mathbb{R}^{n \times n}$ is called **nonsingular** or **invertible** if there exists another matrix $B \in \mathbb{R}^{n \times n}$ so that $AB = I$.

1. Recall that I is the identity matrix in $\mathbb{R}^{n \times n}$.
2. We can prove (later) that matrices commute with their inverses: $AB = BA = I$.
3. In the above, B is called an **inverse** of A .

Exercise

Show that $\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$ is nonsingular by finding an explicit inverse matrix.

It is: $\begin{bmatrix} 1/2 & 0 \\ 0 & 1/3 \end{bmatrix}$

A Good Example

Let a, b, c, d be real numbers and $ad - bc \neq 0$. Verify that

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}.$$

is invertible with inverse

$$\frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Inverse Matrices are Unique

Suppose a non-singular matrix A has two inverses called B and C .

Inverse Matrices are Unique

Suppose a non-singular matrix A has two inverses called B and C .
Then

$$B = BI_n$$

Inverse Matrices are Unique

Suppose a non-singular matrix A has two inverses called B and C .
Then

$$\begin{aligned} B &= BI_n \\ &= B(AC) \end{aligned}$$

Inverse Matrices are Unique

Suppose a non-singular matrix A has two inverses called B and C .
Then

$$\begin{aligned} B &= BI_n \\ &= B(AC) \\ &= (BA)C \end{aligned}$$

Inverse Matrices are Unique

Suppose a non-singular matrix A has two inverses called B and C .
Then

$$\begin{aligned} B &= BI_n \\ &= B(AC) \\ &= (BA)C \\ &= I_n C \end{aligned}$$

Inverse Matrices are Unique

Suppose a non-singular matrix A has two inverses called B and C .
Then

$$\begin{aligned} B &= BI_n \\ &= B(AC) \\ &= (BA)C \\ &= I_n C \\ &= C. \end{aligned}$$

Inverse Matrices are Unique

Suppose a non-singular matrix A has two inverses called B and C .
Then

$$\begin{aligned} B &= BI_n \\ &= B(AC) \\ &= (BA)C \\ &= I_n C \\ &= C. \end{aligned}$$

So $B = C$. Therefore it is ok to say B is THE inverse of A .

Inverse Matrices are Unique

Suppose a non-singular matrix A has two inverses called B and C .
Then

$$\begin{aligned} B &= BI_n \\ &= B(AC) \\ &= (BA)C \\ &= I_n C \\ &= C. \end{aligned}$$

So $B = C$. Therefore it is ok to say B is THE inverse of A .

Notation: If A is nonsingular then we write A^{-1} for its unique inverse.

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$.

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1})$$

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1}$$

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1}$$

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1}$$

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I.$$

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I.$$

Since $(AB)(B^{-1}A^{-1})$ we know $B^{-1}A^{-1}$ is the inverse of AB .

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I.$$

Since $(AB)(B^{-1}A^{-1})$ we know $B^{-1}A^{-1}$ is the inverse of AB .
So $B^{-1}A^{-1} = (AB)^{-1}$.

2. $(A^{-1})^{-1} = A$.

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I.$$

Since $(AB)(B^{-1}A^{-1})$ we know $B^{-1}A^{-1}$ is the inverse of AB .
So $B^{-1}A^{-1} = (AB)^{-1}$.

2. $(A^{-1})^{-1} = A$. Since

$$AA^{-1} = I$$

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I.$$

Since $(AB)(B^{-1}A^{-1})$ we know $B^{-1}A^{-1}$ is the inverse of AB .
So $B^{-1}A^{-1} = (AB)^{-1}$.

2. $(A^{-1})^{-1} = A$. Since

$$AA^{-1} = I \Leftrightarrow A \text{ is the inverse of } A^{-1}$$

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I.$$

Since $(AB)(B^{-1}A^{-1})$ we know $B^{-1}A^{-1}$ is the inverse of AB .
So $B^{-1}A^{-1} = (AB)^{-1}$.

2. $(A^{-1})^{-1} = A$. Since

$$AA^{-1} = I \Leftrightarrow A \text{ is the inverse of } A^{-1} \Leftrightarrow A = (A^{-1})^{-1}.$$

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I.$$

Since $(AB)(B^{-1}A^{-1})$ we know $B^{-1}A^{-1}$ is the inverse of AB .
So $B^{-1}A^{-1} = (AB)^{-1}$.

2. $(A^{-1})^{-1} = A$. Since

$$AA^{-1} = I \Leftrightarrow A \text{ is the inverse of } A^{-1} \Leftrightarrow A = (A^{-1})^{-1}.$$

3. $(A^{-1})^T = (A^T)^{-1}$.

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I.$$

Since $(AB)(B^{-1}A^{-1})$ we know $B^{-1}A^{-1}$ is the inverse of AB .
So $B^{-1}A^{-1} = (AB)^{-1}$.

2. $(A^{-1})^{-1} = A$. Since

$$AA^{-1} = I \Leftrightarrow A \text{ is the inverse of } A^{-1} \Leftrightarrow A = (A^{-1})^{-1}.$$

3. $(A^{-1})^T = (A^T)^{-1}$. Since by transposing $I = AA^{-1}$:

$$I^T = (AA^{-1})^T$$

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I.$$

Since $(AB)(B^{-1}A^{-1})$ we know $B^{-1}A^{-1}$ is the inverse of AB .
So $B^{-1}A^{-1} = (AB)^{-1}$.

2. $(A^{-1})^{-1} = A$. Since

$$AA^{-1} = I \Leftrightarrow A \text{ is the inverse of } A^{-1} \Leftrightarrow A = (A^{-1})^{-1}.$$

3. $(A^{-1})^T = (A^T)^{-1}$. Since by transposing $I = AA^{-1}$:

$$I^T = (AA^{-1})^T \Leftrightarrow I = (A^{-1})^T A^T$$

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I.$$

Since $(AB)(B^{-1}A^{-1})$ we know $B^{-1}A^{-1}$ is the inverse of AB .
So $B^{-1}A^{-1} = (AB)^{-1}$.

2. $(A^{-1})^{-1} = A$. Since

$$AA^{-1} = I \Leftrightarrow A \text{ is the inverse of } A^{-1} \Leftrightarrow A = (A^{-1})^{-1}.$$

3. $(A^{-1})^T = (A^T)^{-1}$. Since by transposing $I = AA^{-1}$:

$$I^T = (AA^{-1})^T \Leftrightarrow I = (A^{-1})^T A^T \Leftrightarrow (A^{-1})^T \text{ is the inverse of } A^T$$

Properties of A^{-1}

Let A and B be nonsingular $n \times n$ matrices.

1. $(AB)^{-1} = B^{-1}A^{-1}$. Since

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I.$$

Since $(AB)(B^{-1}A^{-1}) = I$ we know $B^{-1}A^{-1}$ is the inverse of AB .
So $B^{-1}A^{-1} = (AB)^{-1}$.

2. $(A^{-1})^{-1} = A$. Since

$$AA^{-1} = I \Leftrightarrow A \text{ is the inverse of } A^{-1} \Leftrightarrow A = (A^{-1})^{-1}.$$

3. $(A^{-1})^T = (A^T)^{-1}$. Since by transposing $I = AA^{-1}$:

$$I^T = (AA^{-1})^T \Leftrightarrow I = (A^{-1})^T A^T \Leftrightarrow (A^{-1})^T \text{ is the inverse of } A^T \Leftrightarrow (A^{-1})^T = (A^T)^{-1}.$$

What is Special About Non-Singular Matrices

Let Suppose $AB = AC$ for matrices A , B , and C . Moreover, assume that A is non-singular with inverse A^{-1} . Then:

$$AB = AC$$

What is Special About Non-Singular Matrices

Let Suppose $AB = AC$ for matrices A , B , and C . Moreover, assume that A is non-singular with inverse A^{-1} . Then:

$$\begin{aligned}AB &= AC \\ A^{-1}AB &= A^{-1}AC\end{aligned}$$

What is Special About Non-Singular Matrices

Let Suppose $AB = AC$ for matrices A , B , and C . Moreover, assume that A is non-singular with inverse A^{-1} . Then:

$$\begin{aligned}AB &= AC \\A^{-1}AB &= A^{-1}AC \\B &= C!\end{aligned}$$

What is Special About Non-Singular Matrices

Let Suppose $AB = AC$ for matrices A , B , and C . Moreover, assume that A is non-singular with inverse A^{-1} . Then:

$$\begin{aligned}AB &= AC \\ A^{-1}AB &= A^{-1}AC \\ B &= C!\end{aligned}$$

Therefore: If matrices are non-singular, then they can be “canceled” in matrix multiplication!

What is Special About Non-Singular Matrices

Let Suppose $AB = AC$ for matrices A , B , and C . Moreover, assume that A is non-singular with inverse A^{-1} . Then:

$$\begin{aligned}AB &= AC \\A^{-1}AB &= A^{-1}AC \\B &= C!\end{aligned}$$

Therefore: If matrices are non-singular, then they can be “canceled” in matrix multiplication!

In particular: If A is an invertible matrix then

$$Ax = b \Leftrightarrow x = A^{-1}b$$

Summary

If the system matrix A for a linear system $Ax = b$ is non-singular, then the linear system has the unique solution $x = A^{-1}b$!

Summary

If the system matrix A for a linear system $Ax = b$ is non-singular, then the linear system has the unique solution $x = A^{-1}b$!

This gives us a naive algorithm for solving some linear systems:

Summary

If the system matrix A for a linear system $Ax = b$ is non-singular, then the linear system has the unique solution $x = A^{-1}b$!

This gives us a naive algorithm for solving some linear systems:

1. Express the system in the form $Ax = b$.

Summary

If the system matrix A for a linear system $Ax = b$ is non-singular, then the linear system has the unique solution $x = A^{-1}b$!

This gives us a naive algorithm for solving some linear systems:

1. Express the system in the form $Ax = b$.
2. Show that the system matrix is non-singular by finding A^{-1} .

Summary

If the system matrix A for a linear system $Ax = b$ is non-singular, then the linear system has the unique solution $x = A^{-1}b$!

This gives us a naive algorithm for solving some linear systems:

1. Express the system in the form $Ax = b$.
2. Show that the system matrix is non-singular by finding A^{-1} .
3. Compute $x = A^{-1}b$.

Summary

If the system matrix A for a linear system $Ax = b$ is non-singular, then the linear system has the unique solution $x = A^{-1}b$!

This gives us a naive algorithm for solving some linear systems:

1. Express the system in the form $Ax = b$.
2. Show that the system matrix is non-singular by finding A^{-1} .
3. Compute $x = A^{-1}b$.

Exercise

Solve the linear system

$$\begin{aligned} 2x + y &= 4 \\ -2x + y &= -4 \end{aligned}$$

Result

Result:

$$\left\{ \begin{array}{l} 2x + y = 4 \\ -2x + y = -4 \end{array} \right\} \Leftrightarrow \begin{bmatrix} 2 & 1 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ -4 \end{bmatrix} \Leftrightarrow Ax = b$$

Result

Result:

$$\left\{ \begin{array}{l} 2x + y = 4 \\ -2x + y = -4 \end{array} \right\} \Leftrightarrow \begin{bmatrix} 2 & 1 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ -4 \end{bmatrix} \Leftrightarrow Ax = b$$

So

$$x = A^{-1}b = \begin{bmatrix} 1/4 & -1/4 \\ 1/2 & 1/2 \end{bmatrix} \begin{bmatrix} 4 \\ -4 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$$

Result

Result:

$$\left\{ \begin{array}{l} 2x + y = 4 \\ -2x + y = -4 \end{array} \right\} \Leftrightarrow \begin{bmatrix} 2 & 1 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ -4 \end{bmatrix} \Leftrightarrow Ax = b$$

So

$$x = A^{-1}b = \begin{bmatrix} 1/4 & -1/4 \\ 1/2 & 1/2 \end{bmatrix} \begin{bmatrix} 4 \\ -4 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$$

So the solution set is $\{(x, y) = (2, 0)\}$.

If there is time

A very natural thing to do is *partition* matrices into *submatrices*.

If there is time

A very natural thing to do is *partition* matrices into *submatrices*.
For example, consider the following matrix:

$$A = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}$$

What is interesting about it?

If there is time

A very natural thing to do is *partition* matrices into *submatrices*.
For example, consider the following matrix:

$$A = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}$$

What is interesting about it? It looks like four little identity matrices put together:

$$A = \left[\begin{array}{cc|cc} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ \hline 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{array} \right]$$

If there is time

A very natural thing to do is *partition* matrices into *submatrices*. For example, consider the following matrix:

$$A = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}$$

What is interesting about it? It looks like four little identity matrices put together:

$$A = \left[\begin{array}{cc|cc} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ \hline 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{array} \right]$$

So we might write:

$$A = \begin{bmatrix} I_2 & I_2 \\ I_2 & I_2 \end{bmatrix}$$

Partitioned Matrices Results

In short: Operations on partition matrices work the same as on regular matrices provided the sizes of the partitions make sense. There are some exercises in the HW which explore this.