

Warm Up Exercises

Let

$$A = \begin{bmatrix} -1 & 2 & -5 \\ 2 & -1 & 6 \\ 2 & -2 & 7 \end{bmatrix}, B = \begin{bmatrix} 1 & -2 & 5 \\ 0 & 3 & -4 \\ 0 & 0 & -1 \end{bmatrix}$$

1. Show that A and B are row equivalent. What sequence of row operations did you use?
2. What is the reduced row echelon form of A (and B)?
3. Explain why A (and B) must be invertible.

Answers

1. Do it on the board.

Answers

1. Do it on the board.
2. The identity matrix.

Answers

1. Do it on the board.
2. The identity matrix.
3. A can be written as the product of invertible matrices:

$$A = \prod_{i=1}^n E_i I_3.$$

Answers

1. Do it on the board.
2. The identity matrix.
3. A can be written as the product of invertible matrices:

$$A = \prod_{i=1}^n E_i I_3.$$

Here the E_i are all elementary row operation matrices (which are invertible).

More Exercises!

Now consider the matrix:

$$A = \begin{bmatrix} 2 & 5 & 1 \\ -1 & -1 & 0 \end{bmatrix}$$

1. Let B be the RREF of A . Find B .
2. Interpret both A and B as an augmented matrix and write down the corresponding system of equations.
3. Show that the solution to the system of equations corresponding to B is also the solution to the system of equations corresponding to A .

Results

We have:

$$\begin{bmatrix} 2 & 5 & 1 \\ -1 & -1 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & -1/3 \\ 0 & 1 & 1/3 \end{bmatrix}.$$

Results

We have:

$$\begin{bmatrix} 2 & 5 & 1 \\ -1 & -1 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & -1/3 \\ 0 & 1 & 1/3 \end{bmatrix}.$$

This gives the two systems:

$$\begin{array}{rcl} 2x + 5y & = & 1 \\ -x - y & = & 0 \end{array} \quad \text{and} \quad \begin{array}{rcl} x & = & -1/3 \\ y & = & 1/3 \end{array}$$

Theorem

Theorem

Let A and B be row equivalent matrices. Then the corresponding system of equations have the same solution set.

Theorem

Theorem

Let A and B be row equivalent matrices. Then the corresponding system of equations have the same solution set.

Proof.

Matrices which are row equivalent differ by a finite sequence of row operations.

Theorem

Theorem

Let A and B be row equivalent matrices. Then the corresponding system of equations have the same solution set.

Proof.

Matrices which are row equivalent differ by a finite sequence of row operations. When interpreted as systems of equations each row operation corresponds to an elimination operation.

Theorem

Theorem

Let A and B be row equivalent matrices. Then the corresponding system of equations have the same solution set.

Proof.

Matrices which are row equivalent differ by a finite sequence of row operations. When interpreted as systems of equations each row operation corresponds to an elimination operation. Systems of equations which differ by a finite collection of elimination operations have the same solution set. □

Gaussian Elimination (aka the Gauss-Jordan Procedure)

We can discuss the process of finding RREF for a matrix as an algorithm.

Gaussian Elimination (aka the Gauss-Jordan Procedure)

We can discuss the process of finding RREF for a matrix as an algorithm. This requires a little definition:

Definition

A **pivot** is a non-zero matrix entry with only zero entries to the left of it.

Algorithm Phase 1 (Forward elimination)

Let $k, \ell = 1$.

Algorithm Phase 1 (Forward elimination)

Let $k, \ell = 1$.

1. While $x_{ij} = 0$ for $i \geq k$ and $j \geq \ell$ set $\ell = \ell + 1$.
2. Working from row k to the bottom find the first pivot in the ℓ -th column.
3. Swap rows to place the row with the first pivot into the k -th row.
4. Make the pivot value a one.
5. Make all values below the pivot a 0.
6. Set $k = k + 1$.
7. If there are more columns go back to (1). If not stop.

Algorithm Phase 1 (Forward elimination)

Let $k, \ell = 1$.

1. While $x_{ij} = 0$ for $i \geq k$ and $j \geq \ell$ set $\ell = \ell + 1$.
2. Working from row k to the bottom find the first pivot in the ℓ -th column.
3. Swap rows to place the row with the first pivot into the k -th row.
4. Make the pivot value a one.
5. Make all values below the pivot a 0.
6. Set $k = k + 1$.
7. If there are more columns go back to (1). If not stop.

At this point the matrix will be in REF.

Algorithm phase 2 (Back Substitution)

1. Moving from right to left make all entries ABOVE the pivots zero.

Algorithm phase 2 (Back Substitution)

1. Moving from right to left make all entries ABOVE the pivots zero.

Now the matrix will be in RREF.

Exercises

Solve the following systems of equations:

$$x + 2y + 3z = 9$$

$$2x - y + z = 8$$

$$3x - z = 3$$

$$x + y + z + w = 0$$

$$x + w = 0$$

$$x + 2y + z = 0$$

$$x + 2y + 3z + 4w = 5$$

$$y + 2z + 3w = 6$$

$$x + 3y + 5z + 7w = 11$$