

From Example to Abstraction

Our fundamental example of a vector space is \mathbb{R}^n with standard addition and scalar multiplication.

From Example to Abstraction

Our fundamental example of a vector space is \mathbb{R}^n with standard addition and scalar multiplication.

Properties:

1. $u + v = v + u$
2. $u + (v + w) = (u + v) + w$
3. $u + 0 = 0 + u = u$
4. $u + (-u) = 0$
5. $r(u + v) = ru + rv$
6. $(r + s)u = ru + su$
7. $r(su) = (rs)u$
8. $1u = u$

From Example to Abstraction

Our fundamental example of a vector space is \mathbb{R}^n with standard addition and scalar multiplication.

Properties:

1. $u + v = v + u$
2. $u + (v + w) = (u + v) + w$
3. $u + 0 = 0 + u = u$
4. $u + (-u) = 0$
5. $r(u + v) = ru + rv$
6. $(r + s)u = ru + su$
7. $r(su) = (rs)u$
8. $1u = u$

From this example, we move to abstract vector spaces by *using the properties as the definition!*

Definition of a Vector Space

Definition

Let F be a field (i.e., the real numbers). A **vector space** over F is a set V (the vectors) and operations

$\oplus : V \times V \rightarrow V$ (vector addition) $\odot : F \times V \rightarrow V$ scalar multiplication

with the following properties:

For all vectors u , v , and w in V and scalar values r and s in F we have

1. $u \oplus v = v \oplus u$
2. $u \oplus (v \oplus w) = (u \oplus v) \oplus w$
3. There is a vector $e \in V$ so that $u \oplus e = u$
4. There is a vector u^{-1} so that $u \oplus u^{-1} = e$.
5. $r \odot (u \oplus v) = (r \odot u) \oplus (r \odot v)$
6. $(r + s) \odot u = (r \odot u) \oplus (s \odot u)$
7. $r \odot (s \odot u) = (rs) \odot u$
8. $1 \odot u = u$

Today:

We will do the same thing with the dot product.

Properties of the Dot Product

Let u , v , and w be vectors in \mathbb{R}^n and c a scalar

1. $u \cdot u \geq 0$ and $u \cdot u = 0 \Leftrightarrow u = 0$.
2. Commutativity: $u \cdot v = v \cdot u$.
3. Distributivity: $u \cdot (v + w) = u \cdot v + u \cdot w$.
4. Scalar associativity: $c(u \cdot v) = (cu) \cdot v$.

Properties of the Dot Product

Let u , v , and w be vectors in \mathbb{R}^n and c a scalar

1. $u \cdot u \geq 0$ and $u \cdot u = 0 \Leftrightarrow u = 0$.
2. Commutativity: $u \cdot v = v \cdot u$.
3. Distributivity: $u \cdot (v + w) = u \cdot v + u \cdot w$.
4. Scalar associativity: $c(u \cdot v) = (cu) \cdot v$.

Notes:

Properties of the Dot Product

Let u , v , and w be vectors in \mathbb{R}^n and c a scalar

1. $u \cdot u \geq 0$ and $u \cdot u = 0 \Leftrightarrow u = 0$.
2. Commutativity: $u \cdot v = v \cdot u$.
3. Distributivity: $u \cdot (v + w) = u \cdot v + u \cdot w$.
4. Scalar associativity: $c(u \cdot v) = (cu) \cdot v$.

Notes:

1. We PROVE these properties FROM the computational definition of the dot product.

Properties of the Dot Product

Let u , v , and w be vectors in \mathbb{R}^n and c a scalar

1. $u \cdot u \geq 0$ and $u \cdot u = 0 \Leftrightarrow u = 0$.
2. Commutativity: $u \cdot v = v \cdot u$.
3. Distributivity: $u \cdot (v + w) = u \cdot v + u \cdot w$.
4. Scalar associativity: $c(u \cdot v) = (cu) \cdot v$.

Notes:

1. We PROVE these properties FROM the computational definition of the dot product.
2. These properties are important because (amongst other reasons) they inform us of (and align with) the geometry of \mathbb{R}^n .

Properties of the Dot Product

Let u , v , and w be vectors in \mathbb{R}^n and c a scalar

1. $u \cdot u \geq 0$ and $u \cdot u = 0 \Leftrightarrow u = 0$.
2. Commutativity: $u \cdot v = v \cdot u$.
3. Distributivity: $u \cdot (v + w) = u \cdot v + u \cdot w$.
4. Scalar associativity: $c(u \cdot v) = (cu) \cdot v$.

Notes:

1. We PROVE these properties FROM the computational definition of the dot product.
2. These properties are important because (amongst other reasons) they inform us of (and align with) the geometry of \mathbb{R}^n .

So let's use these properties to DEFINE geometry in general!

The Inner Product

The Inner Product

Definition

Let V be a vector space over the scalar field F . An **inner product** on V is a function:

$$(\cdot, \cdot) : V \times V \rightarrow F.$$

Written as (v, w) .

The Inner Product

Definition

Let V be a vector space over the scalar field F . An **inner product** on V is a function:

$$(\cdot, \cdot) : V \times V \rightarrow F.$$

Written as (v, w) . It has the following properties:

1. $(u, u) \geq 0$ and $(u, u) = 0 \Leftrightarrow u = 0$.
2. $(u, v) = (v, u)$.
3. $(u, v + w) = (u, v) + (u, w)$.
4. $c(u, v) = (cu, v) = (cv, u)$.

Example (1) of Inner Products

In general there are lots of inner products on vector spaces (more on this later).

Example (1) of Inner Products

In general there are lots of inner products on vector spaces (more on this later). But here are two examples that you should be familiar with:

Example (1) of Inner Products

In general there are lots of inner products on vector spaces (more on this later). But here are two examples that you should be familiar with:

The *dot product* in \mathbb{R}^n is the classic example.

Example (1) of Inner Products

In general there are lots of inner products on vector spaces (more on this later). But here are two examples that you should be familiar with:

The *dot product* in \mathbb{R}^n is the classic example. Can be extended to general finite dimensional vector spaces as follows:

Example (1) of Inner Products

In general there are lots of inner products on vector spaces (more on this later). But here are two examples that you should be familiar with:

The *dot product* in \mathbb{R}^n is the classic example. Can be extended to general finite dimensional vector spaces as follows:

Let $\beta = \{v_1, \dots, v_n\}$ be a basis for a finite dimensional vector space. Let $[\cdot]_\beta$ be its associated coordinate map. Then consider

$$(v, w) = [v]_\beta \cdot [w]_\beta.$$

Example (1) of Inner Products

In general there are lots of inner products on vector spaces (more on this later). But here are two examples that you should be familiar with:

The *dot product* in \mathbb{R}^n is the classic example. Can be extended to general finite dimensional vector spaces as follows:

Let $\beta = \{v_1, \dots, v_n\}$ be a basis for a finite dimensional vector space. Let $[\cdot]_\beta$ be its associated coordinate map. Then consider

$$(v, w) = [v]_\beta \cdot [w]_\beta.$$

That this has the properties of an inner product is evident from its definition involving the dot product.

Inner Products in FD Vector Spaces are (basically) the Dot Product

Let (\cdot, \cdot) be an inner product on a vector space of dimension n .

Inner Products in FD Vector Spaces are (basically) the Dot Product

Let (\cdot, \cdot) be an inner product on a vector space of dimension n .

Let $\beta = \{v_1, v_2, \dots, v_n\}$ be a basis for V .

Inner Products in FD Vector Spaces are (basically) the Dot Product

Let (\cdot, \cdot) be an inner product on a vector space of dimension n .

Let $\beta = \{v_1, v_2, \dots, v_n\}$ be a basis for V .

Compute $C_{ij} = (v_i, v_j)$. Note that this gives an $n \times n$ matrix C .

Inner Products in FD Vector Spaces are (basically) the Dot Product

Let (\cdot, \cdot) be an inner product on a vector space of dimension n .

Let $\beta = \{v_1, v_2, \dots, v_n\}$ be a basis for V .

Compute $C_{ij} = (v_i, v_j)$. Note that this gives an $n \times n$ matrix C .

Theorem

- ▶ C is symmetric.
- ▶ $(v, w) = [v]_{\beta} \cdot C[w]_{\beta}$.

Inner Products in FD Vector Spaces are (basically) the Dot Product

Let (\cdot, \cdot) be an inner product on a vector space of dimension n .

Let $\beta = \{v_1, v_2, \dots, v_n\}$ be a basis for V .

Compute $C_{ij} = (v_i, v_j)$. Note that this gives an $n \times n$ matrix C .

Theorem

- ▶ C is symmetric.
- ▶ $(v, w) = [v]_{\beta} \cdot C[w]_{\beta}$.

Note that symmetry is clear.

Proof

Write

$$v = \sum_{i=1}^n a_i v_i \text{ and } w = \sum_{j=1}^n b_j v_j.$$

Then

$$(v, w) = \left(\sum_{i=1}^n a_i v_i, \sum_{j=1}^n b_j v_j \right) = \sum_{i=1}^n \sum_{j=1}^n a_i b_j (v_i, v_j).$$

Proof

Write

$$v = \sum_{i=1}^n a_i v_i \text{ and } w = \sum_{j=1}^n b_j v_j.$$

Then

$$(v, w) = \left(\sum_{i=1}^n a_i v_i, \sum_{j=1}^n b_j v_j \right) = \sum_{i=1}^n \sum_{j=1}^n a_i b_j (v_i, v_j).$$

Use the definition of C :

$$(v, w) = \sum_{i=1}^n \sum_{j=1}^n a_i C_{ij} b_j = \sum_{i=1}^n a_i \left(\sum_{j=1}^n C_{ij} b_j \right).$$

More Proof

We have

$$(v, w) = \sum_{i=1}^n a_i \underbrace{\left(\sum_{j=1}^n C_{ij} b_j \right)}_{(C[w]_{\beta})_i}$$

More Proof

We have

$$(v, w) = \sum_{i=1}^n a_i \underbrace{\left(\sum_{j=1}^n C_{ij} b_j \right)}_{(C[w]_{\beta})_i} = \sum_{i=1}^n a_i (C[w]_{\beta})_i.$$

So

$$(v, w) = [v]_{\beta} \cdot C[w]_{\beta}.$$

The matrix C is called the *matrix of the inner product with respect to the basis β* .

We have met such a matrix before (and we may again!)

Let

$$f : \mathbb{R}^n \rightarrow \mathbb{R}$$

be a differentiable function.

We have met such a matrix before (and we may again!)

Let

$$f : \mathbb{R}^n \rightarrow \mathbb{R}$$

be a differentiable function. The *Hessian* of f is the matrix

$$(H_f)_{ij} = \frac{\partial^2 f}{\partial x_i \partial x_j}$$

We have met such a matrix before (and we may again!)

Let

$$f : \mathbb{R}^n \rightarrow \mathbb{R}$$

be a differentiable function. The *Hessian* of f is the matrix

$$(H_f)_{ij} = \frac{\partial^2 f}{\partial x_i \partial x_j}$$

The question “Does f have a local minimum” can be answered by asking if H_f is the matrix of an inner product!

Example (2b) of Inner Products

A second example of an inner product:

Example (2b) of Inner Products

A second example of an inner product: Let

$f, g \in P_n[a, b] = \{f : [a, b] \rightarrow \mathbb{R} : f \text{ a polynomial of degree at most } n\}.$

Define

$$(f, g) = \int_b^a fg \, dx.$$

Example (2b) of Inner Products

A second example of an inner product: Let

$f, g \in P_n[a, b] = \{f : [a, b] \rightarrow \mathbb{R} : f \text{ a polynomial of degree at most } n\}.$

Define

$$(f, g) = \int_b^a fg \, dx.$$

We should prove this is an inner product.

Exercise: Find the matrix for the inner product of $P_1[0, 1]$ using the standard basis $\{1, x\}$.

Example (2b) of Inner Products

A second example of an inner product: Let

$f, g \in P_n[a, b] = \{f : [a, b] \rightarrow \mathbb{R} : f \text{ a polynomial of degree at most } n\}.$

Define

$$(f, g) = \int_b^a fg \, dx.$$

We should prove this is an inner product.

Exercise: Find the matrix for the inner product of $P_1[0, 1]$ using the standard basis $\{1, x\}$.

Sln:

$$C = \begin{bmatrix} 1 & 1/2 \\ 1/2 & 1/3 \end{bmatrix}$$

Lengths, Distances, and Angles

If V is a vector space with an inner product (\cdot, \cdot) then we have the following definitions.

Lengths, Distances, and Angles

If V is a vector space with an inner product (\cdot, \cdot) then we have the following definitions.

Definition

- ▶ The **norm** of v is

$$\|v\| := \sqrt{(v, v)}.$$

- ▶ The **distance** between vectors v and w is

$$d(v, w) := \|v - w\|.$$

- ▶ The **angle** between vectors v and w is

$$\cos^{-1} \frac{(v, w)}{\|v\| \|w\|}.$$

Example (2b) of Inner Products

Extend our example to an *infinite dimensional* vector space:

Example (2b) of Inner Products

Extend our example to an *infinite dimensional* vector space: Let

$$f, g \in C[a, b] = \{f : [a, b] \rightarrow \mathbb{R} : f \text{ is continuous}\}.$$

Example (2b) of Inner Products

Extend our example to an *infinite dimensional* vector space: Let

$$f, g \in C[a, b] = \{f : [a, b] \rightarrow \mathbb{R} : f \text{ is continuous}\}.$$

Define

$$(f, g) = \int_a^b fg \, dx.$$

Example (2b) of Inner Products

Extend our example to an *infinite dimensional* vector space: Let

$$f, g \in C[a, b] = \{f : [a, b] \rightarrow \mathbb{R} : f \text{ is continuous}\}.$$

Define

$$(f, g) = \int_a^b fg \, dx.$$

Exercise: Show that $\sin(x)$ and $\cos(x)$ are perpendicular in $C[0, 2\pi]$ under the inner product above.

Example (2b) of Inner Products

Extend our example to an *infinite dimensional* vector space: Let

$$f, g \in C[a, b] = \{f : [a, b] \rightarrow \mathbb{R} : f \text{ is continuous}\}.$$

Define

$$(f, g) = \int_a^b fg \, dx.$$

Exercise: Show that $\sin(x)$ and $\cos(x)$ are perpendicular in $C[0, 2\pi]$ under the inner product above.

Sln:

$$\int_0^{2\pi} \cos x \sin x \, dx = \left. \frac{1}{2} \sin^2(x) \right|_0^{2\pi} = 0.$$

Orthogonal Implies Linear Independence

An *orthogonal set* of vectors is a set

$$\{v_1, v_2, \dots, v_n\}$$

with the property that

$$(v_i, v_j) = 0, \quad i \neq j.$$

Orthogonal Implies Linear Independence

An *orthogonal set* of vectors is a set

$$\{v_1, v_2, \dots, v_n\}$$

with the property that

$$(v_i, v_j) = 0, \quad i \neq j.$$

Theorem

A finite orthogonal set of non-zero vectors is linearly independent.

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$.

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$. We want to show that $a_i = 0$.

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$. We want to show that $a_i = 0$. Take the inner product

$$\left(v_j, \sum_{i=1}^n a_i v_i \right).$$

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$. We want to show that $a_i = 0$. Take the inner product

$$\left(v_j, \sum_{i=1}^n a_i v_i \right).$$

Note that this must be zero.

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$. We want to show that $a_i = 0$. Take the inner product

$$\left(v_j, \sum_{i=1}^n a_i v_i \right).$$

Note that this must be zero.

Also we have

$$\left(v_j, \sum_{i=1}^n a_i v_i \right) = \sum_{i=1}^n a_i (v_j, v_i)$$

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$. We want to show that $a_i = 0$. Take the inner product

$$\left(v_j, \sum_{i=1}^n a_i v_i \right).$$

Note that this must be zero.

Also we have

$$\left(v_j, \sum_{i=1}^n a_i v_i \right) = \sum_{i=1}^n a_i (v_j, v_i) = a_j (v_j, v_j).$$

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$. We want to show that $a_j = 0$. Take the inner product

$$\left(v_j, \sum_{i=1}^n a_i v_i \right).$$

Note that this must be zero.

Also we have

$$\left(v_j, \sum_{i=1}^n a_i v_i \right) = \sum_{i=1}^n a_i (v_j, v_i) = a_j (v_j, v_j).$$

So we have $0 = a_j (v_j, v_j)$.

Proof.

Suppose $\sum_{i=1}^n a_i v_i = 0$. We want to show that $a_j = 0$. Take the inner product

$$\left(v_j, \sum_{i=1}^n a_i v_i \right).$$

Note that this must be zero.

Also we have

$$\left(v_j, \sum_{i=1}^n a_i v_i \right) = \sum_{i=1}^n a_i (v_j, v_i) = a_j (v_j, v_j).$$

So we have $0 = a_j (v_j, v_j)$. Now, $(v_j, v_j) \neq 0$ therefore $a_j = 0$. □

Summary

Inner products allow us to do geometry in places which do not seem “natural.”

Summary

Inner products allow us to do geometry in places which do not seem “natural.” Two important geometry theorems:

Theorem (Cauchy-Schwartz)

For any two vectors in an inner product space V we have

$$|(u, v)| \leq \|u\| \|v\|.$$

Summary

Inner products allow us to do geometry in places which do not seem “natural.” Two important geometry theorems:

Theorem (Cauchy-Schwartz)

For any two vectors in an inner product space V we have

$$|(u, v)| \leq \|u\| \|v\|.$$

Theorem (Triangle)

For any two vectors in an inner product space V we have

$$\|u + v\| \leq \|u\| + \|v\|.$$

Summary

Inner products allow us to do geometry in places which do not seem “natural.” Two important geometry theorems:

Theorem (Cauchy-Schwartz)

For any two vectors in an inner product space V we have

$$|(u, v)| \leq \|u\| \|v\|.$$

Theorem (Triangle)

For any two vectors in an inner product space V we have

$$\|u + v\| \leq \|u\| + \|v\|.$$

These results rely ONLY on the properties of the inner products and not HOW you compute them or WHAT the vectors represent!

Summary

Inner products allow us to do geometry in places which do not seem “natural.” Two important geometry theorems:

Theorem (Cauchy-Schwartz)

For any two vectors in an inner product space V we have

$$|(u, v)| \leq \|u\| \|v\|.$$

Theorem (Triangle)

For any two vectors in an inner product space V we have

$$\|u + v\| \leq \|u\| + \|v\|.$$

These results rely ONLY on the properties of the inner products and not HOW you compute them or WHAT the vectors represent! They are important/useful for doing analytic geometry in inner product spaces.