

# Things left to do with Determinants

# Things left to do with Determinants

- ▶ Prove that cofactor expansion formula **IS** the determinant.

# Things left to do with Determinants

- ▶ Prove that cofactor expansion formula **IS** the determinant.
- ▶ Prove that the cofactor inverse formula **IS** the inverse.

# Things left to do with Determinants

- ▶ Prove that cofactor expansion formula **IS** the determinant.
- ▶ Prove that the cofactor inverse formula **IS** the inverse.
- ▶ Geometric interpretation of the determinant.

# The Cofactor Expansion Formula is the Determinant

Recall that if  $A = [a_{ij}]$  is an  $n \times n$  matrix, then the cofactor expansion formula across the  $i$ -th row is

$$f(A) = \sum_{j=1}^n a_{ij}(-1)^{i+j}f(M_{1j})$$

where  $M_{ij}$  is the  $(i,j)$ -th minor matrix.

# The Cofactor Expansion Formula is the Determinant

Recall that if  $A = [a_{ij}]$  is an  $n \times n$  matrix, then the cofactor expansion formula across the  $i$ -th row is

$$f(A) = \sum_{j=1}^n a_{ij}(-1)^{i+j}f(M_{1j})$$

where  $M_{ij}$  is the  $(i,j)$ -th minor matrix. In order to show that  $f = \det$  we need to show:

1.  $f(I) = 1$ .

# The Cofactor Expansion Formula is the Determinant

Recall that if  $A = [a_{ij}]$  is an  $n \times n$  matrix, then the cofactor expansion formula across the  $i$ -th row is

$$f(A) = \sum_{j=1}^n a_{ij}(-1)^{i+j}f(M_{1j})$$

where  $M_{ij}$  is the  $(i,j)$ -th minor matrix. In order to show that  $f = \det$  we need to show:

1.  $f(I) = 1$ .
2.  $f$  is *antisymmetric*.

# The Cofactor Expansion Formula is the Determinant

Recall that if  $A = [a_{ij}]$  is an  $n \times n$  matrix, then the cofactor expansion formula across the  $i$ -th row is

$$f(A) = \sum_{j=1}^n a_{ij}(-1)^{i+j}f(M_{1j})$$

where  $M_{ij}$  is the  $(i,j)$ -th minor matrix. In order to show that  $f = \det$  we need to show:

1.  $f(I) = 1$ .
2.  $f$  is *antisymmetric*.
3.  $f$  is *multilinear* across the rows.



# The Cofactor Expansion Formula is the Determinant

Recall that if  $A = [a_{ij}]$  is an  $n \times n$  matrix, then the cofactor expansion formula across the  $i$ -th row is

$$f(A) = \sum_{j=1}^n a_{ij}(-1)^{i+j}f(M_{1j})$$

where  $M_{ij}$  is the  $(i,j)$ -th minor matrix. In order to show that  $f = \det$  we need to show:

1.  $f(I) = 1$ .
2.  $f$  is *antisymmetric*.
3.  $f$  is *multilinear* across the rows.

If we can do this, then  $f = \det$  from uniqueness.

# The Cofactor Expansion Formula is the Determinant

Recall that if  $A = [a_{ij}]$  is an  $n \times n$  matrix, then the cofactor expansion formula across the  $i$ -th row is

$$f(A) = \sum_{j=1}^n a_{ij}(-1)^{i+j} f(M_{1j})$$

where  $M_{ij}$  is the  $(i, j)$ -th minor matrix. In order to show that  $f = \det$  we need to show:

1.  $f(I) = 1$ .
2.  $f$  is *antisymmetric*.
3.  $f$  is *multilinear* across the rows.

If we can do this, then  $f = \det$  from uniqueness.

We will NOT need to repeat the proof for column expansion since we already know  $\det(A) = \det(A^T)$ .

# The Cofactor Expansion Formula is the Determinant

Recall that if  $A = [a_{ij}]$  is an  $n \times n$  matrix, then the cofactor expansion formula across the  $i$ -th row is

$$f(A) = \sum_{j=1}^n a_{ij}(-1)^{i+j} f(M_{1j})$$

where  $M_{ij}$  is the  $(i, j)$ -th minor matrix. In order to show that  $f = \det$  we need to show:

1.  $f(I) = 1$ .
2.  $f$  is *antisymmetric*.
3.  $f$  is *multilinear* across the rows.

If we can do this, then  $f = \det$  from uniqueness.

We will NOT need to repeat the proof for column expansion since we already know  $\det(A) = \det(A^T)$ .

We will make one MAJOR assumption: We will assume that the expansion formula is row independent.

# Method of Proof: Induction

Our proof is based on the SIZE of the matrix.

# Method of Proof: Induction

Our proof is based on the SIZE of the matrix.

We will show that

- ▶ Cofactor expansion works for  $1 \times 1$  matrices.

# Method of Proof: Induction

Our proof is based on the SIZE of the matrix.

We will show that

- ▶ Cofactor expansion works for  $1 \times 1$  matrices.
- ▶ If cofactor expansion works for small matrices, then it must work for bigger ones too.

# Method of Proof: Induction

Our proof is based on the SIZE of the matrix.

We will show that

- ▶ Cofactor expansion works for  $1 \times 1$  matrices.
- ▶ If cofactor expansion works for small matrices, then it must work for bigger ones too.

Therefore:

Since it works for  $1 \times 1$  it must also work for  $2 \times 2$ .

# Method of Proof: Induction

Our proof is based on the SIZE of the matrix.

We will show that

- ▶ Cofactor expansion works for  $1 \times 1$  matrices.
- ▶ If cofactor expansion works for small matrices, then it must work for bigger ones too.

Therefore:

Since it works for  $1 \times 1$  it must also work for  $2 \times 2$ .

Since it works for  $2 \times 2$  it must also work for  $3 \times 3$ .



# Method of Proof: Induction

Our proof is based on the SIZE of the matrix.

We will show that

- ▶ Cofactor expansion works for  $1 \times 1$  matrices.
- ▶ If cofactor expansion works for small matrices, then it must work for bigger ones too.

Therefore:

Since it works for  $1 \times 1$  it must also work for  $2 \times 2$ .

Since it works for  $2 \times 2$  it must also work for  $3 \times 3$ .

Since it works for  $3 \times 3$  it must also work for  $4 \times 4$ .

# Method of Proof: Induction

Our proof is based on the SIZE of the matrix.

We will show that

- ▶ Cofactor expansion works for  $1 \times 1$  matrices.
- ▶ If cofactor expansion works for small matrices, then it must work for bigger ones too.

Therefore:

Since it works for  $1 \times 1$  it must also work for  $2 \times 2$ .

Since it works for  $2 \times 2$  it must also work for  $3 \times 3$ .

Since it works for  $3 \times 3$  it must also work for  $4 \times 4$ . And so on.

# Method of Proof: Induction

Our proof is based on the SIZE of the matrix.

We will show that

- ▶ Cofactor expansion works for  $1 \times 1$  matrices.
- ▶ If cofactor expansion works for small matrices, then it must work for bigger ones too.

Therefore:

Since it works for  $1 \times 1$  it must also work for  $2 \times 2$ .

Since it works for  $2 \times 2$  it must also work for  $3 \times 3$ .

Since it works for  $3 \times 3$  it must also work for  $4 \times 4$ . And so on.

So it should work for all sizes!

# The Cofactor Expansion Formula Works for $n = 1$ .

If  $A \in \mathbb{R}^{n \times n}$  then  $A = [a]$  and  $f(A) = a$ .

# The Cofactor Expansion Formula Works for $n = 1$ .

If  $A \in \mathbb{R}^{n \times n}$  then  $A = [a]$  and  $f(A) = a$ .

1.  $f(I) = f([1]) = 1$ .

# The Cofactor Expansion Formula Works for $n = 1$ .

If  $A \in \mathbb{R}^{n \times n}$  then  $A = [a]$  and  $f(A) = a$ .

1.  $f(I) = f([1]) = 1$ .
2.  $f$  is antisymmetric because there are no rows to switch.

# The Cofactor Expansion Formula Works for $n = 1$ .

If  $A \in \mathbb{R}^{n \times n}$  then  $A = [a]$  and  $f(A) = a$ .

1.  $f(I) = f([1]) = 1$ .
2.  $f$  is antisymmetric because there are no rows to switch.
3.  $f$  is linear since:
  - 3.1  $f([ka]) = ka = k(f([a]))$ .

# The Cofactor Expansion Formula Works for $n = 1$ .

If  $A \in \mathbb{R}^{n \times n}$  then  $A = [a]$  and  $f(A) = a$ .

1.  $f(I) = f([1]) = 1$ .
2.  $f$  is antisymmetric because there are no rows to switch.
3.  $f$  is linear since:
  - 3.1  $f([ka]) = ka = k(f([a]))$ .
  - 3.2  $f([a + b]) = a + b = f([a]) + f([b])$ .



# The Cofactor Expansion Formula Works for $n = 1$ .

If  $A \in \mathbb{R}^{n \times n}$  then  $A = [a]$  and  $f(A) = a$ .

1.  $f(I) = f([1]) = 1$ .
2.  $f$  is antisymmetric because there are no rows to switch.
3.  $f$  is linear since:
  - 3.1  $f([ka]) = ka = k(f([a]))$ .
  - 3.2  $f([a + b]) = a + b = f([a]) + f([b])$ .

So: Cofactor expansion works for  $1 \times 1$  matrices.

Works for small matrix means it also works for a big one

Do this on the board

## Next: The Cofactor Inverse Formula gives the Inverse

Let  $A = [a_{ij}]$  be non-singular and define

$$B = \frac{1}{|A|} \begin{bmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ A_{1n} & A_{2n} & \dots & A_{nn} \end{bmatrix}$$

where

$$A_{ij} = (-1)^{i+j} |M_{ij}|.$$

## Next: The Cofactor Inverse Formula gives the Inverse

Let  $A = [a_{ij}]$  be non-singular and define

$$B = \frac{1}{|A|} \begin{bmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ A_{1n} & A_{2n} & \dots & A_{nn} \end{bmatrix}$$

where

$$A_{ij} = (-1)^{i+j} |M_{ij}|.$$

Lets show  $AB = I$ .

## Compute $AB$

$$AB = \frac{1}{|A|} \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ A_{1n} & A_{2n} & \dots & A_{nn} \end{bmatrix}$$

Diagonal elements:

$$(AB)_{ii} = \frac{1}{|A|} \underbrace{\sum_{k=1}^n a_{ik} A_{ik}}$$

## Compute $AB$

$$AB = \frac{1}{|A|} \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ A_{1n} & A_{2n} & \dots & A_{nn} \end{bmatrix}$$

Diagonal elements:

$$(AB)_{ii} = \frac{1}{|A|} \underbrace{\sum_{k=1}^n a_{ik} A_{ik}}_{|A|}$$

## Compute $AB$

$$AB = \frac{1}{|A|} \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ A_{1n} & A_{2n} & \dots & A_{nn} \end{bmatrix}$$

Diagonal elements:

$$(AB)_{ii} = \frac{1}{|A|} \underbrace{\sum_{k=1}^n a_{ik} A_{ik}}_{|A|} = 1$$

non-diagonal elements:

$$(AB)_{ij} = \frac{1}{|A|} \sum_{k=1}^n a_{ik} A_{jk}$$

# Compute $AB$

$$AB = \frac{1}{|A|} \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ A_{1n} & A_{2n} & \dots & A_{nn} \end{bmatrix}$$

Diagonal elements:

$$(AB)_{ii} = \frac{1}{|A|} \underbrace{\sum_{k=1}^n a_{ik} A_{ik}}_{|A|} = 1$$

non-diagonal elements:

$$(AB)_{ij} = \frac{1}{|A|} \sum_{k=1}^n a_{ik} A_{jk} = 0 \text{ if } i \neq j$$



# Compute $AB$

$$AB = \frac{1}{|A|} \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ A_{1n} & A_{2n} & \dots & A_{nn} \end{bmatrix}$$

Diagonal elements:

$$(AB)_{ii} = \frac{1}{|A|} \underbrace{\sum_{k=1}^n a_{ik} A_{ik}}_{|A|} = 1$$

non-diagonal elements:

$$(AB)_{ij} = \frac{1}{|A|} \sum_{k=1}^n a_{ik} A_{jk} = 0 \text{ if } i \neq j$$

because the sum is the determinant of a matrix with two identical rows!

# The Determinant and Geometry (1)

# The Determinant and Geometry (1)

We can think of an  $n \times n$  matrix as a list of  $n \times 1$  vectors:

$$\begin{bmatrix} v_1 & v_2 & \cdots & v_n \end{bmatrix}.$$

If we plot the list of vectors in  $\mathbb{R}^n$  we can create a *hyperparallelepiped* (HPP)!

# The Determinant and Geometry (1)

We can think of an  $n \times n$  matrix as a list of  $n \times 1$  vectors:

$$\begin{bmatrix} v_1 & v_2 & \cdots & v_n \end{bmatrix}.$$

If we plot the list of vectors in  $\mathbb{R}^n$  we can create a *hyperparallelepiped* (HPP)! (In two dimensions this is a parallelogram.)

# The Determinant and Geometry (1)

We can think of an  $n \times n$  matrix as a list of  $n \times 1$  vectors:

$$\begin{bmatrix} v_1 & v_2 & \cdots & v_n \end{bmatrix}.$$

If we plot the list of vectors in  $\mathbb{R}^n$  we can create a *hyperparallelepiped* (HPP)! (In two dimensions this is a parallelogram.)

Using this notion we can define a function  $\text{vol}_n : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$  by

$\text{vol}_n(A)$  = volume of the HPP defined by the columns of  $A$ .

# The Determinant and Geometry (1)

We can think of an  $n \times n$  matrix as a list of  $n \times 1$  vectors:

$$\begin{bmatrix} v_1 & v_2 & \cdots & v_n \end{bmatrix}.$$

If we plot the list of vectors in  $\mathbb{R}^n$  we can create a *hyperparallelepiped* (HPP)! (In two dimensions this is a parallelogram.)

Using this notion we can define a function  $vol_n : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$  by

$vol_n(A)$  = volume of the HPP defined by the columns of  $A$ .

It turns out that  $vol_n(A) = |\det(A)|$ !

# The Determinant and Geometry (2)

## The Determinant and Geometry (2)

Let  $A$  be an  $n \times n$  matrix and consider  $f_A : \mathbb{R}^n \rightarrow \mathbb{R}^n$  by  $f_A(x) = Ax$ .



## The Determinant and Geometry (2)

Let  $A$  be an  $n \times n$  matrix and consider  $f_A : \mathbb{R}^n \rightarrow \mathbb{R}^n$  by  $f_A(x) = Ax$ . Then for a set  $S \subset \mathbb{R}^n$  we have:

$$\text{vol}_n(f(S)) = |\det(A)| \text{vol}_n(S).$$

## Example

Let  $S$  be the unit circle in  $\mathbb{R}^2$ . The matrix

$$A = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}$$

will make  $f_A(S)$  an ellipse with semi-radii  $a$  and  $b$ .

## Example

Let  $S$  be the unit circle in  $\mathbb{R}^2$ . The matrix

$$A = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}$$

will make  $f_A(S)$  an ellipse with semi-radii  $a$  and  $b$ .

The area of the ellipse is

$$\text{vol}_2(f(S)) = \det(A)\text{vol}_2(S) = \pi ab.$$

## Example

Let  $S$  be the unit circle in  $\mathbb{R}^2$ . The matrix

$$A = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}$$

will make  $f_A(S)$  an ellipse with semi-radii  $a$  and  $b$ .

The area of the ellipse is

$$\text{vol}_2(f(S)) = \det(A)\text{vol}_2(S) = \pi ab.$$

Exercise: What is the volume of an ellipsoid with radii  $a$ ,  $b$ , and  $c$ ?