

## exercises

Use ONLY ELIMINATION to solve the following

1.

$$x_1 - 3x_2 = -7$$

$$2x_1 + x_2 = 7$$

2.

$$x - 3y = -7$$

$$2x - 6y = 7$$

3.

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

# Answers

1.  $(2, 3)$
2. System is not consistent.
3.  $(1, -2, 3)$ .

## Exercise (from text)

A manufacturer makes three types of chemicals:  $A$ ,  $B$ ,  $C$ .

- ▶ A ton of  $A$  needs 2 hours to refine and 2 hours to package.
- ▶ A ton of  $B$  needs 3 hours to refine and 2 hours to package.
- ▶ A ton of  $C$  needs 4 hours to refine and 3 hours to package.

The refining machine is available 80 hours per week, meanwhile the packing machine is available 60 hours per week. How much of  $A$ ,  $B$ , and  $C$  can be manufactured in a week?

1. Give an appropriate system of equations which represents the scenario given in the problem.
2. In your opinion (and without doing any computations) what types of solution sets are possible?

# Solutions

1. Let  $a$  be the amount (in tons) of chemical  $A$ ,  $b$  for  $B$ , and  $c$  for  $C$ . Then we have the system:

$$2a + 3b + 4c = 80$$

$$2a + 2b + 3c = 60$$

# Solutions

1. Let  $a$  be the amount (in tons) of chemical  $A$ ,  $b$  for  $B$ , and  $c$  for  $C$ . Then we have the system:

$$2a + 3b + 4c = 80$$

$$2a + 2b + 3c = 60$$

2. Interpret the linear equations as planes in  $\mathbb{R}^3$ . Then we can see that we have a consistent system with an infinite number of solutions since they are not parallel.

Let's do some elimination on the board over here ←.

Let's do some elimination on the board over here  $\longleftarrow$ .

One thing we notice as we go through the elimination steps is that (so long as we are careful) location determines variables and **ONLY THE COEFFICIENTS AND WHERE THEY ARE MATTER!**

# Matrices

The coefficients of a system of  $m$  equations in  $n$  variables can be described by a mathematical object called a **matrix**:



# Matrices

The coefficients of a system of  $m$  equations in  $n$  variables can be described by a mathematical object called a **matrix**:

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & a_{13}x_3 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & a_{23}x_3 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & & \vdots & & \vdots & & & & \vdots & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & a_{m3}x_3 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array}$$

# Matrices

The coefficients of a system of  $m$  equations in  $n$  variables can be described by a mathematical object called a **matrix**:

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & a_{13}x_3 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & a_{23}x_3 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & & \vdots & & \vdots & & & & \vdots & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & a_{m3}x_3 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array}$$



$$\text{Matrix} \longrightarrow \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix} \Leftrightarrow [a_{ij}].$$

## Example 1 of Matrices

A matrix of air travel times between several airports:

	To: JFK	To: LAX
From: JFK	0	6:20
From: LAX	5:15	0

## Example 1 of Matrices

A matrix of air travel times between several airports:


	To: JFK	To: LAX
From: JFK	0	6:20
From: LAX	5:15	0



$$\begin{bmatrix} 0 & 6.333 \\ 5.25 & 0 \end{bmatrix}$$


## Example 2: Incidence Matrix

An incidence matrix for a **graph** tells you if two nodes are adjacent:



## Example 2: Incidence Matrix

An incidence matrix for a **graph** tells you if two nodes are adjacent:



$$\begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

## Example 3: Markov Matrix

A Markov matrix tells you the probability of changing from one “state” to another:

## Example 3: Markov Matrix

A Markov matrix tells you the probability of changing from one “state” to another:

- ▶ Markov matrix for a “fair” coin:

$$\begin{bmatrix} 1/2 & 1/2 \\ 1/2 & 1/2 \end{bmatrix}$$



## Example 3: Markov Matrix

A Markov matrix tells you the probability of changing from one “state” to another:

- ▶ Markov matrix for a “fair” coin:

$$\begin{bmatrix} 1/2 & 1/2 \\ 1/2 & 1/2 \end{bmatrix}$$

Regardless if your coin is currently heads or tails, it has a 50-50 chance of changing if you flip it.

## Example 3: Markov Matrix

A Markov matrix tells you the probability of changing from one “state” to another:

- ▶ Markov matrix for a “fair” coin:

$$\begin{bmatrix} 1/2 & 1/2 \\ 1/2 & 1/2 \end{bmatrix}$$

Regardless if your coin is currently heads or tails, it has a 50-50 chance of changing if you flip it.

- ▶ Markov matrix for a “loaded” coin:

$$\begin{bmatrix} 1/10 & 1/10 \\ 9/10 & 9/10 \end{bmatrix}$$

# Definition and Standard Terminology

## Definition (Matrix)

Let  $m$  and  $n$  be positive integers. An  $m \times n$  **Matrix** is a rectangular array of elements (for us: real or complex numbers) arranged in  $m$  (horizontal) rows and  $n$  (vertical) columns:

$$M = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix}.$$

# Definition and Standard Terminology

## Definition (Matrix)

Let  $m$  and  $n$  be positive integers. An  $m \times n$  **Matrix** is a rectangular array of elements (for us: real or complex numbers) arranged in  $m$  (horizontal) rows and  $n$  (vertical) columns:

$$M = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix}.$$

In the above,  $a_{ij}$  is called an **element/entry** of  $M$  or the  $(i,j)$ **th element/entry** of  $M$ . Here  $i$  indicates the *row* of  $M$  in which the element is located and  $j$  indicates the *column*.

# Definition and Standard Terminology

## Definition (Matrix)

Let  $m$  and  $n$  be positive integers. An  $m \times n$  **Matrix** is a rectangular array of elements (for us: real or complex numbers) arranged in  $m$  (horizontal) rows and  $n$  (vertical) columns:

$$M = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix}.$$

In the above,  $a_{ij}$  is called an **element/entry** of  $M$  or the  $(i,j)$ **th element/entry** of  $M$ . Here  $i$  indicates the *row* of  $M$  in which the element is located and  $j$  indicates the *column*.

# Notation

The set of all  $m \times n$  matrices that can be made using real numbers as entries is  $\mathbb{R}^{m \times n}$ .

# Notation

The set of all  $m \times n$  matrices that can be made using real numbers as entries is  $\mathbb{R}^{m \times n}$ .

Write down an element of  $\mathbb{R}^{2 \times 3}$ .

# Special Matrices

- ▶ If  $m = n$  then we say the matrix is **square**.



# Special Matrices

- ▶ If  $m = n$  then we say the matrix is **square**.  
Give an example of a  $3 \times 3$  square matrix using only  $0, 1, -1$  as entries.

# Special Matrices

- ▶ If  $m = n$  then we say the matrix is **square**.  
Give an example of a  $3 \times 3$  square matrix using only  $0, 1, -1$  as entries.
- ▶ If a matrix is square and  $a_{ij} = 0$  when  $i \neq j$  then the matrix is **diagonal**.

# Special Matrices

- ▶ If  $m = n$  then we say the matrix is **square**.  
Give an example of a  $3 \times 3$  square matrix using only  $0, 1, -1$  as entries.
- ▶ If a matrix is square and  $a_{ij} = 0$  when  $i \neq j$  then the matrix is **diagonal**.  
Give a diagonal element of a  $\mathbb{R}^{4 \times 4}$  which has only  $0, 1, -1$  as entries.

# Special Matrices

- ▶ If  $m = n$  then we say the matrix is **square**.  
Give an example of a  $3 \times 3$  square matrix using only  $0, 1, -1$  as entries.
- ▶ If a matrix is square and  $a_{ij} = 0$  when  $i \neq j$  then the matrix is **diagonal**.  
Give a diagonal element of a  $\mathbb{R}^{4 \times 4}$  which has only  $0, 1, -1$  as entries.
- ▶ A diagonal matrix with the property that  $a_{ii} = 1$  is called an **identity** matrix. This matrix is often written as  $I$  or  $I_n$ .

# Special Matrices

- ▶ If  $m = n$  then we say the matrix is **square**.  
Give an example of a  $3 \times 3$  square matrix using only  $0, 1, -1$  as entries.
- ▶ If a matrix is square and  $a_{ij} = 0$  when  $i \neq j$  then the matrix is **diagonal**.  
Give a diagonal element of a  $\mathbb{R}^{4 \times 4}$  which has only  $0, 1, -1$  as entries.
- ▶ A diagonal matrix with the property that  $a_{ii} = 1$  is called an **identity** matrix. This matrix is often written as  $I$  or  $I_n$ .  
Write down the identity in  $\mathbb{R}^{2 \times 2}$ .

# Special Matrices

- ▶ If  $m = n$  then we say the matrix is **square**.  
Give an example of a  $3 \times 3$  square matrix using only  $0, 1, -1$  as entries.
- ▶ If a matrix is square and  $a_{ij} = 0$  when  $i \neq j$  then the matrix is **diagonal**.  
Give a diagonal element of a  $\mathbb{R}^{4 \times 4}$  which has only  $0, 1, -1$  as entries.
- ▶ A diagonal matrix with the property that  $a_{ii} = 1$  is called an **identity** matrix. This matrix is often written as  $I$  or  $I_n$ .  
Write down the identity in  $\mathbb{R}^{2 \times 2}$ .
- ▶ Elements of  $\mathbb{R}^{m \times 1}$  are called **vectors** and elements of  $\mathbb{R}^{1 \times n}$  are called **covectors**.

# Special Matrices

- ▶ If  $m = n$  then we say the matrix is **square**.  
Give an example of a  $3 \times 3$  square matrix using only  $0, 1, -1$  as entries.
- ▶ If a matrix is square and  $a_{ij} = 0$  when  $i \neq j$  then the matrix is **diagonal**.  
Give a diagonal element of a  $\mathbb{R}^{4 \times 4}$  which has only  $0, 1, -1$  as entries.
- ▶ A diagonal matrix with the property that  $a_{ii} = 1$  is called an **identity** matrix. This matrix is often written as  $I$  or  $I_n$ .  
Write down the identity in  $\mathbb{R}^{2 \times 2}$ .
- ▶ Elements of  $\mathbb{R}^{m \times 1}$  are called **vectors** and elements of  $\mathbb{R}^{1 \times n}$  are called **covectors**. In general vectors are *columns* and covectors are *rows*.

## Remark

Technically we are using the term *vector* rather colloquially. For example, we will see later in the course that vectors, covectors, and even matrices themselves can be thought of as vectors in a more abstract way.



# Operations on Matrices

Matrices have natural algebraic operations.

- ▶ *Scalar Multiplication*: Let  $r \in \mathbb{R}$  and  $A = [a_{ij}] \in \mathbb{R}^{m \times n}$ . Then the matrix  $rA := [ra_{ij}] \in \mathbb{R}^{m \times n}$ .

# Operations on Matrices

Matrices have natural algebraic operations.

- *Scalar Multiplication*: Let  $r \in \mathbb{R}$  and  $A = [a_{ij}] \in \mathbb{R}^{m \times n}$ . Then the matrix  $rA := [ra_{ij}] \in \mathbb{R}^{m \times n}$ .

$$3 \begin{bmatrix} 2 & 3 \\ 1 & 0 \\ -5 & 1 \end{bmatrix} =$$

# Operations on Matrices

Matrices have natural algebraic operations.

- *Scalar Multiplication*: Let  $r \in \mathbb{R}$  and  $A = [a_{ij}] \in \mathbb{R}^{m \times n}$ . Then the matrix  $rA := [ra_{ij}] \in \mathbb{R}^{m \times n}$ .

$$3 \begin{bmatrix} 2 & 3 \\ 1 & 0 \\ -5 & 1 \end{bmatrix} = \begin{bmatrix} 6 & 9 \\ 3 & 0 \\ -15 & 3 \end{bmatrix}$$

# Operations on Matrices

Matrices have natural algebraic operations.

- ▶ *Scalar Multiplication:* Let  $r \in \mathbb{R}$  and  $A = [a_{ij}] \in \mathbb{R}^{m \times n}$ . Then the matrix  $rA := [ra_{ij}] \in \mathbb{R}^{m \times n}$ .

$$3 \begin{bmatrix} 2 & 3 \\ 1 & 0 \\ -5 & 1 \end{bmatrix} = \begin{bmatrix} 6 & 9 \\ 3 & 0 \\ -15 & 3 \end{bmatrix}$$

- ▶ *Matrix Addition:* Let  $A = [a_{ij}]$  and  $B = [b_{ij}]$  both be matrices of the same type. (That is, both  $A, B \in \mathbb{R}^{n \times m}$ ). Then  $A + B := [a_{ij} + b_{ij}]$ .

# Operations on Matrices

Matrices have natural algebraic operations.

- ▶ *Scalar Multiplication:* Let  $r \in \mathbb{R}$  and  $A = [a_{ij}] \in \mathbb{R}^{m \times n}$ . Then the matrix  $rA := [ra_{ij}] \in \mathbb{R}^{m \times n}$ .

$$3 \begin{bmatrix} 2 & 3 \\ 1 & 0 \\ -5 & 1 \end{bmatrix} = \begin{bmatrix} 6 & 9 \\ 3 & 0 \\ -15 & 3 \end{bmatrix}$$

- ▶ *Matrix Addition:* Let  $A = [a_{ij}]$  and  $B = [b_{ij}]$  both be matrices of the same type. (That is, both  $A, B \in \mathbb{R}^{n \times m}$ ). Then  $A + B := [a_{ij} + b_{ij}]$ .

$$\begin{bmatrix} 2 & 3 \\ 1 & 0 \\ -5 & 1 \end{bmatrix} + \begin{bmatrix} -1 & 2 \\ 0 & -1 \\ 8 & 0 \end{bmatrix} =$$

# Operations on Matrices

Matrices have natural algebraic operations.

- ▶ *Scalar Multiplication:* Let  $r \in \mathbb{R}$  and  $A = [a_{ij}] \in \mathbb{R}^{m \times n}$ . Then the matrix  $rA := [ra_{ij}] \in \mathbb{R}^{m \times n}$ .

$$3 \begin{bmatrix} 2 & 3 \\ 1 & 0 \\ -5 & 1 \end{bmatrix} = \begin{bmatrix} 6 & 9 \\ 3 & 0 \\ -15 & 3 \end{bmatrix}$$

- ▶ *Matrix Addition:* Let  $A = [a_{ij}]$  and  $B = [b_{ij}]$  both be matrices of the same type. (That is, both  $A, B \in \mathbb{R}^{n \times m}$ ). Then  $A + B := [a_{ij} + b_{ij}]$ .

$$\begin{bmatrix} 2 & 3 \\ 1 & 0 \\ -5 & 1 \end{bmatrix} + \begin{bmatrix} -1 & 2 \\ 0 & -1 \\ 8 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 5 \\ 1 & -1 \\ 3 & 1 \end{bmatrix}$$

# Exercise: Linear Combinations

## Exercise

Compute the following **linear combination** of matrices

$$\begin{bmatrix} 2 & -1 & 2 \\ -1 & 0 & 4 \\ -2 & 5 & 2 \end{bmatrix} - 3I$$

# Exercise: Linear Combinations

## Exercise

Compute the following **linear combination** of matrices

$$\begin{bmatrix} 2 & -1 & 2 \\ -1 & 0 & 4 \\ -2 & 5 & 2 \end{bmatrix} - 3I$$

where  $I$  is the identity matrix



# Exercise: Linear Combinations

## Exercise

Compute the following **linear combination** of matrices

$$\begin{bmatrix} 2 & -1 & 2 \\ -1 & 0 & 4 \\ -2 & 5 & 2 \end{bmatrix} - 3I$$

where  $I$  is the identity matrix in  $\mathbb{R}^{3 \times 3}$ .

# Exercise: Linear Combinations

## Exercise

Compute the following **linear combination** of matrices

$$\begin{bmatrix} 2 & -1 & 2 \\ -1 & 0 & 4 \\ -2 & 5 & 2 \end{bmatrix} - 3I$$

where  $I$  is the identity matrix in  $\mathbb{R}^{3 \times 3}$ .

Answer:

$$\begin{bmatrix} -1 & -1 & 2 \\ -1 & -3 & 4 \\ -2 & 5 & -1 \end{bmatrix}$$

# Transpose

Let  $A = [a_{ij}] \in \mathbb{R}^{n \times m}$ . Then the **transpose** of  $A$  is the matrix  $A^T := [a_{ij}^T] \in \mathbb{R}^{m \times n}$  where

$$a_{ij}^T = a_{ji}.$$

# Transpose

Let  $A = [a_{ij}] \in \mathbb{R}^{n \times m}$ . Then the **transpose** of  $A$  is the matrix  $A^T := [a_{ij}^T] \in \mathbb{R}^{m \times n}$  where

$$a_{ij}^T = a_{ji}.$$

The transpose operation essentially “flips” the matrix along its main diagonal.

# Transpose

Let  $A = [a_{ij}] \in \mathbb{R}^{n \times m}$ . Then the **transpose** of  $A$  is the matrix  $A^T := [a_{ij}^T] \in \mathbb{R}^{m \times n}$  where

$$a_{ij}^T = a_{ji}.$$

The transpose operation essentially “flips” the matrix along its main diagonal.

Use the words “vector”, “covector, ” and “transpose” in a sentence:

# Transpose

Let  $A = [a_{ij}] \in \mathbb{R}^{n \times m}$ . Then the **transpose** of  $A$  is the matrix  $A^T := [a_{ij}^T] \in \mathbb{R}^{m \times n}$  where

$$a_{ij}^T = a_{ji}.$$

The transpose operation essentially “flips” the matrix along its main diagonal.

Use the words “vector”, “covector, ” and “transpose” in a sentence:

*The transpose of a vector is a covector.*

# Transpose

Let  $A = [a_{ij}] \in \mathbb{R}^{n \times m}$ . Then the **transpose** of  $A$  is the matrix  $A^T := [a_{ij}^T] \in \mathbb{R}^{m \times n}$  where

$$a_{ij}^T = a_{ji}.$$

The transpose operation essentially “flips” the matrix along its main diagonal.

Use the words “vector”, “covector, ” and “transpose” in a sentence:

*The transpose of a vector is a covector.*

What is  $(A^T)^T$ ?

# Transpose

Let  $A = [a_{ij}] \in \mathbb{R}^{n \times m}$ . Then the **transpose** of  $A$  is the matrix  $A^T := [a_{ij}^T] \in \mathbb{R}^{m \times n}$  where

$$a_{ij}^T = a_{ji}.$$

The transpose operation essentially “flips” the matrix along its main diagonal.

Use the words “vector”, “covector, ” and “transpose” in a sentence:

*The transpose of a vector is a covector.*

What is  $(A^T)^T$ ?  $(A^T)^T = A$ .



# Transpose

Let  $A = [a_{ij}] \in \mathbb{R}^{n \times m}$ . Then the **transpose** of  $A$  is the matrix  $A^T := [a_{ij}^T] \in \mathbb{R}^{m \times n}$  where

$$a_{ij}^T = a_{ji}.$$

The transpose operation essentially “flips” the matrix along its main diagonal.

Use the words “vector”, “covector, ” and “transpose” in a sentence:

*The transpose of a vector is a covector.*

What is  $(A^T)^T$ ?  $(A^T)^T = A$ .

Let  $A \in \mathbb{R}^{2 \times 7}$ . How many rows and columns does  $A^T$  have?

# Transpose

Let  $A = [a_{ij}] \in \mathbb{R}^{n \times m}$ . Then the **transpose** of  $A$  is the matrix  $A^T := [a_{ij}^T] \in \mathbb{R}^{m \times n}$  where

$$a_{ij}^T = a_{ji}.$$

The transpose operation essentially “flips” the matrix along its main diagonal.

Use the words “vector”, “covector, ” and “transpose” in a sentence:

*The transpose of a vector is a covector.*

What is  $(A^T)^T$ ?  $(A^T)^T = A$ .

Let  $A \in \mathbb{R}^{2 \times 7}$ . How many rows and columns does  $A^T$  have?  
 $A^T \in \mathbb{R}^{7 \times 2}$ . So 7 rows and 2 columns.