

Exercise

For each of the following maps, decide if they are or are not linear.

1. $L : P(\mathbb{R}) \rightarrow P(\mathbb{R})$ by $p(x) \mapsto xp(x)$.

2. $L : P(\mathbb{R}) \rightarrow \mathbb{R}$ by $f \mapsto f(0)$.

3. $L : \mathbb{R} \rightarrow \mathbb{R}$ by $x \mapsto |x|$.

4. $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ by $\begin{bmatrix} a \\ b \\ c \end{bmatrix} \mapsto \begin{bmatrix} b \\ 2c \\ 0 \end{bmatrix}$.

Theorem

Let $T : V \rightarrow W$ be linear. Then

1. $L(0) = 0$.

Theorem

Let $T : V \rightarrow W$ be linear. Then

1. $L(0) = 0$. 0 is the identity in the appropriate vector space.

Theorem

Let $T : V \rightarrow W$ be linear. Then

1. $L(0) = 0$. 0 is the identity in the appropriate vector space.
2. $L(u - v) = L(u) - L(v)$ for all $u, v \in V$.

Theorem

Let $T : V \rightarrow W$ be linear. Then

1. $L(0) = 0$. 0 is the identity in the appropriate vector space.
2. $L(u - v) = L(u) - L(v)$ for all $u, v \in V$.

The first result is nice since it gives insight into the *kernel* of the operator!

Big Theorem: Basis vectors characterize linear maps

Suppose V has basis $\beta = \{v_1, \dots, v_n\}$.

Theorem

For any n vectors w_i in W there exists a unique linear map $T : V \rightarrow W$ where

$$T(v_i) = w_i$$

Big Theorem: Basis vectors characterize linear maps

Suppose V has basis $\beta = \{v_1, \dots, v_n\}$.

Theorem

For any n vectors w_i in W there exists a unique linear map $T : V \rightarrow W$ where

$$T(v_i) = w_i$$

This theorem says that if we define the image of a (finite) basis for a linear map, then we completely determine the entire map!

Big Theorem: Basis vectors characterize linear maps

Suppose V has basis $\beta = \{v_1, \dots, v_n\}$.

Theorem

For any n vectors w_i in W there exists a unique linear map $T : V \rightarrow W$ where

$$T(v_i) = w_i$$

This theorem says that if we define the image of a (finite) basis for a linear map, then we completely determine the entire map!

Proof.

For any $x \in V$ write $x = \sum_{i=1}^n \alpha_i v_i$. Then define

$$T(v) := \sum_{i=1}^n \alpha_i w_i.$$

Big Theorem: Basis vectors characterize linear maps

Suppose V has basis $\beta = \{v_1, \dots, v_n\}$.

Theorem

For any n vectors w_i in W there exists a unique linear map $T : V \rightarrow W$ where

$$T(v_i) = w_i$$

This theorem says that if we define the image of a (finite) basis for a linear map, then we completely determine the entire map!

Proof.

For any $x \in V$ write $x = \sum_{i=1}^n \alpha_i v_i$. Then define

$$T(v) := \sum_{i=1}^n \alpha_i w_i.$$

That this map is well defined and unique come from the fact that β is a basis!



Application

Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$.

Application

Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Let $\{e_1, \dots, e_n\}$ be the standard basis for \mathbb{R}^n .

Application

Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Let $\{e_1, \dots, e_n\}$ be the standard basis for \mathbb{R}^n .

Note that

$$L(e_i) = A_i \in \mathbb{R}^m \text{ is a column vector.}$$

Application

Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Let $\{e_1, \dots, e_n\}$ be the standard basis for \mathbb{R}^n .
Note that

$$L(e_i) = A_i \in \mathbb{R}^m \text{ is a column vector.}$$

Note also that for some matrix $A \in \mathbb{R}^{m \times n}$ the product

Ae_i is the i th column of the matrix A .

Application

Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Let $\{e_1, \dots, e_n\}$ be the standard basis for \mathbb{R}^n .
Note that

$$L(e_i) = A_i \in \mathbb{R}^m \text{ is a column vector.}$$

Note also that for some matrix $A \in \mathbb{R}^{m \times n}$ the product

Ae_i is the i th column of the matrix A .

Definition

The **standard matrix representing** L is the matrix

$$A = [A_1 \ A_2 \ \dots \ A_n] = [L(e_1) \ L(e_2) \ \dots \ L(e_n)].$$

Application

Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Let $\{e_1, \dots, e_n\}$ be the standard basis for \mathbb{R}^n .
Note that

$$L(e_i) = A_i \in \mathbb{R}^m \text{ is a column vector.}$$

Note also that for some matrix $A \in \mathbb{R}^{m \times n}$ the product

Ae_i is the i th column of the matrix A .

Definition

The **standard matrix representing** L is the matrix

$$A = [A_1 \ A_2 \ \dots \ A_n] = [L(e_1) \ L(e_2) \ \dots \ L(e_n)].$$

Theorem

With notation as above, the matrix A is the unique matrix with the property that

$$L(x) = Ax.$$

Exercise

Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ by

$$\begin{bmatrix} a \\ b \\ c \end{bmatrix} \mapsto \begin{bmatrix} a + 2b \\ 3b - 2c \end{bmatrix}$$

Find the standard matrix representing L .

Moral(s) and a Question

- ▶ All linear transforms from $\mathbb{R}^n \rightarrow \mathbb{R}^m$ can be represented by multiplication by a matrix in $\mathbb{R}^{m \times n}$.

Moral(s) and a Question

- ▶ All linear transforms from $\mathbb{R}^n \rightarrow \mathbb{R}^m$ can be represented by multiplication by a matrix in $\mathbb{R}^{m \times n}$.
- ▶ BUT ALSO: we know that if V is a finite dimensional vector space, then $V \simeq \mathbb{R}^n$!

Moral(s) and a Question

- ▶ All linear transforms from $\mathbb{R}^n \rightarrow \mathbb{R}^m$ can be represented by multiplication by a matrix in $\mathbb{R}^{m \times n}$.
- ▶ BUT ALSO: we know that if V is a finite dimensional vector space, then $V \simeq \mathbb{R}^n$!
- ▶ So what, if any, relation is there between a map of finite dimensional vector spaces and a matrix operation on the isomorphic real spaces?

Moral(s) and a Question

- ▶ All linear transforms from $\mathbb{R}^n \rightarrow \mathbb{R}^m$ can be represented by multiplication by a matrix in $\mathbb{R}^{m \times n}$.
- ▶ BUT ALSO: we know that if V is a finite dimensional vector space, then $V \simeq \mathbb{R}^n$!
- ▶ So what, if any, relation is there between a map of finite dimensional vector spaces and a matrix operation on the isomorphic real spaces?

But before that, some definitions....

Definitions

Let $L : V \rightarrow W$ be a linear map.

Definition

- ▶ The **kernel** of L is the set

$$\ker(L) = \{v \in V : L(v) = 0\}.$$

Definitions

Let $L : V \rightarrow W$ be a linear map.

Definition

- ▶ The **kernel** of L is the set

$$\ker(L) = \{v \in V : L(v) = 0\}.$$

- ▶ The **range** of L is the set

$$\text{range}(L) = \{w \in W : \exists v \in V, L(v) = w\}.$$

Definitions

Let $L : V \rightarrow W$ be a linear map.

Definition

- ▶ The **kernel** of L is the set

$$\ker(L) = \{v \in V : L(v) = 0\}.$$

- ▶ The **range** of L is the set

$$\text{range}(L) = \{w \in W : \exists v \in V, L(v) = w\}.$$

Theorem

$\ker(L)$ is a subspace of V and $\text{range}(L)$ is a subspace of W .

Exercise

Since $\ker(L)$ is a subspace of V , it must have a basis.

Exercise

Since $\ker(L)$ is a subspace of V , it must have a basis.

Exercise: Let $L_A : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ by multiplication by

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

Find $\ker(L_A)$ and give a basis for it.

Exercise

Since $\ker(L)$ is a subspace of V , it must have a basis.

Exercise: Let $L_A : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ by multiplication by

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

Find $\ker(L_A)$ and give a basis for it.

Solution: We need to solve the homogeneous problem $Ax = 0$.

Exercise

Since $\ker(L)$ is a subspace of V , it must have a basis.

Exercise: Let $L_A : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ by multiplication by

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

Find $\ker(L_A)$ and give a basis for it.

Solution: We need to solve the homogeneous problem $Ax = 0$.

The solution is given by

$$x_1 = -x_3$$

$$x_2 = -x_3$$

$$x_3 = x_3$$

Exercise

Since $\ker(L)$ is a subspace of V , it must have a basis.

Exercise: Let $L_A : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ by multiplication by

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

Find $\ker(L_A)$ and give a basis for it.

Solution: We need to solve the homogeneous problem $Ax = 0$.

The solution is given by

$$x_1 = -x_3$$

$$x_2 = -x_3$$

$$x_3 = x_3$$

So

$$\ker(L_A) = \text{span} \left\{ \begin{bmatrix} -1 \\ -1 \\ 1 \end{bmatrix} \right\}.$$

Characterization of Kernel

Note that in the prior example, the kernel of L_A is the solutions set to the homogeneous problem $Ax = 0$:

Characterization of Kernel

Note that in the prior example, the kernel of L_A is the solutions set to the homogeneous problem $Ax = 0$:

$$\ker(L_A) = \text{null}(A).$$

Characterization of Kernel

Note that in the prior example, the kernel of L_A is the solutions set to the homogeneous problem $Ax = 0$:

$$\ker(L_A) = \text{null}(A).$$

So we already know how to solve this problem in general!

Characterization of Kernel

Note that in the prior example, the kernel of L_A is the solutions set to the homogeneous problem $Ax = 0$:

$$\ker(L_A) = \text{null}(A).$$

So we already know how to solve this problem in general!

So let's look at non-matrix examples too:

Exercise

Let $L : P_2(\mathbb{R}) \rightarrow \mathbb{R}$ by $p \mapsto \int_0^1 p \, dx$. Find the kernel of L , a basis for it, and the dimension for the space.

Exercise

Let $L : P_2(\mathbb{R}) \rightarrow \mathbb{R}$ by $p \mapsto \int_0^1 p \, dx$. Find the kernel of L , a basis for it, and the dimension for the space.

Solution:

Exercise

Let $L : P_2(\mathbb{R}) \rightarrow \mathbb{R}$ by $p \mapsto \int_0^1 p \, dx$. Find the kernel of L , a basis for it, and the dimension for the space.

Solution: We need to find all polynomials $p = ax^2 + bx + c$ so that

$$0 = \int_0^1 ax^2 + bx + c \, dx$$

Exercise

Let $L : P_2(\mathbb{R}) \rightarrow \mathbb{R}$ by $p \mapsto \int_0^1 p \, dx$. Find the kernel of L , a basis for it, and the dimension for the space.

Solution: We need to find all polynomials $p = ax^2 + bx + c$ so that

$$0 = \int_0^1 ax^2 + bx + c \, dx = \frac{a}{3} + \frac{b}{2} + c$$

So $a = -\frac{3b}{2} - 3c$

Exercise

Let $L : P_2(\mathbb{R}) \rightarrow \mathbb{R}$ by $p \mapsto \int_0^1 p \, dx$. Find the kernel of L , a basis for it, and the dimension for the space.

Solution: We need to find all polynomials $p = ax^2 + bx + c$ so that

$$0 = \int_0^1 ax^2 + bx + c \, dx = \frac{a}{3} + \frac{b}{2} + c$$

So $a = -\frac{3b}{2} - 3c$ and

$$p = \left(-\frac{3b}{2} - 3c \right) x^2 + bx + c$$

Exercise

Let $L : P_2(\mathbb{R}) \rightarrow \mathbb{R}$ by $p \mapsto \int_0^1 p \, dx$. Find the kernel of L , a basis for it, and the dimension for the space.

Solution: We need to find all polynomials $p = ax^2 + bx + c$ so that

$$0 = \int_0^1 ax^2 + bx + c \, dx = \frac{a}{3} + \frac{b}{2} + c$$

So $a = -\frac{3b}{2} - 3c$ and

$$p = \left(-\frac{3b}{2} - 3c \right) x^2 + bx + c = b \left(-\frac{3}{2}x^2 + x \right) + c(-3x + 1).$$

Exercise

Let $L : P_2(\mathbb{R}) \rightarrow \mathbb{R}$ by $p \mapsto \int_0^1 p \, dx$. Find the kernel of L , a basis for it, and the dimension for the space.

Solution: We need to find all polynomials $p = ax^2 + bx + c$ so that

$$0 = \int_0^1 ax^2 + bx + c \, dx = \frac{a}{3} + \frac{b}{2} + c$$

So $a = -\frac{3b}{2} - 3c$ and

$$p = \left(-\frac{3b}{2} - 3c \right) x^2 + bx + c = b \left(-\frac{3}{2}x^2 + x \right) + c(-3x + 1).$$

So

$$\ker(L) = \text{span} \left\{ -\frac{3}{2}x^2 + x, -3x + 1 \right\}$$

Exercise

Let $L : P_2(\mathbb{R}) \rightarrow \mathbb{R}$ by $p \mapsto \int_0^1 p \, dx$. Find the kernel of L , a basis for it, and the dimension for the space.

Solution: We need to find all polynomials $p = ax^2 + bx + c$ so that

$$0 = \int_0^1 ax^2 + bx + c \, dx = \frac{a}{3} + \frac{b}{2} + c$$

So $a = -\frac{3b}{2} - 3c$ and

$$p = \left(-\frac{3b}{2} - 3c\right)x^2 + bx + c = b\left(-\frac{3}{2}x^2 + x\right) + c(-3x + 1).$$

So

$$\ker(L) = \text{span} \left\{ -\frac{3}{2}x^2 + x, -3x + 1 \right\}$$

has dimension 2.

Range Example

$\text{range}(L)$ is a subspace of W .

Range Example

$\text{range}(L)$ is a subspace of W . So we can find a basis for it.

Example: Let $L_A : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ by multiplication by

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

Find $\text{range}(L_A)$ and give a basis for it.

Range Example

$\text{range}(L)$ is a subspace of W . So we can find a basis for it.

Example: Let $L_A : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ by multiplication by

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

Find $\text{range}(L_A)$ and give a basis for it.

Solution: The range of L_A is the set of all possible products Ax with $x \in \mathbb{R}^3$.

Range Example

$\text{range}(L)$ is a subspace of W . So we can find a basis for it.

Example: Let $L_A : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ by multiplication by

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

Find $\text{range}(L_A)$ and give a basis for it.

Solution: The range of L_A is the set of all possible products Ax with $x \in \mathbb{R}^3$. That is: it is the *column space of A*! So

$$\text{range}(L_A) = \text{span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$$

Range Example

$\text{range}(L)$ is a subspace of W . So we can find a basis for it.

Example: Let $L_A : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ by multiplication by

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

Find $\text{range}(L_A)$ and give a basis for it.

Solution: The range of L_A is the set of all possible products Ax with $x \in \mathbb{R}^3$. That is: it is the *column space of A*! So

$$\text{range}(L_A) = \text{span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$$

So a basis will be those column vectors which are linearly independent:

Range Example

$\text{range}(L)$ is a subspace of W . So we can find a basis for it.

Example: Let $L_A : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ by multiplication by

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

Find $\text{range}(L_A)$ and give a basis for it.

Solution: The range of L_A is the set of all possible products Ax with $x \in \mathbb{R}^3$. That is: it is the *column space of A*! So

$$\text{range}(L_A) = \text{span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$$

So a basis will be those column vectors which are linearly independent:

$$\text{range}(L_A) = \text{span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}.$$

Range Example

$\text{range}(L)$ is a subspace of W . So we can find a basis for it.

Example: Let $L_A : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ by multiplication by

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

Find $\text{range}(L_A)$ and give a basis for it.

Solution: The range of L_A is the set of all possible products Ax with $x \in \mathbb{R}^3$. That is: it is the *column space of A*! So

$$\text{range}(L_A) = \text{span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$$

So a basis will be those column vectors which are linearly independent:

$$\text{range}(L_A) = \text{span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}.$$

It is two dimensional.

Range Characterization

If L_A is a linear operator defined by matrix multiplication by A then

$$\text{range}(L_A) = \text{col}(A).$$

Range Characterization

If L_A is a linear operator defined by matrix multiplication by A then

$$\text{range}(L_A) = \text{col}(A).$$

Hence, a basis for the range of L_A is the set of linearly independent columns of A !

Range Characterization

If L_A is a linear operator defined by matrix multiplication by A then

$$\text{range}(L_A) = \text{col}(A).$$

Hence, a basis for the range of L_A is the set of linearly independent columns of A !

Again, this is a problem that we have already solved!

Range, Kernel, Invertibility

A linear map $T : V \rightarrow W$ is invertible if and only if it is

Definition

- ▶ **one to one:** If $T(v_1) = T(v_2)$ then $v_1 = v_2$.

Range, Kernel, Invertibility

A linear map $T : V \rightarrow W$ is invertible if and only if it is

Definition

- ▶ **one to one:** If $T(v_1) = T(v_2)$ then $v_1 = v_2$.
- ▶ **onto:** For every element $w \in W$ there is a $v \in V$ so that $T(v) = w$.

Range, Kernel, Invertibility

A linear map $T : V \rightarrow W$ is invertible if and only if it is

Definition

- ▶ **one to one:** If $T(v_1) = T(v_2)$ then $v_1 = v_2$.
- ▶ **onto:** For every element $w \in W$ there is a $v \in V$ so that $T(v) = w$.

One to one and onto can be interpreted in terms of the subspaces discussed:

Theorem

- ▶ T is onto $\Leftrightarrow \text{range}(T) = W$.

Range, Kernel, Invertibility

A linear map $T : V \rightarrow W$ is invertible if and only if it is

Definition

- ▶ **one to one:** If $T(v_1) = T(v_2)$ then $v_1 = v_2$.
- ▶ **onto:** For every element $w \in W$ there is a $v \in V$ so that $T(v) = w$.

One to one and onto can be interpreted in terms of the subspaces discussed:

Theorem

- ▶ T is onto $\Leftrightarrow \text{range}(T) = W$.
- ▶ T is one to one $\Leftrightarrow \ker(T) = \{0\}$.

Proof

- ▶ We won't prove the first statement since it is just a rephrasing of definitions.

Proof

- ▶ We won't prove the first statement since it is just a rephrasing of definitions.
- ▶ Let T be one to one and $v \in \ker(T)$.

Proof

- ▶ We won't prove the first statement since it is just a rephrasing of definitions.
- ▶ Let T be one to one and $v \in \ker(T)$. Then

$$T(v) = 0 = T(0)$$

Proof

- ▶ We won't prove the first statement since it is just a rephrasing of definitions.
- ▶ Let T be one to one and $v \in \ker(T)$. Then

$$T(v) = 0 = T(0) \Rightarrow v = 0.$$

Proof

- ▶ We won't prove the first statement since it is just a rephrasing of definitions.
- ▶ Let T be one to one and $v \in \ker(T)$. Then

$$T(v) = 0 = T(0) \Rightarrow v = 0.$$

So $\ker(T) = \{0\}$.

Proof

- ▶ We won't prove the first statement since it is just a rephrasing of definitions.
- ▶ Let T be one to one and $v \in \ker(T)$. Then

$$T(v) = 0 = T(0) \Rightarrow v = 0.$$

So $\ker(T) = \{0\}$.

- ▶ Next, suppose $\ker(T) = \{0\}$

Proof

- ▶ We won't prove the first statement since it is just a rephrasing of definitions.
- ▶ Let T be one to one and $v \in \ker(T)$. Then

$$T(v) = 0 = T(0) \Rightarrow v = 0.$$

So $\ker(T) = \{0\}$.

- ▶ Next, suppose $\ker(T) = \{0\}$ and $T(v_1) = T(v_2)$.

Proof

- ▶ We won't prove the first statement since it is just a rephrasing of definitions.
- ▶ Let T be one to one and $v \in \ker(T)$. Then

$$T(v) = 0 = T(0) \Rightarrow v = 0.$$

So $\ker(T) = \{0\}$.

- ▶ Next, suppose $\ker(T) = \{0\}$ and $T(v_1) = T(v_2)$. Then:

$$T(v_1 - v_2) = T(v_1) - T(v_2) = 0.$$

Proof

- ▶ We won't prove the first statement since it is just a rephrasing of definitions.
- ▶ Let T be one to one and $v \in \ker(T)$. Then

$$T(v) = 0 = T(0) \Rightarrow v = 0.$$

So $\ker(T) = \{0\}$.

- ▶ Next, suppose $\ker(T) = \{0\}$ and $T(v_1) = T(v_2)$. Then:

$$T(v_1 - v_2) = T(v_1) - T(v_2) = 0.$$

So $v_1 - v_2 \in \ker(T)$

Proof

- ▶ We won't prove the first statement since it is just a rephrasing of definitions.
- ▶ Let T be one to one and $v \in \ker(T)$. Then

$$T(v) = 0 = T(0) \Rightarrow v = 0.$$

So $\ker(T) = \{0\}$.

- ▶ Next, suppose $\ker(T) = \{0\}$ and $T(v_1) = T(v_2)$. Then:

$$T(v_1 - v_2) = T(v_1) - T(v_2) = 0.$$

So $v_1 - v_2 \in \ker(T) \Rightarrow v_1 - v_2 = 0$.

Proof

- ▶ We won't prove the first statement since it is just a rephrasing of definitions.
- ▶ Let T be one to one and $v \in \ker(T)$. Then

$$T(v) = 0 = T(0) \Rightarrow v = 0.$$

So $\ker(T) = \{0\}$.

- ▶ Next, suppose $\ker(T) = \{0\}$ and $T(v_1) = T(v_2)$. Then:

$$T(v_1 - v_2) = T(v_1) - T(v_2) = 0.$$

So $v_1 - v_2 \in \ker(T) \Rightarrow v_1 - v_2 = 0$. Therefore $v_1 = v_2$ and T is one to one.

The Dimension Theorem Again

Theorem

Let V be a finite dimensional vector space and $T : V \rightarrow W$ a linear map. Then

$$\dim(V) = \text{rank}(T) + \text{nullity}(T)$$

where

$$\text{rank}(T) = \dim \text{range } T \quad \text{and} \quad \text{nullity}(T) = \dim \ker T$$