Exercise

Is the vector

$$v = \begin{bmatrix} 6 \\ 6 \\ -8 \end{bmatrix}$$

a linear combination of the vectors

$$\begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix} \quad \begin{bmatrix} 2 \\ 1 \\ -2 \end{bmatrix} \quad \begin{bmatrix} -3 \\ -2 \\ 0 \end{bmatrix}?$$

Exercise

Is the vector

$$v = \begin{bmatrix} 6 \\ 6 \\ -8 \end{bmatrix}$$

a linear combination of the vectors

$$\begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix} \quad \begin{bmatrix} 2 \\ 1 \\ -2 \end{bmatrix} \quad \begin{bmatrix} -3 \\ -2 \\ 0 \end{bmatrix}?$$

Solution: Try to solve

$$a \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix} + b \begin{bmatrix} 2 \\ 1 \\ -2 \end{bmatrix} + c \begin{bmatrix} -3 \\ -2 \\ 0 \end{bmatrix} = \begin{bmatrix} 6 \\ 6 \\ -8 \end{bmatrix}$$

This becomes

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 1 & -2 \\ -3 & -2 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 6 \\ 6 \\ -8 \end{bmatrix}.$$

This becomes

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 1 & -2 \\ -3 & -2 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 6 \\ 6 \\ -8 \end{bmatrix}.$$

Create:

$$\left[\begin{array}{ccc|ccc|c}
1 & 2 & -3 & 6 \\
2 & 1 & -2 & 6 \\
-3 & -2 & 0 & -8
\end{array}\right]$$

This becomes

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 1 & -2 \\ -3 & -2 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 6 \\ 6 \\ -8 \end{bmatrix}.$$

Create:

$$\left[\begin{array}{ccc|ccc|c}
1 & 2 & -3 & 6 \\
2 & 1 & -2 & 6 \\
-3 & -2 & 0 & -8
\end{array}\right]$$

Use row reduction to find a = 2, b = 1, c = -1.

This becomes

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 1 & -2 \\ -3 & -2 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 6 \\ 6 \\ -8 \end{bmatrix}.$$

Create:

$$\left[\begin{array}{ccc|ccc}
1 & 2 & -3 & 6 \\
2 & 1 & -2 & 6 \\
-3 & -2 & 0 & -8
\end{array}\right]$$

Use row reduction to find a=2, b=1, c=-1. So v is a linear combination of the others.

(2) Examine only the system matrix:

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 1 & -2 \\ -3 & -2 & 0 \end{bmatrix}$$

(2) Examine only the system matrix:

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 1 & -2 \\ -3 & -2 & 0 \end{bmatrix}$$

Row reduce it to an upper triangular form with no zeros on the diagonal.

(2) Examine only the system matrix:

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 1 & -2 \\ -3 & -2 & 0 \end{bmatrix}$$

Row reduce it to an upper triangular form with no zeros on the diagonal. Then it is invertible.

(2) Examine only the system matrix:

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 1 & -2 \\ -3 & -2 & 0 \end{bmatrix}$$

Row reduce it to an upper triangular form with no zeros on the diagonal. Then it is invertible. Hence values of a, b, and c must exist.

(2) Examine only the system matrix:

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 1 & -2 \\ -3 & -2 & 0 \end{bmatrix}$$

Row reduce it to an upper triangular form with no zeros on the diagonal. Then it is invertible. Hence values of a, b, and c must exist. v is a linear combination of the others.

(3) Compute

$$\begin{vmatrix} 1 & 2 & -3 \\ 2 & 1 & -2 \\ -3 & -2 & 0 \end{vmatrix} = 6.$$

(2) Examine only the system matrix:

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 1 & -2 \\ -3 & -2 & 0 \end{bmatrix}$$

Row reduce it to an upper triangular form with no zeros on the diagonal. Then it is invertible. Hence values of a, b, and c must exist. v is a linear combination of the others.

(3) Compute

$$\begin{vmatrix} 1 & 2 & -3 \\ 2 & 1 & -2 \\ -3 & -2 & 0 \end{vmatrix} = 6.$$

 $\det \neq 0$ so A is non-singular.

(2) Examine only the system matrix:

$$\begin{bmatrix} 1 & 2 & -3 \\ 2 & 1 & -2 \\ -3 & -2 & 0 \end{bmatrix}$$

Row reduce it to an upper triangular form with no zeros on the diagonal. Then it is invertible. Hence values of a, b, and c must exist. v is a linear combination of the others.

(3) Compute

$$\begin{vmatrix} 1 & 2 & -3 \\ 2 & 1 & -2 \\ -3 & -2 & 0 \end{vmatrix} = 6.$$

 $\det \neq 0$ so A is non-singular. So values of a, b, and c must exist and v is a linear combination of the others.

Span

Recall from last class the following

Definition

Let V be a vector space and $W \subseteq V$. Then the **span** of W is the set of all (finite) linear combinations of elements of W:

$$\mathsf{span}(W) := \left\{ \sum_{i=1}^n \mathsf{a}_i \mathsf{v}_i : \mathsf{a}_i \in \mathsf{F}, \mathsf{v}_i \in W, \mathsf{n} \in \mathbb{N} \right\}$$

Span

Recall from last class the following

Definition

Let V be a vector space and $W \subseteq V$. Then the **span** of W is the set of all (finite) linear combinations of elements of W:

$$\mathsf{span}(W) := \left\{ \sum_{i=1}^n a_i v_i : a_i \in F, v_i \in W, n \in \mathbb{N} \right\}$$

Some comments:

ightharpoonup We already showed before that span(W) is a subspace of V.

Span

Recall from last class the following

Definition

Let V be a vector space and $W \subseteq V$. Then the **span** of W is the set of all (finite) linear combinations of elements of W:

$$\mathsf{span}(W) := \left\{ \sum_{i=1}^n a_i v_i : a_i \in F, v_i \in W, n \in \mathbb{N} \right\}$$

Some comments:

- ightharpoonup We already showed before that span(W) is a subspace of V.
- ► The question "Is v a linear combination of elements of W?" can be rephrased as "Is $v \in \text{span}(W)$?"

Exercises

1. ls

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \in \mathsf{span} \left(\left\{ \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\} \right)?$$

- 2. Is x^2 in the span of $x^2 2x 1$, x + 1, 1?
- 3. Explain why every vector in \mathbb{R}^3 is in the span of

$$\begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix} \quad \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \quad \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}.$$

1. No. Row reduce

$$\left[\begin{array}{cc|c}
2 & 1 & 1 \\
0 & 1 & 0 \\
1 & 1 & 0
\end{array}\right]$$

and find there is no solution.

1. No. Row reduce

$$\left[\begin{array}{cc|c}
2 & 1 & 1 \\
0 & 1 & 0 \\
1 & 1 & 0
\end{array}\right]$$

and find there is no solution.

2. Yes.

$$x^2 = (x^2 - 2x + 1) + 2(x + 1) - 3(1).$$

1. No. Row reduce

$$\left[\begin{array}{cc|c}
2 & 1 & 1 \\
0 & 1 & 0 \\
1 & 1 & 0
\end{array}\right]$$

and find there is no solution.

2. Yes.

$$x^2 = (x^2 - 2x + 1) + 2(x + 1) - 3(1).$$

3. Note that

$$\begin{bmatrix} 2 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

is not singular.

Lingo

In the last case we have that

$$\mathbb{R}^3 = \operatorname{span}\left(\underbrace{\left\{\begin{bmatrix}2\\0\\1\end{bmatrix},\begin{bmatrix}1\\1\\1\end{bmatrix},\begin{bmatrix}0\\1\\0\end{bmatrix}\right\}}_{W}\right)$$

Lingo

In the last case we have that

$$\mathbb{R}^3 = \mathsf{span}\left(\underbrace{\left\{\begin{bmatrix}2\\0\\1\end{bmatrix},\begin{bmatrix}1\\1\\1\end{bmatrix},\begin{bmatrix}0\\1\\0\end{bmatrix}\right\}}_{W}\right)$$

We say

"
$$W$$
 spans \mathbb{R}^3 ." or " \mathbb{R}^3 is spanned by W ."