

Properties of Addition and Scalar Multiplication in \mathbb{R}^n :

Let u , v , and w be vectors in \mathbb{R}^n with scalar numbers r and s in \mathbb{R} . Then

1. $u + v = v + u$
2. $u + (v + w) = (u + v) + w$
3. $u + 0 = 0 + u = u$
4. $u + (-u) = 0$
5. $r(u + v) = ru + rv$
6. $(r + s)u = ru + su$
7. $r(su) = (rs)u$
8. $1u = u$

Question: What is more important? The operations which produce the properties? Or just the properties themselves?

Example: Matrices

Consider the set of matrices $\mathbb{R}^{n \times m}$ with matrix addition and scalar multiplication.

Example: Matrices

Consider the set of matrices $\mathbb{R}^{n \times m}$ with matrix addition and scalar multiplication. Then for matrices A , B , and C ; and real numbers r and s we have:

1. $A + B = B + A$
2. $A + (B + C) = (A + B) + C$
3. $A + 0 = 0 + A = A$ where 0 is the zero matrix.
4. $A + (-A) = 0$
5. $r(A + B) = rA + rB$
6. $(r + s)A = rA + sA$
7. $r(sA) = (rs)A$
8. $1A = A$

Example: Matrices

Consider the set of matrices $\mathbb{R}^{n \times m}$ with matrix addition and scalar multiplication. Then for matrices A , B , and C ; and real numbers r and s we have:

1. $A + B = B + A$
2. $A + (B + C) = (A + B) + C$
3. $A + 0 = 0 + A = A$ where 0 is the zero matrix.
4. $A + (-A) = 0$
5. $r(A + B) = rA + rB$
6. $(r + s)A = rA + sA$
7. $r(sA) = (rs)A$
8. $1A = A$

This is the same collection of properties with different elements (matrices) and operations (matrix addition).

Example: Functions

Consider the set $F(D, \mathbb{R})$ which is the set of all functions with domain D and range \mathbb{R} .

Example: Functions

Consider the set $F(D, \mathbb{R})$ which is the set of all functions with domain D and range \mathbb{R} . Then for functions f and g and real number r we can define

$$(f + g)(x) = f(x) + g(x) \quad \text{and} \quad (rf)(x) = r(f(x)).$$

Example: Functions

Consider the set $F(D, \mathbb{R})$ which is the set of all functions with domain D and range \mathbb{R} . Then for functions f and g and real number r we can define

$$(f + g)(x) = f(x) + g(x) \quad \text{and} \quad (rf)(x) = r(f(x)).$$

For functions f , g , and h ; and real numbers r and s we have:

1. $f + g = g + f$
2. $f + (g + h) = (f + g) + h$
3. $f + e = e + f = f$ where e is the function $f(x) = 0$.
4. $f + (-f) = e$
5. $r(f + g) = rf + rg$
6. $(r + s)f = rf + sf$
7. $r(sf) = (rs)f$
8. $1f = f$

Example: Functions

Consider the set $F(D, \mathbb{R})$ which is the set of all functions with domain D and range \mathbb{R} . Then for functions f and g and real number r we can define

$$(f + g)(x) = f(x) + g(x) \quad \text{and} \quad (rf)(x) = r(f(x)).$$

For functions f , g , and h ; and real numbers r and s we have:

1. $f + g = g + f$
2. $f + (g + h) = (f + g) + h$
3. $f + e = e + f = f$ where e is the function $f(x) = 0$.
4. $f + (-f) = e$
5. $r(f + g) = rf + rg$
6. $(r + s)f = rf + sf$
7. $r(sf) = (rs)f$
8. $1f = f$

This is the same collection of properties with different elements (functions) and operations (function addition).

Moral: These are all examples of the same thing

We have several examples of sets and operations which share the same properties. So let's classify them as examples of one type of mathematical object:

Moral: These are all examples of the same thing

We have several examples of sets and operations which share the same properties. So let's classify them as examples of one type of mathematical object:

Definition

Let F be a field (i.e., the real numbers). A **vector space** over F is a set V (the vectors) and operations

$\oplus : V \times V \rightarrow V$ (vector addition) $\odot : F \times V \rightarrow V$ scalar multiplication

Moral: These are all examples of the same thing

We have several examples of sets and operations which share the same properties. So let's classify them as examples of one type of mathematical object:

Definition

Let F be a field (i.e., the real numbers). A **vector space** over F is a set V (the vectors) and operations

$\oplus : V \times V \rightarrow V$ (vector addition) $\odot : F \times V \rightarrow V$ scalar multiplication

with the following properties:

VS Properties

For all vectors u , v , and w in V and scalar values r and s in F we have

1. $u \oplus v = v \oplus u$
2. $u \oplus (v \oplus w) = (u \oplus v) \oplus w$
3. There is a vector $e \in V$ so that $u \oplus e = u$
4. There is a vector u^{-1} so that $u \oplus u^{-1} = e$.
5. $r \odot (u \oplus v) = (r \odot u) \oplus (r \odot v)$
6. $(r + s) \odot u = (r \odot u) \oplus (s \odot u)$
7. $r \odot (s \odot u) = (rs) \odot u$
8. $1 \odot u = u$

- ▶ Since the range of \oplus and \odot are both V , we say V is *closed* under the operations.

Comments

- ▶ Since the range of \oplus and \odot are both V , we say V is *closed* under the operations.
- ▶ Typically we do not use the \odot and \oplus notation:

$$u \oplus v \leftrightarrow u + v \quad \text{and} \quad r \odot v \leftrightarrow rv.$$

Comments

- ▶ Since the range of \oplus and \odot are both V , we say V is *closed* under the operations.
- ▶ Typically we do not use the \odot and \oplus notation:

$$u \oplus v \leftrightarrow u + v \quad \text{and} \quad r \odot v \leftrightarrow rv.$$

It is meant to emphasize that addition and multiplication are not “regular.”

Comments

- ▶ Since the range of \oplus and \odot are both V , we say V is *closed* under the operations.
- ▶ Typically we do not use the \odot and \oplus notation:

$$u \oplus v \leftrightarrow u + v \quad \text{and} \quad r \odot v \leftrightarrow rv.$$

It is meant to emphasize that addition and multiplication are not “regular.”

- ▶ e is called the *identity* and u^{-1} is the *inverse* of u .

Not everything is a vector space

Not everything is a vector space

For example, consider \mathbb{R}^n with

$$u \oplus v := u - v.$$

Not everything is a vector space

For example, consider \mathbb{R}^n with

$$u \oplus v := u - v.$$

This operation is not commutative, nor associative, and there are issues with distributivity.

Not everything is a vector space

For example, consider \mathbb{R}^n with

$$u \oplus v := u - v.$$

This operation is not commutative, nor associative, and there are issues with distributivity. (But it does have (one sided) identity and inverses!)

Not everything is a vector space

For example, consider \mathbb{R}^n with

$$u \oplus v := u - v.$$

This operation is not commutative, nor associative, and there are issues with distributivity. (But it does have (one sided) identity and inverses!)

Second example: Consider polynomials of even degree with function addition and scalar multiplication.

Not everything is a vector space

For example, consider \mathbb{R}^n with

$$u \oplus v := u - v.$$

This operation is not commutative, nor associative, and there are issues with distributivity. (But it does have (one sided) identity and inverses!)

Second example: Consider polynomials of even degree with function addition and scalar multiplication. Then:

$$x^2 + x + 1 \oplus -x^2 + x - 2 = 2x + 3$$

Not everything is a vector space

For example, consider \mathbb{R}^n with

$$u \oplus v := u - v.$$

This operation is not commutative, nor associative, and there are issues with distributivity. (But it does have (one sided) identity and inverses!)

Second example: Consider polynomials of even degree with function addition and scalar multiplication. Then:

$$x^2 + x + 1 \oplus -x^2 + x - 2 = 2x + 3$$

But $2x + 3$ is of odd degree and so the set is not *closed* under addition.

What about other properties?

What about other properties?

In all of our examples of vector spaces, there are other things which can be shown to be true. For example, in \mathbb{R}^n we have

$$0u = 0.$$

What about other properties?

In all of our examples of vector spaces, there are other things which can be shown to be true. For example, in \mathbb{R}^n we have

$$0u = 0.$$

This is true even if we replace u with a function, or a matrix, etc.

What about other properties?

In all of our examples of vector spaces, there are other things which can be shown to be true. For example, in \mathbb{R}^n we have

$$0u = 0.$$

This is true even if we replace u with a function, or a matrix, etc. Why not include this property in our list?

The collection of properties 1-8 form some sort of *minimal* collection of assumptions. All others can (should) be derived from them!

Elementary Derived Properties

Theorem

Suppose V is a vector space. Let $v \in V$ and $c \in F$. Then

- 1. The identity is unique.*

Elementary Derived Properties

Theorem

Suppose V is a vector space. Let $v \in V$ and $c \in F$. Then

- 1. The identity is unique.*
- 2. v has a unique inverse.*

Elementary Derived Properties

Theorem

Suppose V is a vector space. Let $v \in V$ and $c \in F$. Then

- 1. The identity is unique.*
- 2. v has a unique inverse.*
- 3. $0 \odot v = e$.*

Elementary Derived Properties

Theorem

Suppose V is a vector space. Let $v \in V$ and $c \in F$. Then

- 1. The identity is unique.*
- 2. v has a unique inverse.*
- 3. $0 \odot v = e$.*
- 4. $c \odot e = e$.*

Elementary Derived Properties

Theorem

Suppose V is a vector space. Let $v \in V$ and $c \in F$. Then

- 1. The identity is unique.*
- 2. v has a unique inverse.*
- 3. $0 \odot v = e$.*
- 4. $c \odot e = e$.*
- 5. If $c \odot v = e$ then either $c = 0$ or $v = e$.*

Elementary Derived Properties

Theorem

Suppose V is a vector space. Let $v \in V$ and $c \in F$. Then

- 1. The identity is unique.*
- 2. v has a unique inverse.*
- 3. $0 \odot v = e$.*
- 4. $c \odot e = e$.*
- 5. If $c \odot v = e$ then either $c = 0$ or $v = e$.*
- 6. $(-1) \odot v = v^{-1}$.*

Elementary Derived Properties

Theorem

Suppose V is a vector space. Let $v \in V$ and $c \in F$. Then

- 1. The identity is unique.*
- 2. v has a unique inverse.*
- 3. $0 \odot v = e$.*
- 4. $c \odot e = e$.*
- 5. If $c \odot v = e$ then either $c = 0$ or $v = e$.*
- 6. $(-1) \odot v = v^{-1}$.*

Some proofs

We won't prove all of these, but here are some arguments:

- 1 The identity is unique.

Some proofs

We won't prove all of these, but here are some arguments:

- 1 The identity is unique.

Proof:

Some proofs

We won't prove all of these, but here are some arguments:

- 1 The identity is unique.

Proof: Suppose on the contrary that there are two identity elements in V .

Some proofs

We won't prove all of these, but here are some arguments:

- 1 The identity is unique.

Proof: Suppose on the contrary that there are two identity elements in V . Call them e_1 and e_2 .

Some proofs

We won't prove all of these, but here are some arguments:

- 1 The identity is unique.

Proof: Suppose on the contrary that there are two identity elements in V . Call them e_1 and e_2 . Then:

$$e_1$$

Some proofs

We won't prove all of these, but here are some arguments:

- 1 The identity is unique.

Proof: Suppose on the contrary that there are two identity elements in V . Call them e_1 and e_2 . Then:

$$e_1 = e_1 \oplus e_2$$

Some proofs

We won't prove all of these, but here are some arguments:

- 1 The identity is unique.

Proof: Suppose on the contrary that there are two identity elements in V . Call them e_1 and e_2 . Then:

$$e_1 = e_1 \oplus e_2 = e_2.$$

Some proofs

We won't prove all of these, but here are some arguments:

- 1 The identity is unique.

Proof: Suppose on the contrary that there are two identity elements in V . Call them e_1 and e_2 . Then:

$$e_1 = e_1 \oplus e_2 = e_2.$$

So $e_1 = e_2$ and the identity is unique. Now that we know it is unique, we often write 0 for the identity.

$$3 \quad 0 \odot v = e.$$

3 $0 \odot v = e.$

Proof:

3 $0 \odot v = e.$

Proof: In order to prove this, we need to show that $w \oplus (0 \odot v) = w$ for any vector w .

3 $0 \odot v = e$.

Proof: In order to prove this, we need to show that $w \oplus (0 \odot v) = w$ for any vector w . Then $0 \oplus v = e$ from the uniqueness of e .

3 $0 \odot v = e.$

Proof: In order to prove this, we need to show that $w \oplus (0 \odot v) = w$ for any vector w . Then $0 \oplus v = e$ from the uniqueness of e . To begin:

$$v = 1 \odot v$$

3 $0 \odot v = e.$

Proof: In order to prove this, we need to show that $w \oplus (0 \odot v) = w$ for any vector w . Then $0 \oplus v = e$ from the uniqueness of e . To begin:

$$v = 1 \odot v = (1 + 0) \odot v$$

3 $0 \odot v = e$.

Proof: In order to prove this, we need to show that $w \oplus (0 \odot v) = w$ for any vector w . Then $0 \oplus v = e$ from the uniqueness of e . To begin:

$$v = 1 \odot v = (1 + 0) \odot v = (1 \odot v) \oplus (0 \odot v)$$

3 $0 \odot v = e$.

Proof: In order to prove this, we need to show that $w \oplus (0 \odot v) = w$ for any vector w . Then $0 \oplus v = e$ from the uniqueness of e . To begin:

$$v = 1 \odot v = (1 + 0) \odot v = (1 \odot v) \oplus (0 \odot v) = v \oplus (0 \odot v).$$

3 $0 \odot v = e$.

Proof: In order to prove this, we need to show that $w \oplus (0 \odot v) = w$ for any vector w . Then $0 \oplus v = e$ from the uniqueness of e . To begin:

$$v = 1 \odot v = (1 + 0) \odot v = (1 \odot v) \oplus (0 \odot v) = v \oplus (0 \odot v).$$

This shows $v = v \oplus (0 \odot v)$ for any v .

3 $0 \odot v = e$.

Proof: In order to prove this, we need to show that $w \oplus (0 \odot v) = w$ for any vector w . Then $0 \oplus v = e$ from the uniqueness of e . To begin:

$$v = 1 \odot v = (1 + 0) \odot v = (1 \odot v) \oplus (0 \odot v) = v \oplus (0 \odot v).$$

This shows $v = v \oplus (0 \odot v)$ for any v .

So for any w we can write

$$w = w \oplus v^{-1} \oplus v$$

3 $0 \odot v = e$.

Proof: In order to prove this, we need to show that $w \oplus (0 \odot v) = w$ for any vector w . Then $0 \oplus v = e$ from the uniqueness of e . To begin:

$$v = 1 \odot v = (1 + 0) \odot v = (1 \odot v) \oplus (0 \odot v) = v \oplus (0 \odot v).$$

This shows $v = v \oplus (0 \odot v)$ for any v .

So for any w we can write

$$w = w \oplus v^{-1} \oplus v = w \oplus v^{-1} \oplus v \oplus (0 \odot v)$$

3 $0 \odot v = e$.

Proof: In order to prove this, we need to show that $w \oplus (0 \odot v) = w$ for any vector w . Then $0 \oplus v = e$ from the uniqueness of e . To begin:

$$v = 1 \odot v = (1 + 0) \odot v = (1 \odot v) \oplus (0 \odot v) = v \oplus (0 \odot v).$$

This shows $v = v \oplus (0 \odot v)$ for any v .

So for any w we can write

$$w = w \oplus v^{-1} \oplus v = w \oplus v^{-1} \oplus v \oplus (0 \odot v) = w \oplus (0 \odot v).$$

An Interesting Example

Let $A \in \mathbb{R}^{n \times m}$. Consider the set

$$H = \{x \in \mathbb{R}^m : Ax = 0\}.$$

An Interesting Example

Let $A \in \mathbb{R}^{n \times m}$. Consider the set

$$H = \{x \in \mathbb{R}^m : Ax = 0\}.$$

This is the set of all solutions to the homogeneous problem.

An Interesting Example

Let $A \in \mathbb{R}^{n \times m}$. Consider the set

$$H = \{x \in \mathbb{R}^m : Ax = 0\}.$$

This is the set of all solutions to the homogeneous problem.

Since $H \subset \mathbb{R}^m$ we have addition and scalar multiplication defined for it.

An Interesting Example

Let $A \in \mathbb{R}^{n \times m}$. Consider the set

$$H = \{x \in \mathbb{R}^m : Ax = 0\}.$$

This is the set of all solutions to the homogeneous problem.

Since $H \subset \mathbb{R}^m$ we have addition and scalar multiplication defined for it.

This means that all of the algebraic properties 1-8 hold for H assuming it is closed!

Set of Homogeneous Solutions is a Vector Space!

Set of Homogeneous Solutions is a Vector Space!

So:

If $x, y \in H$ then $A(x + y) = Ax + Ay = 0 + 0 = 0$.

Set of Homogeneous Solutions is a Vector Space!

So:

If $x, y \in H$ then $A(x + y) = Ax + Ay = 0 + 0 = 0$. Therefore $x + y \in H$ and H is closed under addition.

Set of Homogeneous Solutions is a Vector Space!

So:

If $x, y \in H$ then $A(x + y) = Ax + Ay = 0 + 0 = 0$. Therefore $x + y \in H$ and H is closed under addition.

If $x \in H$ and $r \in \mathbb{R}$ then $A(rx) = rAx = r0 = 0$.

Set of Homogeneous Solutions is a Vector Space!

So:

If $x, y \in H$ then $A(x + y) = Ax + Ay = 0 + 0 = 0$. Therefore $x + y \in H$ and H is closed under addition.

If $x \in H$ and $r \in \mathbb{R}$ then $A(rx) = rAx = r0 = 0$. Therefore $rx \in H$ and H is closed under scalar multiplication.

Set of Homogeneous Solutions is a Vector Space!

So:

If $x, y \in H$ then $A(x + y) = Ax + Ay = 0 + 0 = 0$. Therefore $x + y \in H$ and H is closed under addition.

If $x \in H$ and $r \in \mathbb{R}$ then $A(rx) = rAx = r0 = 0$. Therefore $rx \in H$ and H is closed under scalar multiplication.

In conclusion: H is a vector space.

Set of Homogeneous Solutions is a Vector Space!

So:

If $x, y \in H$ then $A(x + y) = Ax + Ay = 0 + 0 = 0$. Therefore $x + y \in H$ and H is closed under addition.

If $x \in H$ and $r \in \mathbb{R}$ then $A(rx) = rAx = r0 = 0$. Therefore $rx \in H$ and H is closed under scalar multiplication.

In conclusion: H is a vector space. It is also a subset of another vector space \mathbb{R}^m .

Set of Homogeneous Solutions is a Vector Space!

So:

If $x, y \in H$ then $A(x + y) = Ax + Ay = 0 + 0 = 0$. Therefore $x + y \in H$ and H is closed under addition.

If $x \in H$ and $r \in \mathbb{R}$ then $A(rx) = rAx = r0 = 0$. Therefore $rx \in H$ and H is closed under scalar multiplication.

In conclusion: H is a vector space. It is also a subset of another vector space \mathbb{R}^m . Such an object is called a **subspace**.