

Theoretical Result

Recall that a basis is a set which is both spanning and linearly independent.

Theoretical Result

Recall that a basis is a set which is both spanning and linearly independent.

In the last class we proved the following result:

Theoretical Result

Recall that a basis is a set which is both spanning and linearly independent.

In the last class we proved the following result:

Theorem

Let S be a nonempty finite subset of a vector space V . Then there is a set β in S with β a basis for $\text{span}(S)$.

Theoretical Result

Recall that a basis is a set which is both spanning and linearly independent.

In the last class we proved the following result:

Theorem

Let S be a nonempty finite subset of a vector space V . Then there is a set β in S with β a basis for $\text{span}(S)$.

- ▶ Idea: We can remove vectors from a finite set until we get to a linearly independent case WITHOUT losing the spanning property!

Theoretical Result

Recall that a basis is a set which is both spanning and linearly independent.

In the last class we proved the following result:

Theorem

Let S be a nonempty finite subset of a vector space V . Then there is a set β in S with β a basis for $\text{span}(S)$.

- ▶ Idea: We can remove vectors from a finite set until we get to a linearly independent case WITHOUT losing the spanning property!
- ▶ The proof hinged on the fact that such a basis is the *LARGEST* linearly independent subset of S .

An example

Find a basis for $\text{span}(S)$ where

$$S = \left\{ \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ -2 \end{bmatrix} \right\}.$$

An example

Find a basis for $\text{span}(S)$ where

$$S = \left\{ \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ -2 \end{bmatrix} \right\}.$$

We need to find the largest LI subset.

An example

Find a basis for $\text{span}(S)$ where

$$S = \left\{ \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ -2 \end{bmatrix} \right\}.$$

We need to find the largest LI subset. To find it, consider the homogeneous problem.

$$\begin{bmatrix} 1 & 0 & 1 & 1 & -1 \\ 0 & 1 & 1 & 2 & 1 \\ 1 & 1 & 2 & 1 & -2 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \\ e \end{bmatrix} = 0$$

and use row reduction.

Via row reduction

$$\begin{bmatrix} 1 & 0 & 1 & 1 & -1 \\ 0 & 1 & 1 & 2 & 1 \\ 1 & 1 & 2 & 1 & -2 \end{bmatrix}$$

Via row reduction

$$\begin{bmatrix} 1 & 0 & 1 & 1 & -1 \\ 0 & 1 & 1 & 2 & 1 \\ 1 & 1 & 2 & 1 & -2 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 1 & 0 & -2 \\ 0 & 1 & 1 & 0 & -1 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}.$$

Via row reduction

$$\begin{bmatrix} 1 & 0 & 1 & 1 & -1 \\ 0 & 1 & 1 & 2 & 1 \\ 1 & 1 & 2 & 1 & -2 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 1 & 0 & -2 \\ 0 & 1 & 1 & 0 & -1 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}.$$

Examine the columns with “leading ones”. If we *removed* the others, we would have a LI set!

Via row reduction

$$\begin{bmatrix} 1 & 0 & 1 & 1 & -1 \\ 0 & 1 & 1 & 2 & 1 \\ 1 & 1 & 2 & 1 & -2 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 1 & 0 & -2 \\ 0 & 1 & 1 & 0 & -1 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}.$$

Examine the columns with “leading ones”. If we *removed* the others, we would have a LI set! So

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \right\}$$

is the basis for $\text{span}(S)$.

Exercise

Find the a basis for the span of the following sets of vectors:

$$A = \left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ -6 \\ -3 \end{bmatrix}, \begin{bmatrix} -1 \\ -5 \\ 1 \\ 0 \end{bmatrix} \right\}$$

$$B = \{t^3 + t^2 - 2t + 1, t^2 + 1, t^3 - 2t, 2t^3 + 3t^2 - 4t + 3\}$$

Hint for B : write the coefficients of the polynomials as column vectors.

Solutions

Solutions

Set A :

$$\begin{bmatrix} 1 & 0 & 1 & 1 & -1 \\ 1 & 1 & 0 & 1 & -5 \\ 0 & 2 & 1 & -6 & 1 \\ -1 & 1 & -1 & -3 & 0 \end{bmatrix}$$

Solutions

Set A :

$$\begin{bmatrix} 1 & 0 & 1 & 1 & -1 \\ 1 & 1 & 0 & 1 & -5 \\ 0 & 2 & 1 & -6 & 1 \\ -1 & 1 & -1 & -3 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 1 & 1 & -1 \\ 0 & 1 & -1 & 0 & 4 \\ 0 & 0 & 1 & -2 & 3 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Solutions

Set A :

$$\begin{bmatrix} 1 & 0 & 1 & 1 & -1 \\ 1 & 1 & 0 & 1 & -5 \\ 0 & 2 & 1 & -6 & 1 \\ -1 & 1 & -1 & -3 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 1 & 1 & -1 \\ 0 & 1 & -1 & 0 & 4 \\ 0 & 0 & 1 & -2 & 3 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

So

$$\left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \\ -1 \end{bmatrix} \right\}$$

is the largest LI subset and basis for $\text{span}(A)$.

Solutions

Solutions

Set B : Use:

$$ax^3 + bx^2 + cx + d \leftrightarrow \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}$$

Solutions

Set B : Use:

$$ax^3 + bx^2 + cx + d \leftrightarrow \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}$$

System matrix:

$$\begin{bmatrix} 1 & 0 & 1 & 2 \\ 1 & 1 & 0 & 3 \\ -2 & 0 & -2 & -4 \\ 1 & 1 & 0 & 3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 1 & 2 \\ 0 & 1 & -1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Solutions

Set B : Use:

$$ax^3 + bx^2 + cx + d \leftrightarrow \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}$$

System matrix:

$$\begin{bmatrix} 1 & 0 & 1 & 2 \\ 1 & 1 & 0 & 3 \\ -2 & 0 & -2 & -4 \\ 1 & 1 & 0 & 3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 1 & 2 \\ 0 & 1 & -1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

So our basis is made up of the polynomials corresponding to the first and second columns:

$$\{t^3 + t^2 - 2t + 1, t^2 + 1\}.$$

Where are we now:

- ▶ A basis is both linearly independent and spanning.

Where are we now:

- ▶ A basis is both linearly independent and spanning. We should feel comfortable establishing if a given set is a basis or not by verifying these properties.

Where are we now:

- ▶ A basis is both linearly independent and spanning. We should feel comfortable establishing if a given set is a basis or not by verifying these properties.
- ▶ In general, a vector space can have lots of different bases.

Where are we now:

- ▶ A basis is both linearly independent and spanning. We should feel comfortable establishing if a given set is a basis or not by verifying these properties.
- ▶ In general, a vector space can have lots of different bases. We should feel ok with *finding* a basis for a given space.

Where are we now:

- ▶ A basis is both linearly independent and spanning. We should feel comfortable establishing if a given set is a basis or not by verifying these properties.
- ▶ In general, a vector space can have lots of different bases. We should feel ok with *finding* a basis for a given space.
- ▶ There seems to be some sort of “sweet spot” where a set of vectors becomes a basis.

Where are we now:

- ▶ A basis is both linearly independent and spanning. We should feel comfortable establishing if a given set is a basis or not by verifying these properties.
- ▶ In general, a vector space can have lots of different bases. We should feel ok with *finding* a basis for a given space.
- ▶ There seems to be some sort of “sweet spot” where a set of vectors becomes a basis. Roughly speaking:
 - ▶ Not spanning \sim Not enough vectors.
 - ▶ Not linearly independent \sim Too many vectors.

Where are we now:

- ▶ A basis is both linearly independent and spanning. We should feel comfortable establishing if a given set is a basis or not by verifying these properties.
- ▶ In general, a vector space can have lots of different bases. We should feel ok with *finding* a basis for a given space.
- ▶ There seems to be some sort of “sweet spot” where a set of vectors becomes a basis. Roughly speaking:
 - ▶ Not spanning \sim Not enough vectors.
 - ▶ Not linearly independent \sim Too many vectors.

We will now work on theory to make the “sweet spot” concrete.

Basis gives the maximum number of LI vectors

Theorem

Let V be a vector space with finite basis $\beta = \{v_1, \dots, v_n\}$.

Let $W = \{w_1, \dots, w_r\}$ be a set of linearly independent vectors.

Then $r \leq n$.

Basis gives the maximum number of LI vectors

Theorem

Let V be a vector space with finite basis $\beta = \{v_1, \dots, v_n\}$.

Let $W = \{w_1, \dots, w_r\}$ be a set of linearly independent vectors.

Then $r \leq n$.

Proof.

This is a simple proof (on the board) assuming that each vector can be represented as a column in \mathbb{R}^n . (which we may find later is actually not as huge an assumption as you might think!) Please come chat with me for a proof which does not use this assumption (it uses induction).



Dimension

An immediate corollary of the theorem is that if β and γ are two basis for a vector space V , then

$$|\beta| = |\gamma|.$$

Dimension

An immediate corollary of the theorem is that if β and γ are two basis for a vector space V , then

$$|\beta| = |\gamma|.$$

(they have the same number of vectors).

Dimension

An immediate corollary of the theorem is that if β and γ are two basis for a vector space V , then

$$|\beta| = |\gamma|.$$

(they have the same number of vectors).
Hence the *size* of a basis is unique!

Definition

The **dimension**, $\dim(V)$, of a vector space is the size of a basis for the space.

What does dimension tell us?

Corollary

Let V be a vector space.

What does dimension tell us?

Corollary

Let V be a vector space. Then

- 1. if L is a linearly independent subset of V then $|L| \leq \dim(V)$.*

What does dimension tell us?

Corollary

Let V be a vector space. Then

- 1. if L is a linearly independent subset of V then $|L| \leq \dim(V)$.*
- 2. if D is a subset of V and $|D| > \dim(V)$ then D is linearly dependent.*

What does dimension tell us?

Corollary

Let V be a vector space. Then

- 1. if L is a linearly independent subset of V then $|L| \leq \dim(V)$.*
- 2. if D is a subset of V and $|D| > \dim(V)$ then D is linearly dependent.*
- 3. if G is a spanning set for V then $|G| \geq \dim(V)$.*

What does dimension tell us?

Corollary

Let V be a vector space. Then

- 1. if L is a linearly independent subset of V then $|L| \leq \dim(V)$.*
- 2. if D is a subset of V and $|D| > \dim(V)$ then D is linearly dependent.*
- 3. if G is a spanning set for V then $|G| \geq \dim(V)$.*
- 4. if F is a subset of V and $|F| < \dim(V)$ then F does not span V .*

What does dimension tell us?

Corollary

Let V be a vector space. Then

- 1. if L is a linearly independent subset of V then $|L| \leq \dim(V)$.*
- 2. if D is a subset of V and $|D| > \dim(V)$ then D is linearly dependent.*
- 3. if G is a spanning set for V then $|G| \geq \dim(V)$.*
- 4. if F is a subset of V and $|F| < \dim(V)$ then F does not span V .*

Extension to Basis

A final theorem:

Theorem

If L_1 is a linearly independent subset of a vector space V . Suppose $|L_1| = m$ and $\dim(V) = n$. Then there exists $n - m$ vectors L_2 with $L_1 \cup L_2$ a basis for V .

This theorem tells us that we can always assume that we have a basis with some known LI vectors.