

## Exercise

Let

$$A = \begin{bmatrix} 1 & 0 \\ 2 & 1 \\ 3 & 2 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \end{bmatrix}$$

Compute the following:

$$A - 2B^T.$$

## Exercise

Let

$$A = \begin{bmatrix} 1 & 0 \\ 2 & 1 \\ 3 & 2 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \end{bmatrix}$$

Compute the following:

$$A - 2B^T.$$

Result:

$$A - 2B^T = \begin{bmatrix} -1 & -4 \\ -2 & -1 \\ -3 & -6 \end{bmatrix}.$$

# Matrix Operations We Know:

# Matrix Operations We Know:

- ▶ Addition.

# Matrix Operations We Know:

- ▶ Addition.
- ▶ Scalar multiplication.

# Matrix Operations We Know:

- ▶ Addition.
- ▶ Scalar multiplication.
- ▶ Transposition.

# Matrix Operations We Know:

- ▶ Addition.
- ▶ Scalar multiplication.
- ▶ Transposition.

All of these are rather “obvious” operations.

# Matrix Operations We Know:

- ▶ Addition.
- ▶ Scalar multiplication.
- ▶ Transposition.

All of these are rather “obvious” operations. Today we will talk about a non-obvious operation:



# Matrix Operations We Know:

- ▶ Addition.
- ▶ Scalar multiplication.
- ▶ Transposition.

All of these are rather “obvious” operations. Today we will talk about a non-obvious operation:

Q: “Can we multiply matrices?”

# Matrix Operations We Know:

- ▶ Addition.
- ▶ Scalar multiplication.
- ▶ Transposition.

All of these are rather “obvious” operations. Today we will talk about a non-obvious operation:

Q: “Can we multiply matrices?”

A: Yes, but the definition we use is not obvious...

# Dot Product

The standard inner product on  $\mathbb{R}^n$  is called the **dot product**:

# Dot Product

The standard inner product on  $\mathbb{R}^n$  is called the **dot product**:

## Definition

Let

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{and} \quad \mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}.$$

The **dot product** of  $\mathbf{x}$  and  $\mathbf{y}$  is defined to be

$$\mathbf{x} \cdot \mathbf{y} := x_1 y_1 + x_2 y_2 + \cdots x_n y_n$$

# Dot Product

The standard inner product on  $\mathbb{R}^n$  is called the **dot product**:

## Definition

Let

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{and} \quad \mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}.$$

The **dot product** of  $\mathbf{x}$  and  $\mathbf{y}$  is defined to be

$$\mathbf{x} \cdot \mathbf{y} := x_1 y_1 + x_2 y_2 + \cdots x_n y_n = \sum_{i=1}^n x_i y_i.$$

## Examples/Exercises

Let

$$\mathbf{w} = \begin{bmatrix} .15 \\ .15 \\ .2 \\ .2 \\ .3 \end{bmatrix} \quad \text{and} \quad \mathbf{g} = \begin{bmatrix} 90 \\ 85 \\ 80 \\ 75 \\ 90 \end{bmatrix}$$

## Examples/Exercises

Let

$$\mathbf{w} = \begin{bmatrix} .15 \\ .15 \\ .2 \\ .2 \\ .3 \end{bmatrix} \quad \text{and} \quad \mathbf{g} = \begin{bmatrix} 90 \\ 85 \\ 80 \\ 75 \\ 90 \end{bmatrix}$$

Compute  $\mathbf{w} \cdot \mathbf{g}$ :

## Examples/Exercises

Let

$$\mathbf{w} = \begin{bmatrix} .15 \\ .15 \\ .2 \\ .2 \\ .3 \end{bmatrix} \quad \text{and} \quad \mathbf{g} = \begin{bmatrix} 90 \\ 85 \\ 80 \\ 75 \\ 90 \end{bmatrix}$$

Compute  $\mathbf{w} \cdot \mathbf{g}$ :

Answer:

$$.15 * 90 + .15 * 85 + .2 * 80 + .2 * 75 + .3 * 90$$



## Examples/Exercises

Let

$$\mathbf{w} = \begin{bmatrix} .15 \\ .15 \\ .2 \\ .2 \\ .3 \end{bmatrix} \quad \text{and} \quad \mathbf{g} = \begin{bmatrix} 90 \\ 85 \\ 80 \\ 75 \\ 90 \end{bmatrix}$$

Compute  $\mathbf{w} \cdot \mathbf{g}$ :

Answer:

$$.15 * 90 + .15 * 85 + .2 * 80 + .2 * 75 + .3 * 90 = 84.25.$$

Note:  $\mathbf{w}$  can be thought of as a vector of “weights” and  $\mathbf{g}$  as a vector of information. Then the dot product of the two is a weighted average.

# Another Representation for the Dot product

You can interpret the inner product in two ways:

1. As an operation that is done to two vectors.

# Another Representation for the Dot product

You can interpret the inner product in two ways:

1. As an operation that is done to two vectors.
2. As an “action” a covector does to a vector.

# Another Representation for the Dot product

You can interpret the inner product in two ways:

1. As an operation that is done to two vectors.
2. As an “action” a covector does to a vector.

In the second interpretation the left vector in the dot product is represented as a covector:

$$\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w}.$$

# Another Representation for the Dot product

You can interpret the inner product in two ways:

1. As an operation that is done to two vectors.
2. As an “action” a covector does to a vector.

In the second interpretation the left vector in the dot product is represented as a covector:

$$\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w}.$$

In components:

$$\begin{bmatrix} v_1 & v_2 & \dots & v_n \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \\ \vdots \\ w_n \end{bmatrix} = \sum_{i=1}^n v_i w_i.$$

# Matrix multiplication

## Definition

Let  $A = [a_{ik}] \in \mathbb{R}^{m \times \ell}$  and  $B = [b_{kj}] \in \mathbb{R}^{\ell \times n}$ . Then the **product** of  $A$  and  $B$  is the matrix

$$AB := \left[ \sum_{k=1}^n a_{ik} b_{kj} \right] \in \mathbb{R}^{m \times n}.$$

# Matrix multiplication

## Definition

Let  $A = [a_{ik}] \in \mathbb{R}^{m \times \ell}$  and  $B = [b_{kj}] \in \mathbb{R}^{\ell \times n}$ . Then the **product** of  $A$  and  $B$  is the matrix

$$AB := \left[ \sum_{k=1}^n a_{ik} b_{kj} \right] \in \mathbb{R}^{m \times n}.$$

In words: The  $i, j$ th element of  $AB$  is the action of the  $i$ th row of  $A$  on the  $j$ th column of  $B$ .

## Example

Let

$$A = \begin{bmatrix} 1 & 0 \\ 2 & 1 \\ 3 & 2 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \end{bmatrix}$$



## Example

Let

$$A = \begin{bmatrix} 1 & 0 \\ 2 & 1 \\ 3 & 2 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \end{bmatrix}$$

Then

$$AB =$$

## Example

Let

$$A = \begin{bmatrix} 1 & 0 \\ 2 & 1 \\ 3 & 2 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \end{bmatrix}$$

Then

$$AB = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 10 \\ 7 & 8 & 17 \end{bmatrix}.$$

## Example

Let

$$A = \begin{bmatrix} 1 & 0 \\ 2 & 1 \\ 3 & 2 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \end{bmatrix}$$

Then

$$AB = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 10 \\ 7 & 8 & 17 \end{bmatrix}.$$

and

$$BA =$$

## Example

Let

$$A = \begin{bmatrix} 1 & 0 \\ 2 & 1 \\ 3 & 2 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \end{bmatrix}$$

Then

$$AB = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 10 \\ 7 & 8 & 17 \end{bmatrix}.$$

and

$$BA = \begin{bmatrix} 14 & 8 \\ 16 & 9 \end{bmatrix}.$$

# Comments

1. From our example  $AB \neq BA$ . In general matrix multiplication *does not commute!*

# Comments

1. From our example  $AB \neq BA$ . In general matrix multiplication *does not commute*!
2.  $AB$  only makes sense if number of columns of  $A$  is the same as the number of rows of  $B$ .

# Comments

1. From our example  $AB \neq BA$ . In general matrix multiplication *does not commute!*
2.  $AB$  only makes sense if number of columns of  $A$  is the same as the number of rows of  $B$ .
3. The number of rows of  $AB$  is the same as the number of rows of  $A$ .

# Comments

1. From our example  $AB \neq BA$ . In general matrix multiplication *does not commute*!
2.  $AB$  only makes sense if number of columns of  $A$  is the same as the number of rows of  $B$ .
3. The number of rows of  $AB$  is the same as the number of rows of  $A$ .
4. The number or columns of  $AB$  is the same as the number of columns of  $B$ .



# Why is this Our Definition for Matrix Multiplication?

# Why is this Our Definition for Matrix Multiplication?

This choice of definition for matrix multiplication is USEFUL:

# Why is this Our Definition for Matrix Multiplication?

This choice of definition for matrix multiplication is USEFUL:

- ▶ This definition agrees with the the behavior of how linear operators are composed (TBD in a month or so).

# Why is this Our Definition for Matrix Multiplication?

This choice of definition for matrix multiplication is USEFUL:

- ▶ This definition agrees with the the behavior of how linear operators are composed (TBD in a month or so).
- ▶ The special case of a matrix-vector product gives us useful interpretations.

## Special Case: Matrix-Vector Product

Let  $A \in \mathbb{R}^{m \times n}$  and  $x \in \mathbb{R}^n$ .

## Special Case: Matrix-Vector Product

Let  $A \in \mathbb{R}^{m \times n}$  and  $x \in \mathbb{R}^n$ . Then:

$$Ax = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix}$$

## Special Case: Matrix-Vector Product

Let  $A \in \mathbb{R}^{m \times n}$  and  $x \in \mathbb{R}^n$ . Then:

$$Ax = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix}$$
$$= \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \cdots + a_{mn}x_n \end{bmatrix} \in \mathbb{R}^m.$$

## Interpretation (1): $Ax$ is a linear combination

With  $A$  and  $x$  as before we have:

$$Ax = \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \cdots + a_{mn}x_n \end{bmatrix}$$

$$= x_1 \begin{bmatrix} a_{11} \\ a_{21} \\ a_{31} \\ \vdots \\ a_{m1} \end{bmatrix} + x_2 \begin{bmatrix} a_{12} \\ a_{22} \\ a_{32} \\ \vdots \\ a_{m2} \end{bmatrix} + x_3 \begin{bmatrix} a_{13} \\ a_{23} \\ a_{33} \\ \vdots \\ a_{m3} \end{bmatrix} + \cdots + x_n \begin{bmatrix} a_{1n} \\ a_{2n} \\ a_{3n} \\ \vdots \\ a_{mn} \end{bmatrix}$$



## Interpretation (1): $Ax$ is a linear combination

With  $A$  and  $x$  as before we have:

$$Ax = \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \cdots + a_{mn}x_n \end{bmatrix}$$

$$= x_1 \begin{bmatrix} a_{11} \\ a_{21} \\ a_{31} \\ \vdots \\ a_{m1} \end{bmatrix} + x_2 \begin{bmatrix} a_{12} \\ a_{22} \\ a_{32} \\ \vdots \\ a_{m2} \end{bmatrix} + x_3 \begin{bmatrix} a_{13} \\ a_{23} \\ a_{33} \\ \vdots \\ a_{m3} \end{bmatrix} + \cdots + x_n \begin{bmatrix} a_{1n} \\ a_{2n} \\ a_{3n} \\ \vdots \\ a_{mn} \end{bmatrix}$$

So  $Ax$  is a *linear combination* of the columns of  $A$ .

## Interpretation (2): $Ax = b$ is a system of linear equations

Now let  $A$  and  $x$  be as before and consider a vector  $b = [b_i] \in \mathbb{R}^m$

## Interpretation (2): $Ax = b$ is a system of linear equations

Now let  $A$  and  $x$  be as before and consider a vector  $b = [b_i] \in \mathbb{R}^m$   
Then

$$Ax = b \Leftrightarrow \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \cdots + a_{mn}x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ \vdots \\ b_m \end{bmatrix}$$

$$\Leftrightarrow \begin{array}{ccccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & a_{13}x_3 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & a_{23}x_3 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & & \vdots & & \vdots & & & & \vdots & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & a_{m3}x_3 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array}$$

## Interpretation (2): $Ax = b$ is a system of linear equations

Now let  $A$  and  $x$  be as before and consider a vector  $b = [b_i] \in \mathbb{R}^m$   
Then

$$Ax = b \Leftrightarrow \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \cdots + a_{mn}x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ \vdots \\ b_m \end{bmatrix}$$

$$\Leftrightarrow \begin{array}{ccccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & a_{13}x_3 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & a_{23}x_3 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & & \vdots & & \vdots & & & & \vdots & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & a_{m3}x_3 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array}$$

**$Ax = b$  is a system of linear equations!**

## Example

Consider the system of linear equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

Interpret this system as a matrix-vector equation  $Ax = b$ . Write out each of  $A$ ,  $x$ , and  $b$  explicitly.

## Example

Consider the system of linear equations

$$\begin{aligned}x + 2y + 3z &= 6 \\2x - 3y + 2z &= 14 \\3x + y - z &= -2\end{aligned}$$

Interpret this system as a matrix-vector equation  $Ax = b$ . Write out each of  $A$ ,  $x$ , and  $b$  explicitly.

**Solution:**

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -3 & 2 \\ 3 & 1 & -1 \end{bmatrix} \quad x = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \quad b = \begin{bmatrix} 6 \\ 14 \\ -2 \end{bmatrix}$$

## Example

Consider the system of linear equations

$$\begin{aligned}x + 2y + 3z &= 6 \\2x - 3y + 2z &= 14 \\3x + y - z &= -2\end{aligned}$$

Interpret this system as a matrix-vector equation  $Ax = b$ . Write out each of  $A$ ,  $x$ , and  $b$  explicitly.

**Solution:**

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -3 & 2 \\ 3 & 1 & -1 \end{bmatrix} \quad x = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \quad b = \begin{bmatrix} 6 \\ 14 \\ -2 \end{bmatrix}$$

When interpreted as a system of equations,  $A$  is called the **coefficient matrix** of the system.

# Augmented Matrix for a System of Equations

In the prior example, when interpreted as a system of equations,  $Ax = b$  has two “known” parts:  $A$  and  $b$ .  $x$  is considered to be a vector of “unknowns.”



# Augmented Matrix for a System of Equations

In the prior example, when interpreted as a system of equations,  $Ax = b$  has two “known” parts:  $A$  and  $b$ .  $x$  is considered to be a vector of “unknowns.”

A **augmented matrix** is created from the “known” quantities:

$$\left[ A \mid b \right] = \left[ \begin{array}{ccccc|c} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} & b_2 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} & b_m \end{array} \right]$$

## Example

Write out the system of equations which corresponds to the following augmented matrix:

$$\left[ \begin{array}{ccc|c} 2 & -1 & 3 & 4 \\ 3 & 0 & 2 & 5 \end{array} \right]$$

## Example

Write out the system of equations which corresponds to the following augmented matrix:

$$\left[ \begin{array}{ccc|c} 2 & -1 & 3 & 4 \\ 3 & 0 & 2 & 5 \end{array} \right]$$

**Solution:**

$$2x - y + 3z = 4$$

$$3x + 2z = 5$$