

Review: Geometric Realization of Vectors in \mathbb{R}^n

Recall that vectors in \mathbb{R}^n can be thought of as “arrows.”

Exercise: Graph the vectors u , v , and $u - v$ where:

$$u = \begin{bmatrix} -1 \\ 5 \end{bmatrix} \quad v = \begin{bmatrix} 3 \\ 2 \end{bmatrix}.$$


Review: Geometric Realization of Vectors in \mathbb{R}^n

Recall that vectors in \mathbb{R}^n can be thought of as “arrows.”

Exercise: Graph the vectors u , v , and $u - v$ where:

$$u = \begin{bmatrix} -1 \\ 5 \end{bmatrix} \quad v = \begin{bmatrix} 3 \\ 2 \end{bmatrix}.$$

Sln:



Two Interesting questions:

When considering an “arrow” in \mathbb{R}^n there are two interesting questions:

Two Interesting questions:

When considering an “arrow” in \mathbb{R}^n there are two interesting questions:

1. How long is it?

Two Interesting questions:

When considering an “arrow” in \mathbb{R}^n there are two interesting questions:

1. How long is it?
2. What direction is it pointing?

Two Interesting questions:

When considering an “arrow” in \mathbb{R}^n there are two interesting questions:

1. How long is it?
2. What direction is it pointing?

It turns out that there is a *single* computation which can help us to answer both questions!

Two Interesting questions:

When considering an “arrow” in \mathbb{R}^n there are two interesting questions:

1. How long is it?
2. What direction is it pointing?


It turns out that there is a *single* computation which can help us to answer both questions!

But we will tackle one-at-a-time.

Length of a Vector

Consider the generic vector

$$u = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$$




The “length” of u should be the distance from O to the head of u .
Pythagoras gives:

$$\text{length of } u = \sqrt{(u_1)^2 + (u_2)^2}.$$

Length of a Vector

Consider the generic vector


$$u = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$$



Length of a Vector

Consider the generic vector

$$u = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$$




The “length” of u should be the distance from O to the head of u .

Length of a Vector

Consider the generic vector

$$u = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$$



The “length” of u should be the distance from O to the head of u .
Pythagoras gives:

$$\text{length of } u = \sqrt{(u_1)^2 + (u_2)^2}.$$

Length/Magnitude/Norm

Definition

Let $u = [u_i] \in \mathbb{R}^n$. The (standard) **length** (or **magnitude** or **norm**) of u is

$$\|u\| = \sqrt{\sum_{i=1}^n u_i^2}.$$

Length/Magnitude/Norm

Definition

Let $u = [u_i] \in \mathbb{R}^n$. The (standard) **length** (or **magnitude** or **norm**) of u is

$$\|u\| = \sqrt{\sum_{i=1}^n u_i^2}.$$

Exercise: Compute the lengths of $u = \begin{bmatrix} -1 \\ 5 \end{bmatrix}$, $v = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$ and $u - v$.

Length/Magnitude/Norm

Definition

Let $u = [u_i] \in \mathbb{R}^n$. The (standard) **length** (or **magnitude** or **norm**) of u is

$$\|u\| = \sqrt{\sum_{i=1}^n u_i^2}.$$

Exercise: Compute the lengths of $u = \begin{bmatrix} -1 \\ 5 \end{bmatrix}$, $v = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$ and $u - v$.

Solutions:

$$\|u\| = \sqrt{(-1)^2 + 5^2} = \sqrt{26},$$

Length/Magnitude/Norm

Definition

Let $u = [u_i] \in \mathbb{R}^n$. The (standard) **length** (or **magnitude** or **norm**) of u is

$$\|u\| = \sqrt{\sum_{i=1}^n u_i^2}.$$


Exercise: Compute the lengths of $u = \begin{bmatrix} -1 \\ 5 \end{bmatrix}$, $v = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$ and $u - v$.

Solutions:

$$\|u\| = \sqrt{(-1)^2 + 5^2} = \sqrt{26}, \quad \|v\| = \sqrt{13}, \quad \|u - v\| = \sqrt{25}.$$


Distance Between Vectors

Let's return to our vectors u , v , and $u - v$:



Distance Between Vectors

Let's return to our vectors u , v , and $u - v$:



Definition


The **distance** between the vectors u and v in \mathbb{R}^n is

$$d(u, v) := \|u - v\|.$$

Note: Distance measurements are CRUCIAL to answer questions like “How good is our approximation?”


Direction of a Vector

Return to our generic vector in \mathbb{R}^2 :



Direction of a Vector

Return to our generic vector in \mathbb{R}^2 :




Computing θ is straightforward:

$$\tan(\theta) = \frac{u_2}{u_1}$$

Direction of a Vector

Return to our generic vector in \mathbb{R}^2 :




Computing θ is straightforward:

$$\tan(\theta) = \frac{u_2}{u_1} \quad \text{or} \quad \cos \theta = \frac{u_1}{\sqrt{(u_1)^2 + (u_2)^2}} = \frac{u_1}{\|u\|}.$$

Direction of a Vector

Return to our generic vector in \mathbb{R}^2 :




Computing θ is straightforward:

$$\tan(\theta) = \frac{u_2}{u_1} \quad \text{or} \quad \cos \theta = \frac{u_1}{\sqrt{(u_1)^2 + (u_2)^2}} = \frac{u_1}{\|u\|}.$$

Note: the norm of u in the denominator.

Direction of a Vector

Return to our generic vector in \mathbb{R}^2 :




Computing θ is straightforward:

$$\tan(\theta) = \frac{u_2}{u_1} \quad \text{or} \quad \cos \theta = \frac{u_1}{\sqrt{(u_1)^2 + (u_2)^2}} = \frac{u_1}{\|u\|}.$$

Note: the norm of u in the denominator. Somehow the angle is related to sizes.


Angle Between Two Vectors

A slightly longer computation:



Angle Between Two Vectors

A slightly longer computation:




Apply the “law of cosines”:

$$\|u - v\|^2 = \|u\|^2 + \|v\|^2 - 2\|u\|\|v\|\cos(\theta).$$

Angle Between Two Vectors

A slightly longer computation:



Apply the “law of cosines”:

$$\|u - v\|^2 = \|u\|^2 + \|v\|^2 - 2\|u\|\|v\|\cos(\theta).$$

After some expansion and algebra you can show

$$\cos(\theta) = \frac{u_1 v_1 + u_2 v_2}{\|u\|\|v\|}.$$

An oft-repeated computation:

An oft-repeated computation:

- ▶ Length:

$$\|u\|^2 = (u_1)^2 + (u_2)^2$$

An oft-repeated computation:

- ▶ Length:

$$\|u\|^2 = (u_1)^2 + (u_2)^2 = u_1 u_1 + u_2 u_2.$$

An oft-repeated computation:

- ▶ Length:

$$\|u\|^2 = (u_1)^2 + (u_2)^2 = u_1 u_1 + u_2 u_2.$$

- ▶ Angle:

$$\cos(\theta) = \frac{u_1 v_1 + u_2 v_2}{\|u\| \|v\|}.$$

An oft-repeated computation:

- ▶ Length:

$$\|u\|^2 = (u_1)^2 + (u_2)^2 = u_1 u_1 + u_2 u_2.$$

- ▶ Angle:

$$\cos(\theta) = \frac{u_1 v_1 + u_2 v_2}{\|u\| \|v\|}.$$

- ▶ Multiplying matrices together:

$$(AB)_{ij} = A_{i1}B_{1j} + A_{i2}B_{2j} + \cdots + A_{in}B_{nj}.$$

An oft-repeated computation:

- ▶ Length:

$$\|u\|^2 = (u_1)^2 + (u_2)^2 = u_1 u_1 + u_2 u_2.$$

- ▶ Angle:

$$\cos(\theta) = \frac{u_1 v_1 + u_2 v_2}{\|u\| \|v\|}.$$

- ▶ Multiplying matrices together:

$$(AB)_{ij} = A_{i1}B_{1j} + A_{i2}B_{2j} + \cdots + A_{in}B_{nj}.$$

Multiplying individual vector coordinates and then summing comes up in many contexts.

Dot Product

Definition

Let u and v be vectors in \mathbb{R}^n . The **dot product** or **standard inner product** of u and v is

$$u \cdot v := \sum_{i=1}^n u_i v_i$$

Dot Product

Definition

Let u and v be vectors in \mathbb{R}^n . The **dot product** or **standard inner product** of u and v is

$$u \cdot v := \sum_{i=1}^n u_i v_i$$

Using this, we have some new expressions:

Dot Product

Definition

Let u and v be vectors in \mathbb{R}^n . The **dot product** or **standard inner product** of u and v is

$$u \cdot v := \sum_{i=1}^n u_i v_i$$

Using this, we have some new expressions:

- ▶ Length: $\|u\| = \sqrt{u \cdot u}$.

Dot Product

Definition

Let u and v be vectors in \mathbb{R}^n . The **dot product** or **standard inner product** of u and v is

$$u \cdot v := \sum_{i=1}^n u_i v_i$$

Using this, we have some new expressions:

- ▶ Length: $\|u\| = \sqrt{u \cdot u}$.
- ▶ Angle: $\cos(\theta) = \frac{u \cdot v}{\|u\| \|v\|}$.

Dot Product

Definition

Let u and v be vectors in \mathbb{R}^n . The **dot product** or **standard inner product** of u and v is

$$u \cdot v := \sum_{i=1}^n u_i v_i$$

Using this, we have some new expressions:

- ▶ Length: $\|u\| = \sqrt{u \cdot u}$.
- ▶ Angle: $\cos(\theta) = \frac{u \cdot v}{\|u\| \|v\|}$.

We can also say that the element in the (i, j) -th position in the product matrix AB is the dot product of the i -th row of A and the j -th column of B .

Properties of the Dot Product

Theorem

Let u , v , and w be vectors in \mathbb{R}^n and c a scalar

Properties of the Dot Product

Theorem

Let u , v , and w be vectors in \mathbb{R}^n and c a scalar

1. $u \cdot u \geq 0$.

Properties of the Dot Product

Theorem

Let u , v , and w be vectors in \mathbb{R}^n and c a scalar

1. $u \cdot u \geq 0$.
2. $u \cdot u = 0 \Leftrightarrow u = 0$.

Properties of the Dot Product

Theorem

Let u , v , and w be vectors in \mathbb{R}^n and c a scalar

1. $u \cdot u \geq 0$.
2. $u \cdot u = 0 \Leftrightarrow u = 0$.
3. *Commutativity:* $u \cdot v = v \cdot u$

Properties of the Dot Product

Theorem

Let u , v , and w be vectors in \mathbb{R}^n and c a scalar

1. $u \cdot u \geq 0$.
2. $u \cdot u = 0 \Leftrightarrow u = 0$.
3. *Commutativity:* $u \cdot v = v \cdot u$
4. *Distributivity:* $u \cdot (v + w) = u \cdot v + u \cdot w$.

Properties of the Dot Product

Theorem

Let u , v , and w be vectors in \mathbb{R}^n and c a scalar

1. $u \cdot u \geq 0$.
2. $u \cdot u = 0 \Leftrightarrow u = 0$.
3. *Commutativity:* $u \cdot v = v \cdot u$
4. *Distributivity:* $u \cdot (v + w) = u \cdot v + u \cdot w$.
5. *Scalar associativity:* $c(u \cdot v) = (cu) \cdot v$.

Properties of the Dot Product

Theorem

Let u , v , and w be vectors in \mathbb{R}^n and c a scalar

1. $u \cdot u \geq 0$.
2. $u \cdot u = 0 \Leftrightarrow u = 0$.
3. Commutativity: $u \cdot v = v \cdot u$
4. Distributivity: $u \cdot (v + w) = u \cdot v + u \cdot w$.
5. Scalar associativity: $c(u \cdot v) = (cu) \cdot v$.
6. Orthogonality property: $u \perp v \Leftrightarrow u \cdot v = 0$.

Simple Application: unit vectors

Definition

A **unit** vector is a vector with the property that $\|u\| = 1$.

Simple Application: unit vectors

Definition

A **unit** vector is a vector with the property that $\|u\| = 1$.

Exercises:

1. Show that $u = \begin{bmatrix} 3/5 \\ 4/5 \end{bmatrix}$ is a unit vector.
2. Show that u above and the vector $\begin{bmatrix} 6 \\ 8 \end{bmatrix}$ are parallel. (What is the angle between them?)
3. Find a unit vector in the same direction as $v = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$. (What value can you use to *scale* v ?)