

## Exercise

Compute the following determinant in two ways 1) using elimination to reduce to upper (or lower) triangular form and 2) by cofactor expansion.

$$\begin{vmatrix} 2 & 0 & 4 \\ 1 & 1 & 2 \\ 1 & 0 & 1 \end{vmatrix}$$

# Solutions:

Via elimination:

$$\underbrace{\begin{vmatrix} 2 & 0 & 4 \\ 1 & 1 & 2 \\ 1 & 0 & 1 \end{vmatrix}}_{\det(A)}$$

# Solutions:

Via elimination:

$$\underbrace{\begin{vmatrix} 2 & 0 & 4 \\ 1 & 1 & 2 \\ 1 & 0 & 1 \end{vmatrix}}_{\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 1 & 1 & 2 \\ 2 & 0 & 4 \end{vmatrix}}_{-\det(A)}$$

# Solutions:

Via elimination:

$$\underbrace{\begin{vmatrix} 2 & 0 & 4 \\ 1 & 1 & 2 \\ 1 & 0 & 1 \end{vmatrix}}_{\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 1 & 1 & 2 \\ 2 & 0 & 4 \end{vmatrix}}_{-\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 2 \end{vmatrix}}_{-\det(A)}$$

# Solutions:

Via elimination:

$$\underbrace{\begin{vmatrix} 2 & 0 & 4 \\ 1 & 1 & 2 \\ 1 & 0 & 1 \end{vmatrix}}_{\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 1 & 1 & 2 \\ 2 & 0 & 4 \end{vmatrix}}_{-\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 2 \end{vmatrix}}_{-\det(A)}$$

## Solutions:

Via elimination:

$$\underbrace{\begin{vmatrix} 2 & 0 & 4 \\ 1 & 1 & 2 \\ 1 & 0 & 1 \end{vmatrix}}_{\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 1 & 1 & 2 \\ 2 & 0 & 4 \end{vmatrix}}_{-\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 2 \end{vmatrix}}_{-\det(A)}$$

So

$$-\det(A) = 2 \Leftrightarrow \det(A) = -2.$$

## Solutions:

Via elimination:

$$\underbrace{\begin{vmatrix} 2 & 0 & 4 \\ 1 & 1 & 2 \\ 1 & 0 & 1 \end{vmatrix}}_{\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 1 & 1 & 2 \\ 2 & 0 & 4 \end{vmatrix}}_{-\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 2 \end{vmatrix}}_{-\det(A)}$$

So

$$-\det(A) = 2 \Leftrightarrow \det(A) = -2.$$

Via cofactor expansion.

## Solutions:

Via elimination:

$$\underbrace{\begin{vmatrix} 2 & 0 & 4 \\ 1 & 1 & 2 \\ 1 & 0 & 1 \end{vmatrix}}_{\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 1 & 1 & 2 \\ 2 & 0 & 4 \end{vmatrix}}_{-\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 2 \end{vmatrix}}_{-\det(A)}$$

So

$$-\det(A) = 2 \Leftrightarrow \det(A) = -2.$$

Via cofactor expansion. Which row/column should we use?



# Solutions:

Via elimination:

$$\underbrace{\begin{vmatrix} 2 & 0 & 4 \\ 1 & 1 & 2 \\ 1 & 0 & 1 \end{vmatrix}}_{\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 1 & 1 & 2 \\ 2 & 0 & 4 \end{vmatrix}}_{-\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 2 \end{vmatrix}}_{-\det(A)}$$

So

$$-\det(A) = 2 \Leftrightarrow \det(A) = -2.$$

Via cofactor expansion. Which row/column should we use? 2nd column:

$$\det(A) = (0)(-1) \begin{vmatrix} 1 & 2 \\ 1 & 1 \end{vmatrix} + (1)(1) \begin{vmatrix} 2 & 4 \\ 1 & 1 \end{vmatrix} + (0)(-1) \begin{vmatrix} 2 & 4 \\ 1 & 2 \end{vmatrix}$$

## Solutions:

Via elimination:

$$\underbrace{\begin{vmatrix} 2 & 0 & 4 \\ 1 & 1 & 2 \\ 1 & 0 & 1 \end{vmatrix}}_{\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 1 & 1 & 2 \\ 2 & 0 & 4 \end{vmatrix}}_{-\det(A)} \sim \underbrace{\begin{vmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 2 \end{vmatrix}}_{-\det(A)}$$

So

$$-\det(A) = 2 \Leftrightarrow \det(A) = -2.$$

Via cofactor expansion. Which row/column should we use? 2nd column:

$$\det(A) = (0)(-1) \begin{vmatrix} 1 & 2 \\ 1 & 1 \end{vmatrix} + (1)(1) \begin{vmatrix} 2 & 4 \\ 1 & 1 \end{vmatrix} + (0)(-1) \begin{vmatrix} 2 & 4 \\ 1 & 2 \end{vmatrix}$$

So

$$\det(A) = 1(2 - 4) = -2.$$

# Today: Inverse matrices can be written in terms of cofactors

# Today: Inverse matrices can be written in terms of cofactors

Recall that for  $A = [a_{ij}] \in \mathbb{R}^{2 \times 2}$ :

$$A^{-1} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}^{-1} =$$

# Today: Inverse matrices can be written in terms of cofactors

Recall that for  $A = [a_{ij}] \in \mathbb{R}^{2 \times 2}$ :

$$A^{-1} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}^{-1} = \frac{1}{a_{11}a_{22} - a_{12}a_{21}} \begin{bmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{bmatrix}$$

# Today: Inverse matrices can be written in terms of cofactors

Recall that for  $A = [a_{ij}] \in \mathbb{R}^{2 \times 2}$ :

$$A^{-1} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}^{-1} = \frac{1}{a_{11}a_{22} - a_{12}a_{21}} \begin{bmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{bmatrix}$$

Recall that the  $ij$ -th cofactor is

$$A_{ij} = (-1)^{i+j} |M_{ij}|$$

where  $M_{ij}$  is the *minor* matrix created by removing the  $i$ th row and  $j$ th column from  $A$ .

# Today: Inverse matrices can be written in terms of cofactors

Recall that for  $A = [a_{ij}] \in \mathbb{R}^{2 \times 2}$ :

$$A^{-1} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}^{-1} = \frac{1}{a_{11}a_{22} - a_{12}a_{21}} \begin{bmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{bmatrix}$$

Recall that the  $ij$ -th cofactor is

$$A_{ij} = (-1)^{i+j} |M_{ij}|$$

where  $M_{ij}$  is the *minor* matrix created by removing the  $i$ th row and  $j$ th column from  $A$ .

Exercise: Write  $A^{-1}$  above in terms of  $|A|$  and the cofactors  $A_{ij}$ .

# Cofactor Representation for $2 \times 2$ Inverse

We have

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$



# Cofactor Representation for $2 \times 2$ Inverse

We have

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

The cofactors are:

$$A_{11} = a_{22} \quad A_{12} = -a_{21} \quad A_{21} = -a_{12} \quad A_{22} = a_{11}$$

## Cofactor Representation for $2 \times 2$ Inverse

We have

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

The cofactors are:

$$A_{11} = a_{22} \quad A_{12} = -a_{21} \quad A_{21} = -a_{12} \quad A_{22} = a_{11}$$

So:

$$A^{-1} = \frac{1}{a_{11}a_{22} - a_{12}a_{21}} \begin{bmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{bmatrix} = \frac{1}{|A|} \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}^T.$$

## Cofactor Representation for $2 \times 2$ Inverse

We have

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

The cofactors are:

$$A_{11} = a_{22} \quad A_{12} = -a_{21} \quad A_{21} = -a_{12} \quad A_{22} = a_{11}$$

So:

$$A^{-1} = \frac{1}{a_{11}a_{22} - a_{12}a_{21}} \begin{bmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{bmatrix} = \frac{1}{|A|} \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}^T.$$

Question: If  $A$  is a general non-singular matrix, can we write  $A^{-1}$  in terms of  $|A|$  and its cofactors?

# Cofactor Representation for Inverse Matrices

## Theorem

Let  $A = [a_{ij}]$  be a nonsingular matrix. Then

$$A^{-1} = \frac{1}{\det A} [A_{ij}]^T.$$

# Cofactor Representation for Inverse Matrices

## Theorem

Let  $A = [a_{ij}]$  be a nonsingular matrix. Then

$$A^{-1} = \frac{1}{\det A} [A_{ij}]^T.$$

Exercise: Use this theorem to compute the inverse of

$$A = \begin{bmatrix} 3 & -2 & 1 \\ 5 & 6 & 2 \\ 1 & 0 & -3 \end{bmatrix}.$$

# Cofactor Representation for Inverse Matrices

## Theorem

Let  $A = [a_{ij}]$  be a nonsingular matrix. Then

$$A^{-1} = \frac{1}{\det A} [A_{ij}]^T.$$

Exercise: Use this theorem to compute the inverse of

$$A = \begin{bmatrix} 3 & -2 & 1 \\ 5 & 6 & 2 \\ 1 & 0 & -3 \end{bmatrix}.$$

Stop.

This is a terrible way to compute inverse matrices.

# Computing determinants is EXPENSIVE

# Computing determinants is EXPENSIVE

To compute the determinant of a  $25 \times 25$  matrix



# Computing determinants is EXPENSIVE

To compute the determinant of a  $25 \times 25$  matrix (which is teensy-tiny)

# Computing determinants is EXPENSIVE

To compute the determinant of a  $25 \times 25$  matrix (which is teensy-tiny) via cofactor expansion would take  $25!$  multiplications.

# Computing determinants is EXPENSIVE

To compute the determinant of a  $25 \times 25$  matrix (which is teensy-tiny) via cofactor expansion would take  $25!$  multiplications.

This is approximately  $1.55 \times 10^{25}$  operations.

# Computing determinants is EXPENSIVE

To compute the determinant of a  $25 \times 25$  matrix (which is teensy-tiny) via cofactor expansion would take  $25!$  multiplications.

This is approximately  $1.55 \times 10^{25}$  operations.

The fastest supercomputer in the world can make  $122.3 \times 10^{15}$  computations per second

# Computing determinants is EXPENSIVE

To compute the determinant of a  $25 \times 25$  matrix (which is teensy-tiny) via cofactor expansion would take  $25!$  multiplications.

This is approximately  $1.55 \times 10^{25}$  operations.

The fastest supercomputer in the world can make  $122.3 \times 10^{15}$  computations per second and there are about  $3 \times 10^7$  seconds per year.

# Computing determinants is EXPENSIVE

To compute the determinant of a  $25 \times 25$  matrix (which is teensy-tiny) via cofactor expansion would take  $25!$  multiplications.

This is approximately  $1.55 \times 10^{25}$  operations.

The fastest supercomputer in the world can make  $122.3 \times 10^{15}$  computations per second and there are about  $3 \times 10^7$  seconds per year.

In total:

$$\frac{1.55 \times 10^{25}}{3.67 \times 10^{24}} \sim 4.2 \text{ years!}$$

# Computing determinants is EXPENSIVE

To compute the determinant of a  $25 \times 25$  matrix (which is teensy-tiny) via cofactor expansion would take  $25!$  multiplications.

This is approximately  $1.55 \times 10^{25}$  operations.

The fastest supercomputer in the world can make  $122.3 \times 10^{15}$  computations per second and there are about  $3 \times 10^7$  seconds per year.

In total:

$$\frac{1.55 \times 10^{25}}{3.67 \times 10^{24}} \sim 4.2 \text{ years!}$$

But Matlab will do it in under a second.

# Computing determinants is EXPENSIVE

To compute the determinant of a  $25 \times 25$  matrix (which is teensy-tiny) via cofactor expansion would take  $25!$  multiplications.

This is approximately  $1.55 \times 10^{25}$  operations.

The fastest supercomputer in the world can make  $122.3 \times 10^{15}$  computations per second and there are about  $3 \times 10^7$  seconds per year.

In total:

$$\frac{1.55 \times 10^{25}}{3.67 \times 10^{24}} \sim 4.2 \text{ years!}$$

But Matlab will do it in under a second. Because Matlab uses row reduction to an upper triangular matrix (in fact, it does an LU-factorization).



# Computing determinants is EXPENSIVE

To compute the determinant of a  $25 \times 25$  matrix (which is teensy-tiny) via cofactor expansion would take  $25!$  multiplications.

This is approximately  $1.55 \times 10^{25}$  operations.

The fastest supercomputer in the world can make  $122.3 \times 10^{15}$  computations per second and there are about  $3 \times 10^7$  seconds per year.

In total:

$$\frac{1.55 \times 10^{25}}{3.67 \times 10^{24}} \sim 4.2 \text{ years!}$$

But Matlab will do it in under a second. Because Matlab uses row reduction to an upper triangular matrix (in fact, it does an LU-factorization). So you should too.

# Why Inverses via Cofactors is Useful

Let  $A = [a_{ij}]$  be a non-singular  $n \times n$  matrix and  $b$  a  $n$ -vector.  
Then we know that the unique solution to

$$Ax = b$$

# Why Inverses via Cofactors is Useful

Let  $A = [a_{ij}]$  be a non-singular  $n \times n$  matrix and  $b$  a  $n$ -vector.  
Then we know that the unique solution to

$$Ax = b \text{ is } x = A^{-1}b.$$

# Why Inverses via Cofactors is Useful

Let  $A = [a_{ij}]$  be a non-singular  $n \times n$  matrix and  $b$  a  $n$ -vector. Then we know that the unique solution to

$$Ax = b \text{ is } x = A^{-1}b.$$

Let us write  $A^{-1}$  in terms of the cofactors and  $x$  and  $b$  in terms of their components:

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \frac{1}{|A|} \begin{bmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ A_{1n} & A_{2n} & \dots & A_{nn} \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}.$$

So

$$x_i = \frac{1}{|A|} \sum_{k=1}^n b_k A_{ki}$$

# What is $x_i$ ?

We have

$$x_i = \frac{1}{|A|} (b_1 A_{1i} + b_2 A_{2i} + \cdots + b_n A_{ni})$$

## What is $x_i$ ?

We have

$$\begin{aligned}x_i &= \frac{1}{|A|} (b_1 A_{1i} + b_2 A_{2i} + \cdots + b_n A_{ni}) \\&= \frac{1}{|A|} \underbrace{(b_1 (-1)^{1+i} |M_{1i}| + b_2 (-1)^{2+i} |M_{2i}| + \cdots + b_n (-1)^{n+i} |M_{ni}|)}\end{aligned}$$

# What is $x_i$ ?

We have

$$\begin{aligned}x_i &= \frac{1}{|A|} (b_1 A_{1i} + b_2 A_{2i} + \cdots + b_n A_{ni}) \\&= \frac{1}{|A|} \underbrace{(b_1 (-1)^{1+i} |M_{1i}| + b_2 (-1)^{2+i} |M_{2i}| + \cdots + b_n (-1)^{n+i} |M_{ni}|)}_{\text{This looks like a determinant!}}\end{aligned}$$

# What is $x_i$ ?

We have

$$\begin{aligned}x_i &= \frac{1}{|A|} (b_1 A_{1i} + b_2 A_{2i} + \cdots + b_n A_{ni}) \\&= \frac{1}{|A|} \underbrace{(b_1 (-1)^{1+i} |M_{1i}| + b_2 (-1)^{2+i} |M_{2i}| + \cdots + b_n (-1)^{n+i} |M_{ni}|)}_{\text{This looks like a determinant! But of what?}}\end{aligned}$$



## What is $x_i$ ?

We have

$$\begin{aligned}x_i &= \frac{1}{|A|} (b_1 A_{1i} + b_2 A_{2i} + \cdots + b_n A_{ni}) \\&= \frac{1}{|A|} \underbrace{(b_1 (-1)^{1+i} |M_{1i}| + b_2 (-1)^{2+i} |M_{2i}| + \cdots + b_n (-1)^{n+i} |M_{ni}|)}_{\text{This looks like a determinant! But of what?}}\end{aligned}$$

From the indices, it looks like we are expanding through the  $i$ th column.

# What is $x_i$ ?

We have

$$\begin{aligned}x_i &= \frac{1}{|A|} (b_1 A_{1i} + b_2 A_{2i} + \cdots + b_n A_{ni}) \\&= \frac{1}{|A|} \underbrace{(b_1 (-1)^{1+i} |M_{1i}| + b_2 (-1)^{2+i} |M_{2i}| + \cdots + b_n (-1)^{n+i} |M_{ni}|)}_{\text{This looks like a determinant! But of what?}}\end{aligned}$$

From the indices, it looks like we are expanding through the  $i$ th column. So it should be

$$x_i = \frac{1}{|A|} \begin{vmatrix} a_{11} & a_{12} & \cdots & a_{1,i-1} & b_1 & a_{1,i+1} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2,i-1} & b_2 & a_{2,i+1} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{n,i-1} & b_n & a_{n,i+1} & \cdots & a_{nn} \end{vmatrix}.$$

The  $i$ -th column of  $A$  is replaced by  $b$ .

# Cramer's Rule

## Theorem

*Let  $Ax = b$  be a linear system as before. If  $|A| \neq 0$  then we have*

$$x_i = \frac{|B_i|}{|A|}$$

*where  $B_i$  is the matrix created by replacing the  $i$ -th column of  $A$  by  $b$ .*

# Cramer's Rule

## Theorem

Let  $Ax = b$  be a linear system as before. If  $|A| \neq 0$  then we have

$$x_i = \frac{|B_i|}{|A|}$$

where  $B_i$  is the matrix created by replacing the  $i$ -th column of  $A$  by  $b$ .

This tells us an important thing: the solution to a linear system of equations can be expressed *algebraically* in terms of the coefficients of the system matrix and the right hand side using *continuous* operations.

## Summary Comments

- ▶ Inversion via cofactors/Cramer's rule is important because it tells us that the solution to a linear system can be expressed continuously from the components of the matrix and right hand side vector. This result is used for theoretical results in ODE, PDE, dynamical systems, chaos, etc.

## Summary Comments

- ▶ Inversion via cofactors/Cramer's rule is important because it tells us that the solution to a linear system can be expressed continuously from the components of the matrix and right hand side vector. This result is used for theoretical results in ODE, PDE, dynamical systems, chaos, etc.
- ▶ Both are very inefficient computations at size.

# Summary Comments

- ▶ Inversion via cofactors/Cramer's rule is important because it tells us that the solution to a linear system can be expressed continuously from the components of the matrix and right hand side vector. This result is used for theoretical results in ODE, PDE, dynamical systems, chaos, etc.
- ▶ Both are very inefficient computations at size.

Moral: Know both of these results, but never ever use them to do a computation.

# Summary Comments

- ▶ Inversion via cofactors/Cramer's rule is important because it tells us that the solution to a linear system can be expressed continuously from the components of the matrix and right hand side vector. This result is used for theoretical results in ODE, PDE, dynamical systems, chaos, etc.
- ▶ Both are very inefficient computations at size.

Moral: Know both of these results, but never ever use them to do a computation.

TODO:

- ▶ Prove that cofactor expansion is the determinant.



# Summary Comments

- ▶ Inversion via cofactors/Cramer's rule is important because it tells us that the solution to a linear system can be expressed continuously from the components of the matrix and right hand side vector. This result is used for theoretical results in ODE, PDE, dynamical systems, chaos, etc.
- ▶ Both are very inefficient computations at size.

Moral: Know both of these results, but never ever use them to do a computation.

TODO:

- ▶ Prove that cofactor expansion is the determinant.
- ▶ What does the determinant have to do with geometry?

# Summary Comments

- ▶ Inversion via cofactors/Cramer's rule is important because it tells us that the solution to a linear system can be expressed continuously from the components of the matrix and right hand side vector. This result is used for theoretical results in ODE, PDE, dynamical systems, chaos, etc.
- ▶ Both are very inefficient computations at size.

Moral: Know both of these results, but never ever use them to do a computation.

TODO:

- ▶ Prove that cofactor expansion is the determinant.
- ▶ What does the determinant have to do with geometry? (a bunch!)