

VS Properties

For all vectors u , v , and w in V and scalar values r and s in F we have

1. $u \oplus v = v \oplus u$
2. $u \oplus (v \oplus w) = (u \oplus v) \oplus w$
3. There is a vector $e \in V$ so that $u \oplus e = u$
4. There is a vector u^{-1} so that $u \oplus u^{-1} = e$.
5. $r \odot (u \oplus v) = (r \odot u) \oplus (r \odot v)$
6. $(r + s) \odot u = (r \odot u) \oplus (s \odot u)$
7. $r \odot (s \odot u) = (rs) \odot u$
8. $1 \odot u = u$

VS Properties

For all vectors u , v , and w in V and scalar values r and s in F we have

1. $u \oplus v = v \oplus u$
2. $u \oplus (v \oplus w) = (u \oplus v) \oplus w$
3. There is a vector $e \in V$ so that $u \oplus e = u$
4. There is a vector u^{-1} so that $u \oplus u^{-1} = e$.
5. $r \odot (u \oplus v) = (r \odot u) \oplus (r \odot v)$
6. $(r + s) \odot u = (r \odot u) \oplus (s \odot u)$
7. $r \odot (s \odot u) = (rs) \odot u$
8. $1 \odot u = u$

Examples of vector spaces are matrices, \mathbb{R}^n , functions, polynomials, series, and more.

Homogeneous Solution Set

Let $A \in \mathbb{R}^{n \times m}$. Consider the set

$$H = \{x \in \mathbb{R}^m : Ax = 0\}.$$

Homogeneous Solution Set

Let $A \in \mathbb{R}^{n \times m}$. Consider the set

$$H = \{x \in \mathbb{R}^m : Ax = 0\}.$$

This is the set of all solutions to a homogeneous system of linear equations.

Homogeneous Solution Set

Let $A \in \mathbb{R}^{n \times m}$. Consider the set

$$H = \{x \in \mathbb{R}^m : Ax = 0\}.$$

This is the set of all solutions to a homogeneous system of linear equations.

Homogeneous Solution Set

Let $A \in \mathbb{R}^{n \times m}$. Consider the set

$$H = \{x \in \mathbb{R}^m : Ax = 0\}.$$

This is the set of all solutions to a homogeneous system of linear equations. Note that H is not empty (since $0 \in H$).

Since $H \subset \mathbb{R}^m$ we have addition and scalar multiplication (already) defined for it.

Homogeneous Solution Set

Let $A \in \mathbb{R}^{n \times m}$. Consider the set

$$H = \{x \in \mathbb{R}^m : Ax = 0\}.$$

This is the set of all solutions to a homogeneous system of linear equations. Note that H is not empty (since $0 \in H$).

Since $H \subset \mathbb{R}^m$ we have addition and scalar multiplication (already) defined for it.

This means that all of the algebraic properties 1-8 hold for H assuming it is closed!

Set of Homogeneous Solutions is a Vector Space!

Set of Homogeneous Solutions is a Vector Space!

So:

If $x, y \in H$ then $A(x + y) = Ax + Ay = 0 + 0 = 0$.

Set of Homogeneous Solutions is a Vector Space!

So:

If $x, y \in H$ then $A(x + y) = Ax + Ay = 0 + 0 = 0$. Therefore $x + y \in H$ and H is closed under addition.

Set of Homogeneous Solutions is a Vector Space!

So:

If $x, y \in H$ then $A(x + y) = Ax + Ay = 0 + 0 = 0$. Therefore $x + y \in H$ and H is closed under addition.

If $x \in H$ and $r \in \mathbb{R}$ then $A(rx) = rAx = r0 = 0$.

Set of Homogeneous Solutions is a Vector Space!

So:

If $x, y \in H$ then $A(x + y) = Ax + Ay = 0 + 0 = 0$. Therefore $x + y \in H$ and H is closed under addition.

If $x \in H$ and $r \in \mathbb{R}$ then $A(rx) = rAx = r0 = 0$. Therefore $rx \in H$ and H is closed under scalar multiplication.

Set of Homogeneous Solutions is a Vector Space!

So:

If $x, y \in H$ then $A(x + y) = Ax + Ay = 0 + 0 = 0$. Therefore $x + y \in H$ and H is closed under addition.

If $x \in H$ and $r \in \mathbb{R}$ then $A(rx) = rAx = r0 = 0$. Therefore $rx \in H$ and H is closed under scalar multiplication.

In conclusion: H is a vector space.

Set of Homogeneous Solutions is a Vector Space!

So:

If $x, y \in H$ then $A(x + y) = Ax + Ay = 0 + 0 = 0$. Therefore $x + y \in H$ and H is closed under addition.

If $x \in H$ and $r \in \mathbb{R}$ then $A(rx) = rAx = r0 = 0$. Therefore $rx \in H$ and H is closed under scalar multiplication.

In conclusion: H is a vector space. It is also a subset of another vector space \mathbb{R}^m .

Set of Homogeneous Solutions is a Vector Space!

So:

If $x, y \in H$ then $A(x + y) = Ax + Ay = 0 + 0 = 0$. Therefore $x + y \in H$ and H is closed under addition.

If $x \in H$ and $r \in \mathbb{R}$ then $A(rx) = rAx = r0 = 0$. Therefore $rx \in H$ and H is closed under scalar multiplication.

In conclusion: H is a vector space. It is also a subset of another vector space \mathbb{R}^m . Such an object is called a **subspace**.

Subspace

Definition

Let V be a vector space and H a non-empty subset of V . If H is also a vector space under the vector operations of V then H is a **subspace** of V

Subspace

Definition

Let V be a vector space and H a non-empty subset of V . If H is also a vector space under the vector operations of V then H is a **subspace** of V

Theorem

If V is a vector space and H is a non-empty subset of V then H is a subspace if and only if it is closed under the operations of V .

Subspace

Definition

Let V be a vector space and H a non-empty subset of V . If H is also a vector space under the vector operations of V then H is a **subspace** of V

Theorem

If V is a vector space and H is a non-empty subset of V then H is a subspace if and only if it is closed under the operations of V .

Proof.

- The forward implication is clear.

Subspace

Definition

Let V be a vector space and H a non-empty subset of V . If H is also a vector space under the vector operations of V then H is a **subspace** of V

Theorem

If V is a vector space and H is a non-empty subset of V then H is a subspace if and only if it is closed under the operations of V .

Proof.

- ▶ The forward implication is clear.
- ▶ All of the “purely algebraic” properties (like associativity, distributivity, etc.) hold automatically.

Subspace

Definition

Let V be a vector space and H a non-empty subset of V . If H is also a vector space under the vector operations of V then H is a **subspace** of V

Theorem

If V is a vector space and H is a non-empty subset of V then H is a subspace if and only if it is closed under the operations of V .

Proof.

- ▶ The forward implication is clear.
- ▶ All of the “purely algebraic” properties (like associativity, distributivity, etc.) hold automatically.
- ▶ H is nonempty

Subspace

Definition

Let V be a vector space and H a non-empty subset of V . If H is also a vector space under the vector operations of V then H is a **subspace** of V

Theorem

If V is a vector space and H is a non-empty subset of V then H is a subspace if and only if it is closed under the operations of V .

Proof.

- ▶ The forward implication is clear.
- ▶ All of the “purely algebraic” properties (like associativity, distributivity, etc.) hold automatically.
- ▶ H is nonempty \Rightarrow There is at least one $v \in H$.

Subspace

Definition

Let V be a vector space and H a non-empty subset of V . If H is also a vector space under the vector operations of V then H is a **subspace** of V

Theorem

If V is a vector space and H is a non-empty subset of V then H is a subspace if and only if it is closed under the operations of V .

Proof.

- ▶ The forward implication is clear.
- ▶ All of the “purely algebraic” properties (like associativity, distributivity, etc.) hold automatically.
- ▶ H is nonempty \Rightarrow There is at least one $v \in H$. H closed $\Rightarrow 0v = 0 \in H$.

Subspace

Definition

Let V be a vector space and H a non-empty subset of V . If H is also a vector space under the vector operations of V then H is a **subspace** of V

Theorem

If V is a vector space and H is a non-empty subset of V then H is a subspace if and only if it is closed under the operations of V .

Proof.

- ▶ The forward implication is clear.
- ▶ All of the “purely algebraic” properties (like associativity, distributivity, etc.) hold automatically.
- ▶ H is nonempty \Rightarrow There is at least one $v \in H$. H closed $\Rightarrow 0v = 0 \in H$.
- ▶ H closed $\Rightarrow (-1)v = -v \in H$.

Examples of Subspaces and Not Subspaces

- ▶ As discussed before, the solution set to a homogeneous system of equations is a subspace.

Examples of Subspaces and Not Subspaces

- ▶ As discussed before, the solution set to a homogeneous system of equations is a subspace.
- ▶ Every vector space V has the smallest subspace $\{0\}$.

Examples of Subspaces and Not Subspaces

- ▶ As discussed before, the solution set to a homogeneous system of equations is a subspace.
- ▶ Every vector space V has the smallest subspace $\{0\}$. And biggest subspace V .

Examples of Subspaces and Not Subspaces

- ▶ As discussed before, the solution set to a homogeneous system of equations is a subspace.
- ▶ Every vector space V has the smallest subspace $\{0\}$. And biggest subspace V .
- ▶ Functions have polynomials as a subspace.

Examples of Subspaces and Not Subspaces

- ▶ As discussed before, the solution set to a homogeneous system of equations is a subspace.
- ▶ Every vector space V has the smallest subspace $\{0\}$. And biggest subspace V .
- ▶ Functions have polynomials as a subspace.
- ▶ $P_n(\mathbb{R})$ has $P_{n-1}(\mathbb{R})$ as a subspace.

Examples of Subspaces and Not Subspaces

- ▶ As discussed before, the solution set to a homogeneous system of equations is a subspace.
- ▶ Every vector space V has the smallest subspace $\{0\}$. And biggest subspace V .
- ▶ Functions have polynomials as a subspace.
- ▶ $P_n(\mathbb{R})$ has $P_{n-1}(\mathbb{R})$ as a subspace.
- ▶ The set of all polynomials of even degree is not a subspace.

Examples of Subspaces and Not Subspaces

- ▶ As discussed before, the solution set to a homogeneous system of equations is a subspace.
- ▶ Every vector space V has the smallest subspace $\{0\}$. And biggest subspace V .
- ▶ Functions have polynomials as a subspace.
- ▶ $P_n(\mathbb{R})$ has $P_{n-1}(\mathbb{R})$ as a subspace.
- ▶ The set of all polynomials of even degree is not a subspace. Since

$$(x^4 + 2x) \oplus (-x^4 + x^3) = x^3 + 2x$$

is of odd degree.

Examples of Subspaces and Not Subspaces

- ▶ As discussed before, the solution set to a homogeneous system of equations is a subspace.
- ▶ Every vector space V has the smallest subspace $\{0\}$. And biggest subspace V .
- ▶ Functions have polynomials as a subspace.
- ▶ $P_n(\mathbb{R})$ has $P_{n-1}(\mathbb{R})$ as a subspace.
- ▶ The set of all polynomials of even degree is not a subspace. Since

$$(x^4 + 2x) \oplus (-x^4 + x^3) = x^3 + 2x$$

is of odd degree. So the set is not closed under addition.

Examples of Subspaces and Not Subspaces

- ▶ As discussed before, the solution set to a homogeneous system of equations is a subspace.
- ▶ Every vector space V has the smallest subspace $\{0\}$. And biggest subspace V .
- ▶ Functions have polynomials as a subspace.
- ▶ $P_n(\mathbb{R})$ has $P_{n-1}(\mathbb{R})$ as a subspace.
- ▶ The set of all polynomials of even degree is not a subspace. Since

$$(x^4 + 2x) \oplus (-x^4 + x^3) = x^3 + 2x$$

is of odd degree. So the set is not closed under addition. (it is closed under multiplication though.)

Exercise

Which of the following sets W are/are not subspaces of \mathbb{R}^3 ? Why?

$$W_1 = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} : x = 0 \right\}$$

$$W_2 = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} : z = x + y \right\}$$

$$W_3 = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} : x \geq 0 \right\}$$

Exercise

Which of the following sets W are/are not subspaces of \mathbb{R}^3 ? Why?

$$W_1 = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} : x = 0 \right\}$$

$$W_2 = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} : z = x + y \right\}$$

$$W_3 = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} : x \geq 0 \right\}$$

W_1 and W_3 are. W_2 is not.

Linear Combinations and Span

Given a collection of vectors W in a vector space V , it is not necessarily true that W is always a subspace.

Linear Combinations and Span

Given a collection of vectors W in a vector space V , it is not necessarily true that W is always a subspace. But we can always create a (smallest) subspace of V which contains W .

Linear Combinations and Span

Given a collection of vectors W in a vector space V , it is not necessarily true that W is always a subspace. But we can always create a (smallest) subspace of V which contains W .

Definition

A **(finite) linear combination** of vectors from a vector space V is a sum of the form

$$\sum_{i=1}^n c_i v_i = c_1 v_1 + c_2 v_2 + \cdots + c_n v_n$$

where c_i are from the field and v_i are from V .

Linear Combinations and Span

Given a collection of vectors W in a vector space V , it is not necessarily true that W is always a subspace. But we can always create a (smallest) subspace of V which contains W .

Definition

A **(finite) linear combination** of vectors from a vector space V is a sum of the form

$$\sum_{i=1}^n c_i v_i = c_1 v_1 + c_2 v_2 + \cdots + c_n v_n$$

where c_i are from the field and v_i are from V .

Definition

Let V be a vector space and $W \subset V$. The **span** of W is the set

$$\text{span}(W) = \{v \in V : v \text{ is a linear combination of elements from } W\}.$$

$\text{span}(W)$ is a subspace

- ▶ $\text{span}(W)$ is not empty since $0 \in \text{span}(W)$.

$\text{span}(W)$ is a subspace

- ▶ $\text{span}(W)$ is not empty since $0 \in \text{span}(W)$.
- ▶ Closed under multiplication: Let $v \in \text{span}(W)$ and $r \in F$.

$\text{span}(W)$ is a subspace

- ▶ $\text{span}(W)$ is not empty since $0 \in \text{span}(W)$.
- ▶ Closed under multiplication: Let $v \in \text{span}(W)$ and $r \in F$.
Then

$$rv = r \sum_{i=1}^n c_i v_i = \sum_{i=1}^n (rc) v_i \in \text{span}(W).$$

$\text{span}(W)$ is a subspace

- ▶ $\text{span}(W)$ is not empty since $0 \in \text{span}(W)$.
- ▶ Closed under multiplication: Let $v \in \text{span}(W)$ and $r \in F$. Then

$$rv = r \sum_{i=1}^n c_i v_i = \sum_{i=1}^n (rc) v_i \in \text{span}(W).$$

- ▶ Closed under addition: Let $v, w \in \text{span}(W)$. Without loss of generality we can assume that

$$v = \sum_{i=1}^n a_i v_i \text{ and } w = \sum_{i=1}^n b_i v_i.$$

$\text{span}(W)$ is a subspace

- ▶ $\text{span}(W)$ is not empty since $0 \in \text{span}(W)$.
- ▶ Closed under multiplication: Let $v \in \text{span}(W)$ and $r \in F$. Then

$$rv = r \sum_{i=1}^n c_i v_i = \sum_{i=1}^n (rc) v_i \in \text{span}(W).$$

- ▶ Closed under addition: Let $v, w \in \text{span}(W)$. Without loss of generality we can assume that

$$v = \sum_{i=1}^n a_i v_i \text{ and } w = \sum_{i=1}^n b_i v_i.$$

$$\text{So } v + w = \sum_{i=1}^n (a_i + b_i) v_i \in \text{span}(W)$$