To call in the statistician after the experiment is done may be no more than asking him to perform a post-mortem examination: he may be able to say what the experiment died of.

Sir Ronald Fisher

Presidential Address to the First Indian Statistical Congress, 1938. Sankhya 4, 14-17

(BUT maybe you don't need to call in a statistician if you know some principles)

When assumptions are not met

Timothée Bonnet

Biological Data Science Institute

May 2, 2019

If you haven't enough R in your life

https://www.meetup.com/rladies-canberra/events/dvrjwqyzhbjb/



Monday, May 6, 2019 Tidying and plotting in R








Details

R is primarily known as a tool for statistics and data analysis, but it is also great for "simpler" data wrangling and plotting purposes. As R is becoming more accessible and useful in this regards, it is becoming a more viable alternative to Microsoft Excel and other spreadsheet-based applications for wrangling,



Assumptions are about math properties, but also about scientific logic

- Assumptions are about math properties, but also about scientific logic
- In general assumptions are not perfectly met

- Assumptions are about math properties, but also about scientific logic
- 2 In general assumptions are not perfectly met
- There are many technics to get closer

- Assumptions are about math properties, but also about scientific logic
- In general assumptions are not perfectly met
- There are many technics to get closer
- "there is no free lunch", every model makes assumptions, explicit or implicit

- Assumptions are about math properties, but also about scientific logic
- 2 In general assumptions are not perfectly met
- There are many technics to get closer
- "there is no free lunch", every model makes assumptions, explicit or implicit
- 4 All models are wrong, a few are useful


- Why we need assumptions
- 2 What can go wrong; e.g. Linear models


1. Scientific question


1. Scientific question





2. Model and Statistical question












- Why we need assumptions
- 2 What can go wrong; e.g. Linear models

Simple linear model

$Response = Intercept + Slope \times Predictor + Error$



Linear model assumptions

What are linear model assumptions?

9/14

Linear model assumptions

Assumptions do NOT include:

- Relationship is a straight line ("Linear" means a line on some scale, not any scale)
- Data normality (Only error normality)
- Collinearity (Changes parameter meaning, not the validity)

Linear model assumptions

Linear combination of parameters (including transformation, polynoms, interactions...)
 Risk: biologically meaningless

Predictor not perfectly correlated

Risk: Model won't run, unstable convergence, or huge SE

• Little error in predictors

Risk: bias estimates (underestimate with Gaussian error)

Gaussian error distribution
 Risk: Poor predictions, wrong uncertainty

Homoscedasticity (constant error variance)
 Risk: Poor predictions, wrong uncertainty

Independence of error
 Risk: Bias and over-optimistic uncertainty

Residual distributions

Exercise 1

Residual distributions

Exercise 1

Exercise 2

Non-independence

Exercise 3

Non-independence

Exercise 3

Exercise 4

11 / 14

Let's think about more complex issues

Non-Linear relationships / Thinking
 Solution: do you really need that? / non-linear models

- Non-Linear relationships / Thinking
 Solution: do you really need that? / non-linear models
- Predictor perfectly correlated / Singular fit Solution: Drop redundant variable

- Non-Linear relationships / Thinking
 Solution: do you really need that? / non-linear models
- Predictor perfectly correlated / Singular fit Solution: Drop redundant variable
- Error in predictors / Thinking / multiple measurements
 Solution: Better experiment / measurement error model

- Non-Linear relationships / Thinking
 Solution: do you really need that? / non-linear models
- Predictor perfectly correlated / Singular fit Solution: Drop redundant variable
- Error in predictors / Thinking / multiple measurements
 Solution: Better experiment / measurement error model
- Non-Gaussian error distribution / Plot / experimental design Solution: Transformation / GLM

- Non-Linear relationships / Thinking
 Solution: do you really need that? / non-linear models
- Predictor perfectly correlated / Singular fit Solution: Drop redundant variable
- Error in predictors / Thinking / multiple measurements
 Solution: Better experiment / measurement error model
- Non-Gaussian error distribution / Plot / experimental design Solution: Transformation / GLM
- Changing error variance Plot / experimental design Solution: Transformation / GLM

- Non-Linear relationships / Thinking
 Solution: do you really need that? / non-linear models
- Predictor perfectly correlated / Singular fit Solution: Drop redundant variable
- Error in predictors / Thinking / multiple measurements
 Solution: Better experiment / measurement error model
- Non-Gaussian error distribution / Plot / experimental design Solution: Transformation / GLM
- Changing error variance Plot / experimental design Solution: Transformation / GLM
- Correlated errors Thinking (/plot)
 Solution: Experimental design / control variables / mixed models

- Assumptions are about math properties, but also about scientific logic
- In general assumptions are not perfectly met
- There are many technics to get closer
- "there is no free lunch", every model makes assumptions, explicit or implicit
- 4 All models are wrong, a few are useful