## 1

## Assignment 8

## Ananthoju Pranav Sai - AI20BTECH11004

Download the latex code from

https://github.com/Ananthoju-Pranav-Sai/AI1103/blob/main/Assignment 8/main.tex

UGC June 2017 Math set A Q 57

Suppose  $(X_1, X_2)$  follows a bivariate normal distribution with

$$E(X_1) = E(X_2) = 0$$
 (0.0.1)

$$V(X_1) = V(X_2) = 2 (0.0.2)$$

$$Cov(X_1, X_2) = -1$$
 (0.0.3)

If  $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} e^{-\frac{y^2}{2}} dy$ , then  $\Pr(X_1 - X_2 > 6) = ?$ 

- 1)  $\Phi(-1)$
- 2)  $\Phi(-3)$
- 3)  $\Phi(\sqrt{6})$
- 4)  $\Phi(-\sqrt{6})$

## Solution

Let  $\mathbf{x} \sim \mathcal{N}(\mu_{\mathbf{x}}, \Sigma_{\mathbf{x}})$  be a bivariate random vector.

$$\mathbf{x} = \begin{pmatrix} X_1 \\ X_2 \end{pmatrix} \tag{0.0.4}$$

$$\mu_{\mathbf{x}} = \begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix} \tag{0.0.5}$$

$$\Sigma_{\mathbf{x}} = \begin{pmatrix} \sigma_1^2 & \rho \sigma_1 \sigma_2 \\ \rho \sigma_1 \sigma_2 & \sigma_2^2 \end{pmatrix} \tag{0.0.6}$$

where  $\Sigma_{\mathbf{x}}$  is covariance matrix of  $\mathbf{x}$  and  $\rho$  is correlation of  $X_1$  and  $X_2$  which is given by

$$\rho = \frac{Cov(X_1, X_2)}{\sigma_1 \sigma_2} \tag{0.0.7}$$

**Theorem 0.1.** If  $(X_1, X_2)$  follow bivariate distribution then  $Pr(X_1 - X_2 > \alpha)$  is given by

$$\Phi\left(\frac{(\mu_1-\mu_2)-\alpha}{\sqrt{\sigma_1^2+\sigma_2^2-2\rho\sigma_1\sigma_2}}\right).$$

*Proof.* Let 
$$\mathbf{u} = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$
 and  $\mathbf{x} = \begin{pmatrix} X_1 \\ X_2 \end{pmatrix}$ 
$$X_2 - X_1 = \mathbf{u}^{\mathsf{T}} \mathbf{x} \tag{0.0.8}$$

Now consider a random variable Y defined as follows

$$Y = X_2 - X_1 \tag{0.0.9}$$

$$\implies \mathbf{y} = \mathbf{u}^{\mathsf{T}} \mathbf{x} \tag{0.0.10}$$

then y has normal distribution with mean

$$\mu_{\mathbf{v}} = \mathbf{u}^{\mathsf{T}} \mu_{\mathbf{x}} \tag{0.0.11}$$

$$\implies \mu_{\mathbf{y}} = \left(-1 \ 1\right) \begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix} \tag{0.0.12}$$

$$\implies \mu_{\mathbf{y}} = \left(\mu_2 - \mu_1\right) \tag{0.0.13}$$

and covariance matrix is given by

$$\Sigma_{\mathbf{v}} = \mathbf{u}^{\mathsf{T}} \Sigma_{\mathbf{x}} (\mathbf{u}^{\mathsf{T}})^{\mathsf{T}}$$
 (0.0.14)

$$\implies \Sigma_{\mathbf{y}} = \begin{pmatrix} -1 & 1 \end{pmatrix} \begin{pmatrix} \sigma_1^2 & \rho \sigma_1 \sigma_2 \\ \rho \sigma_1 \sigma_2 & \sigma_2^2 \end{pmatrix} \begin{pmatrix} -1 \\ 1 \end{pmatrix} \quad (0.0.15)$$

$$\implies \Sigma_{\mathbf{y}} = \left(\sigma_1^2 + \sigma_2^2 - 2\rho\sigma_1\sigma_2\right) \tag{0.0.16}$$

$$\therefore Y \sim \mathcal{N}(\mu = \mu_2 - \mu_1, \sigma^2 = \sigma_1^2 + \sigma_2^2 - 2\rho\sigma_1\sigma_2)$$

The Standard Normal, often written Z, is a Normal with  $\mu = 0$  and  $\sigma^2 = 1$ . Thus,  $Z \sim \mathcal{N}(\mu = 0, \sigma^2 = 1)$ 

Now  $Pr(X_1 - X_2 > \alpha)$  can be written as  $Pr(Y < -\alpha)$ 

$$\Pr(Y < -\alpha) = \Pr\left(\frac{Y - \mu_y}{\sigma_y} < \frac{-\alpha - \mu_y}{\sigma_y}\right)$$

(0.0.17)

$$\Rightarrow \Pr(Y < -\alpha) = \Pr\left(Z < \frac{-\alpha - (\mu_2 - \mu_1)}{\sqrt{\sigma_1^2 + \sigma_2^2 - 2\rho\sigma_1\sigma_2}}\right)$$

$$(0.0.18)$$

$$\therefore \Pr(X_1 - X_2 > \alpha) = \Phi\left(\frac{(\mu_1 - \mu_2) - \alpha}{\sqrt{\sigma_1^2 + \sigma_2^2 - 2\rho\sigma_1\sigma_2}}\right)$$
(0.0.19)

Given  $(X_1, X_2)$  follow a bivariate normal distribution with

$$\mu_1 = \mu_2 = 0 \tag{0.0.20}$$

$$\sigma_1^2 = \sigma_2^2 = 2 \tag{0.0.21}$$

$$Cov(X_1, X_2) = -1$$
 (0.0.22)

$$\rho = \frac{Cov(X_1, X_2)}{\sigma_1 \sigma_2} = \frac{-1}{2}$$
 (0.0.23)

for  $Pr(X_1 - X_2 > 6)$  we can use the above theorem as follows

$$\Pr(X_1 - X_2 > \alpha) = \Phi\left(\frac{(\mu_1 - \mu_2) - \alpha}{\sqrt{\sigma_1^2 + \sigma_2^2 - 2\rho\sigma_1\sigma_2}}\right)$$
(0.0.24)

$$\implies \Pr(X_1 - X_2 > 6) = \Phi\left(\frac{(0-0) - 6}{\sqrt{2 + 2 - 2\left(\frac{-1}{2}\right)2}}\right)$$
(0.0.25)

$$\implies \Pr(X_1 - X_2 > 6) = \Phi\left(\frac{-6}{\sqrt{6}}\right)$$
 (0.0.26)

$$\therefore \Pr(X_1 - X_2 > 6) = \Phi(-\sqrt{6}) \tag{0.0.27}$$