

ASSIGNMENT 5

Ananthoju Pranav Sai
AI20BTECH11004

Download all python codes from

<https://github.com/Ananthoju-Pranav-Sai/EE3900/blob/main/Assignment-5/codes/Assignment-5.py>

and latex-tikz codes from

<https://github.com/Ananthoju-Pranav-Sai/EE3900/tree/main/Assignment-5/Assignment-5.tex>

1 QUADRATIC FORMS Q.2.62

Find a point on the curve $y = (x - 2)^2$ at which the tangent is parallel to the chord joining the points $\begin{pmatrix} 2 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 4 \\ 4 \end{pmatrix}$.

2 SOLUTION

Equation of the given conic in vector form is

$$\mathbf{x}^T \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \mathbf{x} + 2 \begin{pmatrix} -2 & -\frac{1}{2} \end{pmatrix} \mathbf{x} + 4 = 0 \quad (2.0.1)$$

Therefore we have

$$\mathbf{V} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad (2.0.2)$$

$$\mathbf{u} = \begin{pmatrix} -2 \\ -\frac{1}{2} \end{pmatrix} \quad (2.0.3)$$

Given the tangent is parallel to the chord joining the points $\begin{pmatrix} 2 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 4 \\ 4 \end{pmatrix}$ then the tangent is given by

$$\mathbf{x} = \mathbf{q} + \mu \mathbf{m} \quad (2.0.4)$$

where,

$$\mathbf{m} = \begin{pmatrix} 4 \\ 4 \end{pmatrix} - \begin{pmatrix} 2 \\ 0 \end{pmatrix} \quad (2.0.5)$$

$$\mathbf{m} = \begin{pmatrix} 2 \\ 4 \end{pmatrix} \quad (2.0.6)$$

and \mathbf{q} is the point of contact of tangent on the curve.

Lemma 2.1. *If the line*

$$L : \mathbf{x} = \mathbf{q} + \mu \mathbf{m} \quad (2.0.7)$$

is tangent to the conic

$$\mathbf{x}^T \mathbf{V} \mathbf{x} + 2 \mathbf{u}^T \mathbf{x} + f = 0 \quad (2.0.8)$$

then

$$\mathbf{m}^T (\mathbf{V} \mathbf{q} + \mathbf{u}) = 0 \quad (2.0.9)$$

So from lemma we have

$$\begin{pmatrix} 2 & 4 \end{pmatrix} \left(\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \mathbf{q} + \begin{pmatrix} -2 \\ -\frac{1}{2} \end{pmatrix} \right) = 0 \quad (2.0.10)$$

Let $\mathbf{q} = \begin{pmatrix} q_1 \\ q_2 \end{pmatrix}$ then

$$\begin{pmatrix} 2 & 4 \end{pmatrix} \begin{pmatrix} q_1 - 2 \\ -\frac{1}{2} \end{pmatrix} = 0 \quad (2.0.11)$$

$$q_1 = 3 \quad (2.0.12)$$

As \mathbf{q} also lies on conic from (2.0.1) we get

$$\mathbf{q} = \begin{pmatrix} 3 \\ 1 \end{pmatrix} \quad (2.0.13)$$




Fig. 0: Plot of the tangent and parabola