



OPEN ACCESS

EDITED AND REVIEWED BY

Wei Peng,
Chengdu University of Traditional Chinese
Medicine, China

*CORRESPONDENCE

Peibo Li,
✉ lipeibo@mail.sysu.edu.cn

RECEIVED 27 March 2026

ACCEPTED 02 April 2026

PUBLISHED 21 April 2026

CITATION

Xu Z, Wu H, Fan W, Meng F, Hu S, Zou M,
Chen Y, Su W and Li P (2026) Correction:
Naoxintong capsule decreases circulating
exosomes of miR-382-5p to protect LPS-
induced vascular endothelial cell injury by
targeting STC1 *in vitro*.

Front. Pharmacol. 17:1827794.

doi: 10.3389/fphar.2026.1827794

COPYRIGHT

© 2026 Xu, Wu, Fan, Meng, Hu, Zou, Chen,
Su and Li. This is an open-access article
distributed under the terms of the [Creative
Commons Attribution License \(CC BY\)](#).
The use, distribution or reproduction in
other forums is permitted, provided the
original author(s) and the copyright
owner(s) are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does not
comply with these terms.

Correction: Naoxintong capsule decreases circulating exosomes of miR-382-5p to protect LPS-induced vascular endothelial cell injury by targeting STC1 *in vitro*

Ziyan Xu, Hao Wu, Weiyang Fan, Fenzhao Meng, Shuangfei Hu,
Muhan Zou, Yixuan Chen, Weiwei Su and Peibo Li*

Guangdong Engineering and Technology Research Center for Quality and Efficacy Re-Evaluation of Post-Marketed TCM, Guangdong Provincial Key Laboratory of Plant Stress Biology, State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-sen University, Guangzhou, China

KEYWORDS

circulating exosomes, miR-382-5p, naoxintong capsule (NXT), protective effect, stanniocalcin-1 (STC1), TLR4/TRIF/NF- κ B pathway, vascular endothelial injury

A Correction on

Naoxintong capsule decreases circulating exosomes of miR-382-5p to protect LPS-induced vascular endothelial cell injury by targeting STC1 *in vitro*

by Xu Z, Wu H, Fan W, Meng F, Hu S, Zou M, Chen Y, Su W and Li P (2026). *Front. Pharmacol.* 17: 1655883. doi: 10.3389/fphar.2026.1655883

In the original publication, there were a few mistakes in [Figures 1, 2, 7, 8](#). In [Figure 1D](#), the kDa value for the protein ALIX was incorrectly stated as “80” instead of the correct “96”. In [Figure 2G](#), the kDa value for the protein Cleaved caspase 3 was incorrectly stated as “23” instead of the correct “17”. In [Figure 7I](#), the kDa value for the protein Cleaved caspase 3 was incorrectly stated as “16” instead of the correct “17”. In [Figure 8H](#), the kDa value for the protein Cleaved caspase 3 was incorrectly stated as “23” instead of the correct “17”. The corrected [Figures 1, 2, 7, 8](#) appear below.

The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.





FIGURE 1
Body weight map and biological identification, concentration determination, and internalization of circulating exosomes. **(A)** Daily body weight of SD rats during the experiment. **(B)** Circulating exosomes scanning results of TEM (Scale bar, 50 nm). **(C)** The particle diameter distribution of circulating exosomes tested by NTA. **(D)** The circulating exosome markers (ALIX, TSG101, CD9) and the negative control marker (β -actin) detected by Western blot. **(E)** The concentrations of circulating exosomes tested by BCA. **(F)** The internalization of circulating exosomes from SD rats in HMEC-1 imaged by confocal microscopy (Scale bar, 10 μ m). PKH-26 (Ex: 551 nm, Em: 562~580 nm) labeled circulating exosomes, Dio (Ex: 484 nm, Em: 496~520 nm) labeled cell membrane, and Hoechst 33342 (Ex: 346 nm, Em: 455~470 nm) labeled cell nuclei. Values are expressed as mean $\pm$ SD, n = 3. * p < 0.05, ** p < 0.01 and ns indicated no significant statistical difference.






FIGURE 7

miR-382-5p downregulated the expression of STC1 and the effect of STC1 on the endothelial protection of NXT-Exo. (A) The mRNA expression level of STC1 in HMEC-1 transfected with miR-382-5p mimic or miR-382-5p mimic-NC was detected by RT-qPCR. (B) The mRNA expression level of STC1 in HMEC-1 treated with NXT-Exo or NXT-Exo combined with miR-382-5p was detected by RT-qPCR. (C) The transfection of si-STC1 was verified by RT-qPCR. (D,E) The secretion levels of IL-6 and IL-8 were measured by ELISA. (F) The release of LDH was detected by microplate method. (G,H) The production level of ROS and expression level of MDA were tested by fluorescence. (I) The effects of miR-382-5p and STC1 treatment on the relative protein expression levels of Bax, Bcl-2, Cleaved caspase 3 and STC1 in HMEC-1 were detected by Western blot. Values are expressed as mean $\pm$ SD, $n = 3$. * $p < 0.05$, ** $p < 0.01$.

