

18.404/6.840 Lecture 16

Last time:

- NP-completeness
- $3SAT \leq_p CLIQUE$
- $3SAT \leq_p HAMPATH$

Today:

- Cook-Levin Theorem: SAT is NP-complete
- $3SAT$ is NP-complete

Quick Review

Defn: B is NP-complete if

- 1) $B \in \text{NP}$
- 2) For all $A \in \text{NP}$, $A \leq_P B$

If B is NP-complete and $B \in \text{P}$ then $\text{P} = \text{NP}$.

Importance of NP-completeness

- 1) Evidence of computational intractability.
- 2) Gives a good candidate for proving $\text{P} \neq \text{NP}$.

To show some language C is NP-complete, show $3\text{SAT} \leq_P C$.

— or some other previously shown NP-complete language

Check-in 16.1

The big sigma notation means summing over some set.

$$\sum_{1 \leq i \leq n} i = 1 + 2 + \dots + n$$

The big AND (or OR) notation has a similar meaning.

For example, if $x = x_1 \dots x_n$ and $y = y_1 \dots y_n$ are two strings of length n , when does the following hold?

$$\left(\bigwedge_{1 \leq i \leq n} x_i = y_i \right) = \text{TRUE}$$

- (a) Whenever x and y agree on some symbol.
- (b) Whenever $x = y$.

Cook-Levin Theorem (idea)

Theorem: SAT is NP-complete

Proof: 1) $SAT \in NP$ (done)

2) Show that for each $A \in NP$ we have $A \leq_p SAT$:

Let $A \in NP$ be decided by NTM M in time n^k .

Give a polynomial-time reduction f mapping A to SAT .

$f: \Sigma^* \rightarrow \text{formulas}$

$f(w) = \langle \phi_{M,w} \rangle$


$w \in A$ iff $\phi_{M,w}$ is satisfiable

Idea: $\phi_{M,w}$ simulates M on w . Design $\phi_{M,w}$ to “say” M accepts w .

Satisfying assignment to $\phi_{M,w}$ is a computation history for M on w .

Tableau for M on w

Defn: An (accepting) tableau for NTM M on w is an $n^k \times n^k$ table representing an computation history for M on w on an accepting branch of the nondeterministic computation.




Construct $\phi_{M,w}$ to "say" M accepts w .

$\phi_{M,w}$ "says" a tableau for M on w exists.

$$\phi_{M,w} = \phi_{\text{cell}} \wedge \phi_{\text{start}} \wedge \phi_{\text{move}} \wedge \phi_{\text{accept}}$$

Constructing $\phi_{M,w}$: ϕ_{cell}



The variables of $\phi_{M,w}$ are $x_{i,j,\sigma}$ for $1 \leq i, j \leq n^k$ and $\sigma \in \Gamma \cup Q$.

$x_{i,j,\sigma} = \text{TRUE}$ means cell i, j contains σ .

Check-in 16.2

How many variables does $\phi_{M,w}$ have?

Recall that $n = |w|$.

- (a) $O(n)$
- (b) $O(n^2)$
- (c) $O(n^k)$
- (d) $O(n^{2k})$

Constructing $\phi_{M,w}$: ϕ_{start} and ϕ_{accept}

	1	2	3	n^k		
1	q_0	w_1	w_2	w_3	\cdots	w_n	$\sqcup \cdots \sqcup$	← Start configuration
	a	q_7	w_2	\cdots				
n^k				\cdots	q_{accept}	\cdots		← Accepting configuration

$\phi_{M,w}$ “says” a tableau for M on w exists.

$$\phi_{M,w} = \phi_{\text{cell}} \wedge \phi_{\text{start}} \wedge \phi_{\text{move}} \wedge \phi_{\text{accept}}$$


ϕ_{cell} done ✓

$$\phi_{\text{start}} =$$

$$\phi_{\text{accept}} = \bigvee_{1 \leq j \leq n^k} x_{n^k, j, q_{\text{accept}}}$$



Constructing $\phi_{M,w}$: ϕ_{move}

	j							
q_0	w_1	w_2	w_3	\cdots	w_n	\sqsubset	\cdots	\sqsubset
a	q_7	w_2	\cdots					
i								
			\cdots	q_{accept}	\cdots			

2 × 3 neighborhood

$\phi_{M,w}$ “says” a tableau for M on w exists.

$$\phi_{M,w} = \underbrace{\phi_{\text{cell}}}_{\checkmark} \wedge \underbrace{\phi_{\text{start}}}_{\checkmark} \wedge \phi_{\text{move}} \wedge \underbrace{\phi_{\text{accept}}}_{\checkmark}$$

Legal neighborhoods: consistent with M 's transition function

potential
examples:

a	q_7	b
q_3	a	c

a	b	c
a	b	c

a	b	c
a	b	q_5

a	b	c
d	b	c

Illegal neighborhoods: not consistent with M 's transition function

examples:

a	b	c
a	d	c

a	b	c
a	q_2	c

a	q_7	c
a	b	c

a	q_7	c
q_3	d	q_4

Claim: If every 2×3 neighborhood is legal then tableau corresponds to a computation history.

$$\phi_{\text{move}} = \bigwedge_{1 \leq i, j \leq n^k} \left(\bigvee_{\substack{\text{Legal} \\ \begin{array}{|c|c|c|} \hline r & s & t \\ \hline v & y & z \\ \hline \end{array}}} (x_{i,j-1,r} \wedge x_{i,j,s} \wedge x_{i,j+1,t} \wedge x_{i+1,j-1,v} \wedge x_{i+1,j,y} \wedge x_{i+1,j+1,z}) \right)$$

Conclusion: SAT is NP-complete

Diagram illustrating a matrix structure, likely representing a quantum circuit or a data layout. The matrix is labeled with dimensions n^k on both the horizontal and vertical axes.

The matrix is divided into four quadrants by a vertical and a horizontal line. The top-left quadrant contains the labels q_0 , w_1 , w_2 , w_3 , \dots , w_n , \dots . The top-right quadrant contains the label \dots . The bottom-left quadrant contains the label a , q_7 , w_2 , \dots . The bottom-right quadrant contains the label \dots , q_{accept} , \dots .

Summary:

For $A \in \text{NP}$, decided by NTM M ,
we gave a reduction f from A to SAT :

$$f: \Sigma^* \rightarrow \text{formulas}$$
$$f(w) = \langle \phi_{M,w} \rangle$$
$$w \in A \text{ iff } \phi_{M,w} \text{ is satisfiable.}$$

$$\phi_{M,w} = \phi_{\text{cell}} \wedge \phi_{\text{start}} \wedge \phi_{\text{move}} \wedge \phi_{\text{accept}}$$

The size of $\phi_{M,w}$ is roughly the size of the tableau for M on w , so size is $O(n^k \times n^k) = O(n^{2k})$.

Therefore f is computable in polynomial time.

3SAT is NP-complete

a	b	$a \vee b = c$	
1	1	1	$(a \wedge b) \rightarrow c$
0	1	1	$(\bar{a} \wedge b) \rightarrow c$
1	0	1	$(a \wedge \bar{b}) \rightarrow c$
0	0	0	$(\bar{a} \wedge \bar{b}) \rightarrow \bar{c}$

Theorem: 3SAT is NP-complete


Proof: Show $SAT \leq_p 3SAT$

Give reduction f converting formula ϕ to 3CNF formula ϕ' , preserving satisfiability.

(Note: ϕ and ϕ' are not logically equivalent)

Example: Say $\phi = ((a \wedge b) \vee c) \wedge (\bar{a} \vee b)$

Tree structure for ϕ :



Logical equivalence: $(A \rightarrow B)$ and $(\bar{A} \vee B)$ $\overline{(A \wedge B)}$ and $(\bar{A} \vee \bar{B})$

$$\begin{aligned} \phi' = & ((a \wedge b) \rightarrow z_1) \wedge ((\bar{a} \wedge b) \rightarrow \bar{z}_1) \wedge ((a \wedge \bar{b}) \rightarrow \bar{z}_1) \wedge ((\bar{a} \wedge \bar{b}) \rightarrow \bar{z}_1) \\ & \wedge ((z_1 \wedge c) \rightarrow z_2) \wedge ((\bar{z}_1 \wedge c) \rightarrow z_2) \wedge ((z_1 \wedge \bar{c}) \rightarrow z_2) \wedge ((\bar{z}_1 \wedge \bar{c}) \rightarrow \bar{z}_2) \\ & \vdots \text{ repeat for each } z_i \end{aligned}$$

$\wedge (z_4)$

Check-in 16.3

If ϕ has k operations (\wedge and \vee), how many clauses has ϕ' ?

- | | |
|--------------|------------|
| (a) $k + 1$ | (c) k^2 |
| (b) $4k + 1$ | (d) $2k^2$ |

Quick review of today

1. SAT is NP-complete
2. $3SAT$ is NP-complete