

18.404/6.840 Lecture 9

Last time:

- A_{TM} is undecidable
- The diagonalization method
- $\overline{A_{TM}}$ is T-unrecognizable
- The Reducibility Method, preview

Today:

- The Reducibility Method for proving undecidability and T-unrecognizability.
- General reducibility
- Mapping reducibility

Posted: Problem Set 2 solutions and Problem Set 3.

TAs are available to answer questions during chat-breaks!

The Reducibility Method

If we know that some problem (say A_{TM}) is undecidable, we can use that to show other problems are undecidable.

Defn: $HALT_{\text{TM}} = \{\langle M, w \rangle \mid M \text{ halts on input } w\}$

Recall Theorem: $HALT_{\text{TM}}$ is undecidable

Proof by contradiction, showing that A_{TM} is reducible to $HALT_{\text{TM}}$:

Assume that $HALT_{\text{TM}}$ is decidable and show that A_{TM} is decidable (false!).

Let TM R decide $HALT_{\text{TM}}$.

Construct TM S deciding A_{TM} .

$S =$ “On input $\langle M, w \rangle$

1. Use R to test if M on w halts. If not, *reject*.
2. Simulate M on w until it halts (as guaranteed by R).
3. If M has accepted then *accept*.
If M has rejected then *reject*.

TM S decides A_{TM} , a contradiction. Therefore $HALT_{\text{TM}}$ is undecidable.

Reducibility – Concept

If we have two languages (or problems) A and B , then A is reducible to B means that we can use B to solve A .

Example 1: Measuring the area of a rectangle is reducible to measuring the lengths of its sides.

Example 2: We showed that A_{NFA} is reducible to A_{DFA} .

Example 3: From Pset 2, $PUSHER$ is reducible to E_{CFG} .
(Idea- Convert push states to accept states.)

If A is reducible to B then solving B gives a solution to A .

- then B is easy $\rightarrow A$ is easy.

- then A is hard $\rightarrow B$ is hard.

this is the form we will use

Check-in 9.1

Is Biology reducible to Physics?

- (a) Yes, all aspects of the physical world may be explained in terms of Physics, at least in principle.
- (b) No, some things in the world, maybe life, the brain, or consciousness, are beyond the realm of Physics.
- (c) I'm on the fence on this question!

E_{TM} is undecidable

Let $E_{\text{TM}} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) = \emptyset\}$

Theorem: E_{TM} is undecidable

Proof by contradiction. Show that A_{TM} is reducible to E_{TM} .

Assume that E_{TM} is decidable and show that A_{TM} is decidable (false!).

Let TM R decide E_{TM} .

Construct TM S deciding A_{TM} .

$S =$ "On input $\langle M, w \rangle$

1. Transform M to new TM $M_w =$ "On input x
 1. If $x \neq w$, *reject*.
 2. else run M on w
 3. *Accept* if M accepts."
2. Use R to test whether $L(M_w) = \emptyset$
3. If YES [so M rejects w] then *reject*.
If NO [so M accepts w] then *accept*.


M_w works like M except that it always rejects strings x where $x \neq w$.

So $L(M_w) = \begin{cases} \{w\} & \text{if } M \text{ accepts } w \\ \emptyset & \text{if } M \text{ rejects } w \end{cases}$

Mapping Reducibility

Defn: Function $f: \Sigma^* \rightarrow \Sigma^*$ is computable if there is a TM F where F on input w halts with $f(w)$ on its tape, for all strings w .

Defn: A is mapping-reducible to B ($A \leq_m B$) if there is a computable function f where $w \in A$ iff $f(w) \in B$.



Example: $A_{\text{TM}} \leq_m \overline{E_{\text{TM}}}$

The computable reduction function f is $f(\langle M, w \rangle) = \langle M_w \rangle$

Because $\langle M, w \rangle \in A_{\text{TM}}$ iff $\langle M_w \rangle \in \overline{E_{\text{TM}}}$
 (M accepts w iff $L(\langle M_w \rangle) \neq \emptyset$)

Recall TM $M_w =$ "On input x

1. If $x \neq w$, *reject*.
2. else run M on w
3. *Accept* if M accepts."

Mapping Reductions - properties


Theorem: If $A \leq_m B$ and B is decidable then so is A

Proof: Say TM R decides B .

Construct TM S deciding A :

$S =$ "On input w

1. Compute $f(w)$
2. Run R on $f(w)$ to test if $f(w) \in B$
3. If R halts then output same result."



Corollary: If $A \leq_m B$ and A is undecidable then so is B

Theorem: If $A \leq_m B$ and B is T-recognizable then so is A

Proof: Same as above.

Corollary: If $A \leq_m B$ and A is T-unrecognizable then so is B

Check-in 9.2

Suppose $A \leq_m B$.

What can we conclude?

Check all that apply.


- (a) $B \leq_m A$
- (b) $\bar{A} \leq_m \bar{B}$
- (c) None of the above



Mapping vs General Reducibility


Mapping Reducibility of A to B : Translate A -questions to B -questions.

- A special type of reducibility
- Useful to prove T-unrecognizability



(General) Reducibility of A to B : Use B solver to solve A .

- May be conceptually simpler
- Useful to prove undecidability



Noteworthy difference:

- A is reducible to \overline{A}
- A may not be mapping reducible to \overline{A} .

For example $\overline{A_{TM}} \not\leq_m A_{TM}$

Check-in 9.3

We showed that if $A \leq_m B$ and B is T-recognizable then so is A .

Is the same true if we use general reducibility instead of mapping reducibility?

- (a) Yes
- (b) No

Check-in 9.3

Reducibility – Templates

To prove B is undecidable:

- Show undecidable A is reducible to B . (often A is A_{TM})
- Template: Assume TM R decides B .
Construct TM S deciding A . Contradiction.

To prove B is T-unrecognizable:

- Show T-unrecognizable A is mapping reducible to B . (often A is $\overline{A_{\text{TM}}}$)
- Template: give reduction function f .

E_{TM} is T-unrecognizable

Recall $E_{\text{TM}} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) = \emptyset\}$

Theorem: E_{TM} is T-unrecognizable


Proof: Show $\overline{A_{\text{TM}}} \leq_m E_{\text{TM}}$

Reduction function: $f(\langle M, w \rangle) = \langle M_w \rangle$ Recall TM $M_w =$ “On input x

Explanation: $\langle M, w \rangle \in \overline{A_{\text{TM}}}$ iff $\langle M_w \rangle \in E_{\text{TM}}$

M rejects w iff $L(\langle M_w \rangle) = \emptyset$

1. If $x \neq w$, *reject*.
2. else run M on w
3. *Accept* if M accepts.”



EQ_{TM} and $\overline{EQ_{TM}}$ are T-unrecognizable

$$EQ_{TM} = \{ \langle M_1, M_2 \rangle \mid M_1 \text{ and } M_2 \text{ are TMs and } L(M_1) = L(M_2) \}$$

Theorem: Both EQ_{TM} and $\overline{EQ_{TM}}$ are T-unrecognizable

Proof: (1) $\overline{A_{TM}} \leq_m EQ_{TM}$
(2) $A_{TM} \leq_m \overline{EQ_{TM}}$

For any w let $T_w =$ “On input x T_w acts on all inputs the way M acts on w .

1. Ignore x .
2. Simulate M on w .”

(1) Here we give f which maps $\overline{A_{TM}}$ problems (of the form $\langle M, w \rangle$) to EQ_{TM} problems (of the form $\langle T_1, T_2 \rangle$).

$$f(\langle M, w \rangle) = \langle T_w, T_{\text{reject}} \rangle \quad T_{\text{reject}} \text{ is a TM that always rejects.}$$

(2) Similarly $f(\langle M, w \rangle) = \langle T_w, T_{\text{accept}} \rangle \quad T_{\text{accept}}$ always accepts.

Reducibility terminology

Why do we use the term “reduce”?

When we reduce A to B , we show how to solve A by using B and conclude that A is no harder than B . (suggests the \leq_m notation)

Possibility 1: We bring A 's difficulty down to B 's difficulty.

Possibility 2: We bring B 's difficulty up to A 's difficulty.

Quick review of today

1. Introduced The Reducibility Method to prove undecidability and T-unrecognizability.
2. Defined mapping reducibility as a type of reducibility.
3. E_{TM} is undecidable.
4. E_{TM} is T-unrecognizable.
5. EQ_{TM} and $\overline{EQ_{TM}}$ are T-unrecognizable.