

Section (2)

Physics (I)

Revision question

Suppose $A = BC$, where A has the dimension L/M and C has the dimension L/T . Then B has the dimension:

- A. T/M
- B. L^2/TM
- C. TM/L^2
- D. L^2T/M
- E. M/L^2T

Ans: A

- 1) The speed-skating record for women is held by Bonnie Blair, who completed a 500-m race in 39.10 s. What is the corresponding average speed? Give the answer in m/s and in km/h.

Solution:

$$\text{Average speed} = \frac{\text{total distance}}{\text{time taken}} = \frac{500}{39.1} = 12.8 \frac{\text{m}}{\text{s}} = 12.8 \times \frac{3600}{1000} = 46.0 \frac{\text{km}}{\text{h}}$$

2) A particle moves along the x axis, with the following equation for the position as a function of time:

$$x = 2.0 + 6.0t - 3.0t^2$$

where x is measured in meters and t is measured in seconds.

- (a) What is the position of the particle at $t = 0.50$ s?
- (b) What is the instantaneous velocity at this time?
- (c) What is the instantaneous acceleration at this time?

Solution:

(a) The position at $t = 0.5$ s is: $x = 2.0 + 6.0 \times 0.5 - 3.0 \times 0.5^2 = 4.3$ m

(b) The instantaneous velocity at $t = 0.5$ s is:

$$v = \frac{dx}{dt} = 6.0 - 6.0t = 6 - 6 \times 0.5 = 3.00 \text{ m/s}$$

(c) The instantaneous acceleration at $t = 0.5$ s is:

$$a = \frac{dv}{dt} = -6.0 \text{ m/s}^2$$

101. A particle is initially at rest. Beginning at $t = 0$, it begins moving, with an acceleration given by $a = a_o \{1 - [t^2/(4.0s^2)]\}$ for $0 \leq t \leq 2 \text{ s}$ and $a = 0$ thereafter. The initial value is $a_o = 20 \text{ m/s}^2$. What is the particle's velocity after 1.0 s? After a long time? How far has the particle traveled after 2.0 s?

Answer:

$$v = \int a \, dt = \int a_o \left\{ 1 - \left[\frac{t^2}{(4.0s^2)} \right] \right\} dt$$

$$v = a_o \left\{ t - \left[\frac{t^3}{(12s^2)} \right] \right\}$$

After 1.0 s:

$$v = 20 \left\{ 1 - \left[\frac{1}{(12)} \right] \right\} = 18.3 \frac{m}{s}$$

$$v = 20 \left\{ 2 - \left[\frac{8}{(12)} \right] \right\} = 26.7 \text{ m/s}$$

$$x = \int v \, dt = \int a_o \left\{ t - \left[\frac{t^3}{(12s^2)} \right] \right\} dt = a_o \left\{ \frac{t^2}{2} - \left[\frac{t^4}{(48s^2)} \right] \right\}$$

At $t = 2 \text{ s}$:

$$x = 20 \left\{ \frac{2^2}{2} - \left[\frac{2^4}{(48)} \right] \right\} = 33.3 \text{ m}$$

A particle moves along the x axis from x_i to x_f . Of the following values of the initial and final coordinates, which results in the displacement with the largest magnitude?

- A. $x_i = 4 \text{ m}, x_f = 6 \text{ m}$
- B. $x_i = -4 \text{ m}, x_f = -8 \text{ m}$
- C. $x_i = -4 \text{ m}, x_f = 2 \text{ m}$
- D. $x_i = 4 \text{ m}, x_f = -2 \text{ m}$
- E. $x_i = -4 \text{ m}, x_f = 4 \text{ m}$

Ans: E

A particle moves along the x axis from x_i to x_f . Of the following values of the initial and final coordinates, which results in a negative displacement?

- A. $x_i = 4 \text{ m}, x_f = 6 \text{ m}$
- B. $x_i = -4 \text{ m}, x_f = -8 \text{ m}$
- C. $x_i = -4 \text{ m}, x_f = 2 \text{ m}$
- D. $x_i = -4 \text{ m}, x_f = -2 \text{ m}$
- E. $x_i = -4 \text{ m}, x_f = 4 \text{ m}$

Ans: B

Each of four particles move along an x axis. Their coordinates (in meters) as functions of time (in seconds) are given by

particle 1: $x(t) = 3.5 - 2.7t^3$

particle 2: $x(t) = 3.5 + 2.7t^3$

particle 3: $x(t) = 3.5 + 2.7t^2$

particle 4: $x(t) = 3.5 - 3.4t - 2.7t^2$

Which of these particles have constant acceleration?

- A. All four
- B. Only 1 and 2
- C. Only 2 and 3
- D. Only 3 and 4
- E. None of them

Ans: D