



Physics (2)


Section 08

Faculty of Information Technology
Egyptian E-Learning University

Spring 2021



Electric current (I)



We can assume that all electrons travel with constant drift velocity

- The current is defined as the net charge (dq) that passes (perpendicular) through a given area per unit time.

$$I = \frac{dq}{dt}$$

Scalar quantity

$$[I] = \frac{[q]}{[t]} = \frac{C}{Sec} = A$$

Materials that obeys Ohm's law are called Ohmic. Example : Metals

$$R = \frac{\Delta V}{I} : \text{Ohm's Law}$$

$$I = \frac{1}{R} \Delta V$$

Conductivity (σ)

$$R = \frac{\ell}{\sigma A}$$

$$\sigma = \frac{1}{R} \frac{\ell}{A}$$

Units of $\sigma = \Omega^{-1} \cdot m^{-1}$

Resistivity (ρ)

$$\rho = \frac{1}{\sigma}$$

$$\rho = R \frac{A}{\ell}$$

$$R = \rho \frac{\ell}{A}$$

Units of $\rho = \Omega \cdot m$

The power P is defined as the rate at which the charge loses energy

$$P = \frac{\Delta U}{\Delta t}$$

$$P = I \Delta V = \frac{\Delta V^2}{R} = I^2 R$$

Q1. In the Bohr model of the hydrogen atom, an electron in the lowest energy state moves at a speed of 2.19×10^6 m/s in a circular path of radius 5.29×10^{-11} m. What is the effective current associated with this orbiting electron?

Solution

The period of the electron in its orbit is $T = 2\pi r / v$, and the current represented by the orbiting electron is

$$\begin{aligned} I &= \frac{\Delta Q}{\Delta t} = \frac{|e|}{T} = \frac{v|e|}{2\pi r} \\ &= \frac{(2.19 \times 10^6 \text{ m/s})(1.60 \times 10^{-19} \text{ C})}{2\pi (5.29 \times 10^{-11} \text{ m})} \\ &= 1.05 \times 10^{-3} \text{ C/s} = \boxed{1.05 \text{ mA}} \end{aligned}$$

Q2. A 0.900 V potential difference is maintained across a 1.50 m length of tungsten wire that has a cross sectional area of 0.600 mm². What is the current in the wire? Note: $\rho_{tungsten} = 5.60 \times 10^{-8} \Omega \cdot m$

Solution

$\Delta V = IR$ and $R = \frac{\rho \ell}{A}$. The area is

$$A = (0.600 \text{ mm}^2) \left(\frac{1.00 \text{ m}}{1\,000 \text{ mm}} \right)^2 = 6.00 \times 10^{-7} \text{ m}^2$$

From the potential difference, we can solve for the current, which gives

$$\Delta V = \frac{I \rho \ell}{A} \rightarrow I = \frac{\Delta V A}{\rho \ell} = \frac{(0.900 \text{ V})(6.00 \times 10^{-7} \text{ m}^2)}{(5.60 \times 10^{-8} \Omega \cdot m)(1.50 \text{ m})}$$

$$I = \boxed{6.43 \text{ A}}$$

Q3. A 100 W lightbulb connected to a 120 V source experiences a voltage surge that produces 140 V for a moment. By what percentage does its power output increase? Assume its resistance does not change.

Solution

From $P = (\Delta V)^2 / R$, we find that

$$R = \frac{(\Delta V_i)^2}{P} = \frac{(120 \text{ V})^2}{100 \text{ W}} = 144 \Omega$$

The final current is

$$I_f = \frac{\Delta V_f}{R} = \frac{140 \text{ V}}{144 \Omega} = 0.972 \text{ A}$$

The power during the surge is

$$P = \frac{(\Delta V_f)^2}{R} = \frac{(140 \text{ V})^2}{144 \Omega} = \boxed{136 \text{ W}}$$

So the percentage increase is

$$\frac{136 \text{ W} - 100 \text{ W}}{100 \text{ W}} = 0.361 = \boxed{36.1\%}$$

Q4. Batteries are rated in terms of ampere-hours (A . h). For example, a battery that can produce a current of 2.00 A for 3.00 h is rated at 6.00 A . h. (a) What is the total energy, in kilowatt-hours, stored in a 12.0 V battery rated at 55.0 A . h? (b) At \$0.110 per kilowatt-hour, what is the value of the electricity at dollar that produced by this battery?

Solution

(a) The total energy stored in the battery is

$$\begin{aligned}\Delta U_E &= q(\Delta V) = It(\Delta V) \\ &= (55.0 \text{ A} \cdot \text{h})(12.0 \text{ V}) \left(\frac{1 \text{ C}}{1 \text{ A} \cdot \text{s}} \right) \left(\frac{1 \text{ J}}{1 \text{ V} \cdot \text{C}} \right) \left(\frac{1 \text{ W} \cdot \text{s}}{1 \text{ J}} \right) \\ &= 660 \text{ W} \cdot \text{h} = \boxed{0.660 \text{ kWh}}\end{aligned}$$

(b) The value of the electricity is

$$\text{Cost} = (0.660 \text{ kWh}) \left(\frac{\$0.110}{1 \text{ kWh}} \right) = \boxed{\$0.0726}$$

MSQ

(1)

How much energy is dissipated as heat during a two-minute time interval by a 1.5-k Ω resistor which has a constant 20-V potential difference across its leads?

- a. 58 J
- b. 46 J
- c. 32 J
- d. 72 J
- e. 16 J

Answer (C)

(2)

How many electrons pass through a 20- Ω resistor in 10 min if there is a potential drop of 30 volts across it?

- a. 5.6×10^{21}
- b. 7.5×10^{21}
- c. 9.4×10^{21}
- d. 1.1×10^{21}
- e. 3.8×10^{21}

Answer (a)

(3)

A cook plugs a 500 W crockpot and a 1000 W kettle into a 240 V power supply, all operating on direct current. When we compare the two, we find that

- a. $I_{\text{crockpot}} < I_{\text{kettle}}$ and $R_{\text{crockpot}} < R_{\text{kettle}}$.
- b. $I_{\text{crockpot}} < I_{\text{kettle}}$ and $R_{\text{crockpot}} > R_{\text{kettle}}$.
- c. $I_{\text{crockpot}} = I_{\text{kettle}}$ and $R_{\text{crockpot}} = R_{\text{kettle}}$.
- d. $I_{\text{crockpot}} > I_{\text{kettle}}$ and $R_{\text{crockpot}} < R_{\text{kettle}}$.
- e. $I_{\text{crockpot}} > I_{\text{kettle}}$ and $R_{\text{crockpot}} > R_{\text{kettle}}$.

Answer (b)

(4)

If 5.0×10^{21} electrons pass through a $20\text{-}\Omega$ resistor in 10 min, what is the potential difference across the resistor?

- a. 21 V
- b. 32 V
- c. 27 V
- d. 37 V
- e. 54 V

Answer (c)