

Fitting a generalized Pareto distribution to burned area data

Anas Mourahib


December 16, 2024

August complex wildfire

- List of California wildfires from 2003 to 2024


August complex wildfire

- List of California wildfires from 2003 to 2024
- The August Complex (August-November 2020, Northern California) burned more than 1 million acres



August complex wildfire


- List of California wildfires from 2003 to 2024
- The August Complex (August-November 2020, Northern California) burned more than 1 million acres



- **Question** : How unusual is the August Complex wildfire?

August complex wildfire


- List of California wildfires from 2003 to 2024
- The August Complex (August-November 2020, Northern California) burned more than 1 million acres



- **Question** : How unusual is the August Complex wildfire?
- One can say that such an event can happen once every 20 years...

August complex wildfire

- List of California wildfires from 2003 to 2024
- The August Complex (August-November 2020, Northern California) burned more than 1 million acres



- **Question** : How unusual is the August Complex wildfire?
- One can say that such an event can happen once every 20 years...
- however this estimation is based on small sample of observations


Peaks over thresholds

- Let X_1, \dots, X_n denote the number of acres burned on California wildfires from 2003 to 2024
- We use the following model

$$\lim_{u \rightarrow \infty} P(X - u \leq x \mid X > u) = G(x) \quad (1)$$

→ Condition $X > u$: “peaks over thresholds”

→ $Y = X - u$ is the exceedance loss above the high threshold u



Generalized Pareto distribution

- Question: which kind of distribution G can appear in Equation (1)?

Generalized Pareto distribution

- **Question:** which kind of distribution G can appear in Equation (1)?

Theorem (Gnedenko (1943); Fisher and Tippett (1928))

*The limit distribution G in (1) is necessarily a **generalized Pareto distribution** (gpd):*

$$G_{\gamma, \alpha}(x) = \begin{cases} 1 - (1 + \gamma x / \alpha)^{-1/\gamma}, & \gamma \neq 0, \\ 1 - e^{-x/\alpha}, & \gamma = 0, \end{cases}$$

where $\alpha > 0$, and $1 + \gamma x / \alpha > 0$ when $\gamma \neq 0$.

Fitting a gpd to exceedances

- Recall X_1, \dots, X_n the number of acres burned on California wildfires from 2003 to 2024

Fitting a gpd to exceedances

- Recall X_1, \dots, X_n the number of acres burned on California wildfires from 2003 to 2024
- A threshold u , typically chosen as the 0.9 quantile in extreme value theory

Fitting a gpd to exceedances

- Recall X_1, \dots, X_n the number of acres burned on California wildfires from 2003 to 2024
- A threshold u , typically chosen as the 0.9 quantile in extreme value theory
- A random number N_u will exceed the threshold u

Fitting a gpd to exceedances

- Recall X_1, \dots, X_n the number of acres burned on California wildfires from 2003 to 2024
- A threshold u , typically chosen as the 0.9 quantile in extreme value theory
- A random number N_u will exceed the threshold u
- Using Equation (1), we assume that the exceedance losses $Y_1, \dots, Y_n \sim G_{\gamma, \alpha}$, and estimate γ and α using maximum likelihood (Bücher and Segers, 2017)

Fitting a gpd to exceedances

- Recall X_1, \dots, X_n the number of acres burned on California wildfires from 2003 to 2024
- A threshold u , typically chosen as the 0.9 quantile in extreme value theory
- A random number N_u will exceed the threshold u
- Using Equation (1), we assume that the exceedance losses $Y_1, \dots, Y_n \sim G_{\gamma, \alpha}$, and estimate γ and α using maximum likelihood (Bücher and Segers, 2017)
- We obtain the log-likelihood function

$$l((\gamma, \alpha); Y_1, \dots, Y_{N_u}) = \begin{cases} -N_u \ln(\alpha) - (1 + 1/\gamma) \sum_{i=1}^{N_u} \ln(1 + \gamma Y_i / \alpha), & \gamma \neq 0, \\ \alpha^{-1} \left(N_u(1 - u) - \sum_{i=1}^{N_u} Y_i \right), & \gamma = 0, \end{cases} \quad (2)$$

which we maximize subject to $\alpha > 0$, and $1 + \gamma Y_i / \alpha > 0$ for all i , when $\gamma \neq 0$

QQ-plot of the fitting: Part 1

- We minimize the minus likelihood function in (2) using the R function `'optim()'` from the `'stats'` package

QQ-plot of the fitting: Part 1

- We minimize the minus likelihood function in (2) using the R function 'optim()' from the 'stats' package
- We obtain $\hat{\gamma} \approx 0.419$ and $\hat{\alpha} \approx 62\,886$

QQ-plot of the fitting: Part 1

- We minimize the minus likelihood function in (2) using the R function 'optim()' from the 'stats' package
- We obtain $\hat{\gamma} \approx 0.419$ and $\hat{\alpha} \approx 62\,886$
- The quantile function of a gpd is given by

$$Q_{\gamma,\alpha}(p) = \frac{\gamma}{\alpha} \left((1-p)^{-\gamma} - 1 \right) \quad (3)$$

QQ-plot of the fitting: Part 1

- We minimize the minus likelihood function in (2) using the R function 'optim()' from the 'stats' package
- We obtain $\hat{\gamma} \approx 0.419$ and $\hat{\alpha} \approx 62\,886$
- The quantile function of a gpd is given by

$$Q_{\gamma,\alpha}(p) = \frac{\gamma}{\alpha} \left((1-p)^{-\gamma} - 1 \right) \quad (3)$$


- We consider the QQ-plot

$$\left(Q_{\hat{\gamma},\hat{\alpha}}\left(\frac{i}{N_u+1}\right), Y_{i,N_u} \right), \quad i = \dots, N_u,$$

where Y_{i,N_u} denote the i -th ranked observation

- See the plot on the next slide

QQ-plot of the fitting: Part 2



How unusual is the August Complex wildfire?

- Again, we assume that the exceedance losses $Y_1, \dots, Y_{N_u} \sim G_{\hat{\gamma}, \hat{\alpha}}$
- Let $x > u$. In our example, x is the number of acres burned in the August Complex wildfire and recall u , the 0.9 quantile

$$\begin{aligned} P(X > x) &= P(X > u) P(X > x \mid X > u) \\ &= 0.9 \times P(X - u > x - u \mid X > u) \\ &= 0.9 \times P(Y > x - u \mid X > u) \\ &= 0.9 \times \left(1 + \hat{\gamma} \frac{x - u}{\hat{\alpha}}\right)^{-1/\hat{\gamma}} \approx 0.006648479 \end{aligned}$$

Bibliography

- Bücher, A. and J. Segers (2017). On the maximum likelihood estimator for the generalized extreme-value distribution. *Extremes* 20, 839–872.
- Fisher, R. A. and L. H. C. Tippett (1928). Limiting forms of the frequency distribution of the largest or smallest member of a sample. In *Mathematical proceedings of the Cambridge philosophical society*, Volume 24, pp. 180–190. Cambridge University Press.
- Gnedenko, B. (1943). Sur la distribution limite du terme maximum d'une serie aleatoire. *Annals of mathematics* 44(3), 423–453.