

# Topoi with enough points

Ivan Di Liberti

Logic Colloquium 24  
June 2024, Gothenburg.



This talk is based on a joint work with *Morgan Rogers*.

- **Topoi with enough points**, ArXiv:2403.15338.



This talk is based on a joint work with *Morgan Rogers*.

- **Topoi with enough points**, ArXiv:2403.15338.

## Plan



This talk is based on a joint work with *Morgan Rogers*.

- **Topoi with enough points**, ArXiv:2403.15338.

## Plan

- 1 Logical aspects of duality theory,



This talk is based on a joint work with *Morgan Rogers*.

- **Topoi with enough points**, ArXiv:2403.15338.

## Plan

- 1 Logical aspects of duality theory,
- 2 Classifying topoi,



This talk is based on a joint work with *Morgan Rogers*.

- **Topoi with enough points**, ArXiv:2403.15338.

## Plan

- 1 Logical aspects of duality theory,
- 2 Classifying topoi,
- 3 Topoi with enough points



This talk is based on a joint work with *Morgan Rogers*.

- **Topoi with enough points**, ArXiv:2403.15338.

## Plan

- 1 Logical aspects of duality theory,
- 2 Classifying topoi,
- 3 Topoi with enough points (new stuff),



This talk is based on a joint work with *Morgan Rogers*.

- **Topoi with enough points**, ArXiv:2403.15338.

## Plan

- 1 Logical aspects of duality theory,
- 2 Classifying topoi,
- 3 Topoi with enough points (new stuff),





## Boolean algebras



## Boolean algebras

Boole introduced boolean algebras in 1847,



## Boolean algebras

Boole introduced boolean algebras in 1847, starting a program that nowadays we may call *algebraization of logic*.



## Boolean algebras

Boole introduced boolean algebras in 1847, starting a program that nowadays we may call *algebraization of logic*.

$$\mathbb{B} = (B, \top, \perp, \neg, \vee, \wedge).$$



## Boolean algebras

Boole introduced boolean algebras in 1847, starting a program that nowadays we may call *algebraization of logic*.

$$\mathbb{B} = (B, \top, \perp, \neg, \vee, \wedge).$$

Motto:



## Boolean algebras

Boole introduced boolean algebras in 1847, starting a program that nowadays we may call *algebraization of logic*.

$$\mathbb{B} = (B, \top, \perp, \neg, \vee, \wedge).$$

Motto:propositional first order theories  $\equiv$  Boolean algebras



## Models vs homomorphisms



## Models vs homomorphisms

Models of a propositional theory (boolean algebras) are assignments of truth values to its formulas,





## Models vs homomorphisms

Models of a propositional theory (boolean algebras) are assignments of truth values to its formulas, and can be understood as homomorphisms into  $\mathbb{T} = \{0 < 1\}$ .

$$M : \mathbb{B} \rightarrow \mathbb{T}$$



## Models vs homomorphisms

Models of a propositional theory (boolean algebras) are assignments of truth values to its formulas, and can be understood as homomorphisms into  $\mathbb{T} = \{0 < 1\}$ .

$$M : \mathbb{B} \rightarrow \mathbb{T}$$

$$\phi \mapsto [\phi]$$



## Models vs homomorphisms

Models of a propositional theory (boolean algebras) are assignments of truth values to its formulas, and can be understood as homomorphisms into  $\mathbb{T} = \{0 < 1\}$ .

$$M : \mathbb{B} \rightarrow \mathbb{T}$$

$$\phi \mapsto [\phi]$$



## Completeness theorem



## Completeness theorem

**Question:** can we recover a boolean algebra (a propositional theory) from its models?



## Completeness theorem

**Question:** can we recover a boolean algebra (a propositional theory) from its models?

**Yes, we can.**



## Completeness theorem

**Question:** can we recover a boolean algebra (a propositional theory) from its models?

**Yes, we can.** Provability can indeed be tested semantically.



## Completeness theorem

**Question:** can we recover a boolean algebra (a propositional theory) from its models?

**Yes, we can.** Provability can indeed be tested semantically.  
(Completeness theorem.)





## Completeness theorem

**Question:** can we recover a boolean algebra (a propositional theory) from its models?

**Yes, we can.** Provability can indeed be tested semantically.  
(Completeness theorem.)

## Stone duality



## Completeness theorem

**Question:** can we recover a boolean algebra (a propositional theory) from its models?

**Yes, we can.** Provability can indeed be tested semantically.  
(Completeness theorem.)

## Stone duality

There is a dual adjunction between boolean algebras (theories) and stone spaces



## Completeness theorem

**Question:** can we recover a boolean algebra (a propositional theory) from its models?

**Yes, we can.** Provability can indeed be tested semantically.  
(Completeness theorem.)

## Stone duality

There is a dual adjunction between boolean algebras (theories) and stone spaces (sets of models *equipped with a topology*).




## Completeness theorem

**Question:** can we recover a boolean algebra (a propositional theory) from its models?

**Yes, we can.** Provability can indeed be tested semantically.  
(Completeness theorem.)

## Stone duality

There is a dual adjunction between boolean algebras (theories) and stone spaces (sets of models *equipped with a topology*).



## Stone-type dualities



## Stone-type dualities

- Stone-type dualities is a cluster of theorems that relates classes of posets (propositional logics) to classes of spaces (their models)



## Stone-type dualities

- Stone-type dualities is a cluster of theorems that relates classes of posets (propositional logics) to classes of spaces (their models) .
- One of the aims is to reconstruct the theory from its models (completeness theorem)



## Stone-type dualities

- Stone-type dualities is a cluster of theorems that relates classes of posets (propositional logics) to classes of spaces (their models) .
- One of the aims is to reconstruct the theory from its models (completeness theorem) .
- Boolean algebras are the posets of propositional first order logic.





## Stone-type dualities

- Stone-type dualities is a cluster of theorems that relates classes of posets (propositional logics) to classes of spaces (their models) .
- One of the aims is to reconstruct the theory from its models (completeness theorem) .
- Boolean algebras are the posets of propositional first order logic.

## Isbell duality



## Stone-type dualities

- Stone-type dualities is a cluster of theorems that relates classes of posets (propositional logics) to classes of spaces (their models) .
- One of the aims is to reconstruct the theory from its models (completeness theorem) .
- Boolean algebras are the posets of propositional first order logic.

## Isbell duality

There is a very general duality in which every other embeds,




## Stone-type dualities

- Stone-type dualities is a cluster of theorems that relates classes of posets (propositional logics) to classes of spaces (their models) .
- One of the aims is to reconstruct the theory from its models (completeness theorem) .
- Boolean algebras are the posets of propositional first order logic.

## Isbell duality

There is a very general duality in which every other embeds, the duality between frames and topological spaces.



## Frames



## Frames

A frame is a complete lattice where the *infinitary distributivity rule* holds,

$$(\bigvee x_i) \wedge y = \bigvee (x_i \wedge y).$$

A frame homomorphism is a morphism  $f$  preserving suprema ( $\bigvee$ ) and finite meets ( $\wedge$ ).



## Frames

A frame is a complete lattice where the *infinitary distributivity rule* holds,

$$(\bigvee x_i) \wedge y = \bigvee (x_i \wedge y).$$

A frame homomorphism is a morphism  $f$  preserving suprema ( $\bigvee$ ) and finite meets ( $\wedge$ ).

## Topologies are frames



## Frames

A frame is a complete lattice where the *infinitary distributivity rule* holds,

$$(\bigvee x_i) \wedge y = \bigvee (x_i \wedge y).$$

A frame homomorphism is a morphism  $f$  preserving suprema ( $\bigvee$ ) and finite meets ( $\wedge$ ).

## Topologies are frames

The canonical example of frame is a topology. When  $(X, \tau)$  is a topological space,  $\tau$  is a frame.

$$\mathcal{O} : \mathbf{Top} \rightarrow \mathbf{Frm}^{\text{op}}$$



## Models or Formal points





## Models or Formal points

For  $\mathcal{L}$  a frame we can study its models, as we did for boolean algebras,

$$M : \mathcal{L} \rightarrow \mathbb{T}$$



## Models or Formal points

For  $\mathcal{L}$  a frame we can study its models, as we did for boolean algebras,

$$M : \mathcal{L} \rightarrow \mathbb{T}$$

this will be a frame homomorphism into truth values. The set of models can be topologised, providing a dual adjunction.

## Isbell duality




## Models or Formal points

For  $\mathcal{L}$  a frame we can study its models, as we did for boolean algebras,

$$M : \mathcal{L} \rightarrow \mathbb{T}$$

this will be a frame homomorphism into truth values. The set of models can be topologised, providing a dual adjunction.

## Isbell duality



## Frames and logic



## Frames and logic

Boolean algebras are to first order logic what frames are to *geometric logic*, a logic of infinitary disjunction, finitary conjunction and infinitary distributivity rule.



## Frames and logic

Boolean algebras are to first order logic what frames are to *geometric logic*, a logic of infinitary disjunction, finitary conjunction and infinitary distributivity rule.



## Frames and logic

Boolean algebras are to first order logic what frames are to *geometric logic*, a logic of infinitary disjunction, finitary conjunction and infinitary distributivity rule.

**Question:** Can we recover a frame from its space of models?



## Frames and logic

Boolean algebras are to first order logic what frames are to *geometric logic*, a logic of infinitary disjunction, finitary conjunction and infinitary distributivity rule.

**Question:** Can we recover a frame from its space of models?  
**Not always** this adjunction is too general to hope for such a result. But...





## Frames and logic

Boolean algebras are to first order logic what frames are to *geometric logic*, a logic of infinitary disjunction, finitary conjunction and infinitary distributivity rule.

**Question:** Can we recover a frame from its space of models?  
**Not always** this adjunction is too general to hope for such a result. But...

**Definition:** frames with enough points



## Frames and logic

Boolean algebras are to first order logic what frames are to *geometric logic*, a logic of infinitary disjunction, finitary conjunction and infinitary distributivity rule.

**Question:** Can we recover a frame from its space of models?  
**Not always** this adjunction is too general to hope for such a result. But...

### Definition: frames with enough points

A frame has enough points when given two (different) formulas in it  $\phi < \psi$  there exists a model  $M$  such that  $M(\phi) = 0$  and  $M(\psi) = 1$ .



## Frames and logic

Boolean algebras are to first order logic what frames are to *geometric logic*, a logic of infinitary disjunction, finitary conjunction and infinitary distributivity rule.

**Question:** Can we recover a frame from its space of models?  
**Not always** this adjunction is too general to hope for such a result. But...

### Definition: frames with enough points

A frame has enough points when given two (different) formulas in it  $\phi < \psi$  there exists a model  $M$  such that  $M(\phi) = 0$  and  $M(\psi) = 1$ .

### Thm



## Frames and logic

Boolean algebras are to first order logic what frames are to *geometric logic*, a logic of infinitary disjunction, finitary conjunction and infinitary distributivity rule.

**Question:** Can we recover a frame from its space of models?  
**Not always** this adjunction is too general to hope for such a result. But...

### Definition: frames with enough points

A frame has enough points when given two (different) formulas in it  $\phi < \psi$  there exists a model  $M$  such that  $M(\phi) = 0$  and  $M(\psi) = 1$ .

### Thm

When a frame has enough points, then  $\text{Opt}(\mathcal{L}) \cong \mathcal{L}$ .



## From propositional to first order logic



## From propositional to first order logic

Posets are perfect to handle *propositional* logic.



## From propositional to first order logic

Posets are perfect to handle *propositional* logic.

**Question:** What algebraic gadget can we use to handle *first order logic*?



## From propositional to first order logic

Posets are perfect to handle *propositional* logic.

**Question:** What algebraic gadget can we use to handle *first order logic*?

**Question:** Can we recover a Stone-type duality for first order logic?





## From propositional to first order logic

Posets are perfect to handle *propositional* logic.

**Question:** What algebraic gadget can we use to handle *first order logic*?

**Question:** Can we recover a Stone-type duality for first order logic?

## Classifying topoi



## From propositional to first order logic

Posets are perfect to handle *propositional* logic.

**Question:** What algebraic gadget can we use to handle *first order logic*?

**Question:** Can we recover a Stone-type duality for first order logic?

## Classifying topoi

Yes.



## From propositional to first order logic

Posets are perfect to handle *propositional* logic.

**Question:** What algebraic gadget can we use to handle *first order logic*?

**Question:** Can we recover a Stone-type duality for first order logic?

## Classifying topoi

**Yes.** The previous picture can be recovered by replacing frames with **topoi**.



## From propositional to first order logic

Posets are perfect to handle *propositional* logic.

**Question:** What algebraic gadget can we use to handle *first order logic*?

**Question:** Can we recover a Stone-type duality for first order logic?

## Classifying topoi

**Yes.** The previous picture can be recovered by replacing frames with **topoi**. A topos is a category (as opposed to poset) with all colimits (suprema), finite limits (wedges) and a good interaction between them (infinitary distributivity rule).



## From propositional to first order logic

Posets are perfect to handle *propositional* logic.

**Question:** What algebraic gadget can we use to handle *first order logic*?

**Question:** Can we recover a Stone-type duality for first order logic?

## Classifying topoi

**Yes.** The previous picture can be recovered by replacing frames with **topoi**. A topos is a category (as opposed to poset) with all colimits (suprema), finite limits (wedges) and a good interaction between them (infinitary distributivity rule).

## DL 2020, a Stone-type duality



## From propositional to first order logic

Posets are perfect to handle *propositional* logic.


**Question:** What algebraic gadget can we use to handle *first order* logic?

**Question:** Can we recover a Stone-type duality for first order logic?

## Classifying topoi

**Yes.** The previous picture can be recovered by replacing frames with **topoi**. A topos is a category (as opposed to poset) with all colimits (suprema), finite limits (wedges) and a good interaction between them (infinitary distributivity rule).

## DL 2020, a Stone-type duality



## Definition: topoi with enough points



**Definition: topoi with enough points**

A topos  $\mathcal{E}$  has enough points when given a proper subobject  $f : X \hookrightarrow Y$  there exists a model  $M : \mathcal{E} \rightarrow \mathbf{Set}$  such that  $M(f)$  is not an isomorphism.





**Definition: topoi with enough points**

A topos  $\mathcal{E}$  has enough points when given a proper subobject  $f : X \hookrightarrow Y$  there exists a model  $M : \mathcal{E} \rightarrow \mathbf{Set}$  such that  $M(f)$  is not an isomorphism.

**Deligne (1970s), Godel completeness theorem**

**Definition: topoi with enough points**

A topos  $\mathcal{E}$  has enough points when given a proper subobject  $f : X \hookrightarrow Y$  there exists a model  $M : \mathcal{E} \rightarrow \mathbf{Set}$  such that  $M(f)$  is not an isomorphism.

**Deligne (1970s), Godel completeness theorem**

Coherent topoi have enough points.



**Definition: topoi with enough points**

A topos  $\mathcal{E}$  has enough points when given a proper subobject  $f : X \hookrightarrow Y$  there exists a model  $M : \mathcal{E} \rightarrow \mathbf{Set}$  such that  $M(f)$  is not an isomorphism.

**Deligne (1970s), Godel completeness theorem**

Coherent topoi have enough points.

**Makkai and Reyes (1980s), Fourman-Grayson completeness theorem**

**Definition: topoi with enough points**

A topos  $\mathcal{E}$  has enough points when given a proper subobject  $f : X \hookrightarrow Y$  there exists a model  $M : \mathcal{E} \rightarrow \mathbf{Set}$  such that  $M(f)$  is not an isomorphism.

**Deligne (1970s), Godel completeness theorem**

Coherent topoi have enough points.

**Makkai and Reyes (1980s), Fourman-Grayson completeness theorem**

Separable topoi have enough points.



**Definition: topoi with enough points**

A topos  $\mathcal{E}$  has enough points when given a proper subobject  $f : X \hookrightarrow Y$  there exists a model  $M : \mathcal{E} \rightarrow \mathbf{Set}$  such that  $M(f)$  is not an isomorphism.

**Deligne (1970s), Godel completeness theorem**

Coherent topoi have enough points.

**Makkai and Reyes (1980s), Fourman-Grayson completeness theorem**

Separable topoi have enough points.

**Question:** Are these the same theorem?



**Definition: topoi with enough points**

A topos  $\mathcal{E}$  has enough points when given a proper subobject  $f : X \hookrightarrow Y$  there exists a model  $M : \mathcal{E} \rightarrow \mathbf{Set}$  such that  $M(f)$  is not an isomorphism.

**Deligne (1970s), Godel completeness theorem**

Coherent topoi have enough points.

**Makkai and Reyes (1980s), Fourman-Grayson completeness theorem**

Separable topoi have enough points.

**Question:** Are these the same theorem?

**Question:** Can we push to the extreme the technology of Deligne and unify every completeness theorem?



**Thm. DL and Rogers.**



**Thm. DL and Rogers.**

**Yes.**





**Thm. DL and Rogers.**

**Yes.** We generalise the technology of Deligne and offer a general theorem that implies both Deligne's and Makkai-Reyes theorem.



**Thm. DL and Rogers.**

**Yes.** We generalise the technology of Deligne and offer a general theorem that implies both Deligne's and Makkai-Reyes theorem.

**Thm. DL and Rogers.**

**Thm. DL and Rogers.**

**Yes.** We generalise the technology of Deligne and offer a general theorem that implies both Deligne's and Makkai-Reyes theorem.

**Thm. DL and Rogers.**

A closed subtopos of a topos with enough points has enough points.



**Thm. DL and Rogers.**

**Yes.** We generalise the technology of Deligne and offer a general theorem that implies both Deligne's and Makkai-Reyes theorem.

**Thm. DL and Rogers.**

A closed subtopos of a topos with enough points has enough points.

